

- Scott, B.H., 767  
 Screation, N.J., 668  
 Sear, J.W., 281  
 Seppälä, T., 748  
 Settergren, G., 64  
 Shah, A.S., 24S  
 Shams, M., 313  
 Shanewise, J.S., 657, 715  
 Sharma, J., 304  
 Sharma, S., 759, 769  
 Sharma, V.K., 759  
 Shayan, C., 68  
 Sheikh F., 824  
 Shime, N., 218  
 Shimode, Y., 479  
 Shivananda, N., 392, 673  
 Shorten, G., 592  
 Sidebotham, D., 671  
 Sidery, M., 668  
 Sie, H.T., 309  
 Silvey, G., 248, 390, 535, 678, 817  
 Singh, A., 43  
 Singh, A.K., 719  
 Singh, S., 769  
 Sinha, A., 396  
 Sinha, P.K., 123, 469, 632  
 Sisillo, E., 256  
 Skidmore, K.L., 780  
 Sleilaty, G.S., 59  
 Slooff, M.J.H., 31S  
 Smith, B.E., 620  
 Smith, P.K., 24S  
 Smyrmiotis, V., 30  
 Sørensen, L.H., 709  
 Spahr, T., 587  
 Spain, D., 475  
 Sparicio, D., 704  
 Spiess, B.D., 1S  
 Spinale, F.G., 25  
 Sreedhar, R., 197  
 Sreeram, G.M., 548  
 Stafford-Smith, M., 548  
 Stamm, C., 415  
 Staples, E.D., 563  
 Staples, J.R., 715  
 Steenbeck, J., 529  
 Steinfath, M., 185  
 Story, D.A., 613  
 Stratmann, G., 175  
 Straub, U., 731  
 Stuart-Smith, K., 317, 482  
 Sugahara, K., 806  
 Sugita, M., 479  
 Sukernik, M.R., 101, 824  
 Sullivan, E., 228  
 Sungur, Z., 152  
 Svenarud, P., 180  
 Svendsen, J.H., 709  
 Swafford, J., 396  
 Swaminathan, M., 107, 404  
 Swoboda, H., 336  
 Szokol, J.W., 812  
 Tanaka, K., 288  
 Tanaka, K.A., 715  
 Tarkka, M.R., 799  
 Taylor, K.M., 93  
 Taylor, R., 317  
 Telci, L., 446  
 Tempe, D.K., 657  
 Thackray, M., 540  
 Thakar, D., 396  
 Thangathurai, D., 822  
 Theodoraki, K., 30  
 Thimmannagowda, P., 512  
 Thingnam, S.K., 122  
 Thomas, J.J., 765  
 Thomassen, L., 709  
 Thompson, B., 47  
 Tilak, M.D., 75  
 Tobias, J.D., 207  
 Tokumine, J., 806  
 Tomar, A.S., 657  
 Tonner, P.H., 185  
 Toscana, D., 148  
 Tran, Q.H.D., 788  
 Trehan, N., 304  
 Trivedi, U., 539  
 Troster, E.J., 160  
 Tseng, C.-C.A., 552  
 Tugrul, M., 152, 446  
 Tunaoglu, F.S., 775  
 Tür, A., 461  
 Tuttle-Newhall, E., 438  
 Uckunkaya, N., 755  
 Uezu, T., 806  
 Urdaneta, F., 624  
 Üstün, E., 461  
 Vales, R., Jr, 353  
 van der Linden, J., 180  
 van der Starre, P.J.A., 309, 466  
 van Rooyen-Butijn, M., 309  
 Varma, P.K., 197, 632  
 Varner, J., 475  
 Veeken, C., 506  
 Veit, M., 53  
 Verkkala, K., 748  
 Via, G., 242  
 Vinluan, J., 573  
 Vogt, S., 275  
 Votapka, T.V., 812  
 Vuylsteke, A., 394, 668  
 Wagner, R., 777  
 Waikar, H.D., 332  
 Waldenberger, F., 336  
 Wall, M.H., 238  
 Walton, S., 25  
 Wang, L., 210  
 Wasnick, J.D., 573  
 Waters, P.F., 228  
 Watnik, M., 43  
 Weidemann, F., 415  
 Weightman, W., 136, 540  
 Weiner, J., 725  
 Weiss, S.J., 521, 725  
 Wells, J., 763  
 White, W.D., 548  
 Wilkes, M.M., 429  
 Wilkin, T., 812  
 Willert, J., 624  
 Wilson, W.C., 175  
 Winning, J., 731  
 Wippermann, J., 527  
 Worthmann, F., 185  
 Wu, G.-J., 114  
 Wulf, H., 275  
 Yalcin, O., 166  
 Yamashiro, S., 806  
 Yang, H., 620  
 Yans, T., 637  
 Yasumoto, M., 121  
 Yazigi, A.G., 59  
 Yetkin, U., 339  
 Yey, H.-M., 114  
 Yilik, L., 339  
 Young, N., 43  
 Yppäriälä, H., 559  
 Yuskevich, B., 451  
 Zabeeda, D., 156  
 Zalunardo, M.P., 587  
 Zangrillo, A., 34, 38, 442, 704  
 Zhuravlev, I., 573  
 Ziser, A., 458  
 Zollinger, A., 587  
 Zurakowski, D., 415

## SUBJECT INDEX

- Acinetobacter*, in central venous catheter-related infections, 304–308  
 Activated coagulation time, intraoperative, arterial vs. venous, 573–580  
 Acupressure, wristbands for, for postoperative nausea and vomiting, 68–71  
 Acute respiratory distress syndrome  
 in off-pump CABG surgery, 491  
 Acute respiratory distress syndrome (*Continued*)  
 in severe chest trauma, extracorporeal lung assist for, 777–779  
 Adhesion molecules, after CABG surgery, simvastatin effects on, 605–609  
 Adrenal insufficiency, delayed-onset, postoperative, 80–81  
 Age factors, in atrial fibrillation, after off-pump CABG surgery, 704–708

- Aggregometry, platelet, for platelet dysfunction assessment, 136–140
- Air embolism  
in cardiopulmonary bypass, 112–113  
cerebral, bispectral index value in, 82–84  
prevention of, carbon dioxide deairing in, 180–184
- Airway  
anomalies of, difficult intubation in, 610–612  
obstruction of, after lung resection, from subglottic granuloma, 479–481
- Albumin, for pump priming, vs. crystalloid, 429–437
- Aldosterone, heparin and, 96–97
- Aminocaproic acid, for blood loss prevention, in pediatric patients, 141–143
- Amiodarone  
for atrial fibrillation, after CABG surgery, 709–714  
multisystem organ failure related to, 366–374
- Amrinone, for postoperative care, in pediatric patients, 220
- Analgesia  
postoperative. *See* Postoperative pain  
for video-assisted thoracoscopy, 317–321
- Anaphylaxis, from aprotinin, 39S–40S
- Anemia, postoperative, treatment of, 59–63
- Anesthesia. *See specific agents and procedures*
- Aneurysm, aortic  
perioperative beta-blockade in, 7–13  
stent placement for, transesophageal echocardiography in, 466–468  
transmediastinal repair of, cerebrospinal fluid drainage catheters for, 624–627
- Angioplasty, coronary, brain natriuretic peptide and, 782–783
- Angiotensin-converting enzyme inhibitors, intraoperative, hemodynamic results of, 47–52
- Animal studies  
arrhythmias in magnesium deficiency, 313–316  
cardiac output measurement in one-lung ventilation, 190–193  
left ventricular pressure-area relationship indices, 415–422  
minimum alveolar concentration of isoflurane in cardiopulmonary bypass, 620–623  
vena cava obstruction detection, 293–303
- Antithrombin III, heparin binding to, 93, 131–135
- Aorta  
aneurysm of  
perioperative beta-blockade in, 7–13  
stent placement for, transesophageal echocardiography in, 466–468  
transmediastinal repair of, cerebrospinal fluid drainage catheters for, 624–627  
atherosclerosis of, ultrasonography in, 715–718  
coarctation of  
coronary artery anomalous origin and, 327–331  
jet ventilation during, 207–209  
cross-clamping of, pulmonary artery perfusion in, 166–174  
dissection of, repair of, patent foramen ovale formation in, 763–764  
interruption of, in cardiopulmonary bypass, 469–471  
prosthetic valve dehiscence into, 812–813  
pseudoaneurysm of, echocardiography in, 527–528  
stent placement in, transesophageal echocardiography in, 466–468  
transit-time ultrasound of, in cardiac output measurement, before and after cardiopulmonary bypass, 563–572
- Aortic root, perfusion of, for ventricular arrhythmias, after aortic valve replacement, 197–200
- Aortic valve  
bicuspid, sinus of Valsalva thrombosis with, 384–386  
papillary fibroelastoma of, echocardiography in, 114  
prosthetic, dehiscence of, aortic obstruction by, 812–813
- Aortic valve (*Continued*)  
regurgitation of, brain natriuretic peptide and, 783  
replacement of  
atrial natriuretic peptide levels in, 30–33  
cerebral air embolism in, 82–84  
delayed-onset adrenal insufficiency in, 80–81  
gauze object left during, 242–244  
pulmonary artery catheter venous infusion port interruption in, 72–74  
refractory ventricular arrhythmias after, controlled aortic root perfusion for, 197–200  
stenosis of, brain natriuretic peptide and, 783
- Aprotinin  
for blood conservation, 24S–28S  
benefits of, 34S–36S  
clinical trials of, 24S–26S  
cost-effectiveness of, 26S–27S  
dose for, 25S  
heparin and, 38S–42S  
in liver transplantation, 31S–37S  
mechanisms of action of, 24S  
for orthopedic surgery, 29S–30S, 40S–41S  
safety of, 25S–26S, 33S–34S  
for cardiopulmonary bypass, stroke after, in pediatric patients, 194–196
- Argatroban, for heparin-induced thrombocytopenia, 95
- Arrhythmias  
atrial  
after CABG surgery, 309–312, 704–708, 709–714  
in off-pump CABG surgery, 492–493  
intraoperative, 641–644  
in magnesium deficiency, 313–316  
prevention of, ischemic preconditioning in, 800  
ventricular  
after aortic valve replacement, controlled aortic root perfusion for, 197–200  
after CABG surgery, 366
- Arterial switch operation, postoperative care in, 219–220
- Arteries. *See specific arteries*
- Aspirin, platelet dysfunction due to, assessment of, 136–140
- Atenolol, perioperative, in aortic aneurysm surgery, cost-effectiveness of, 7–13
- Atherosclerosis, aortic, ultrasonography in, 715–718
- Atrial appendage thrombus, dislodgment of, in femoral embolectomy, 809–811
- Atrial fibrillation  
after CABG surgery, amiodarone and epidural analgesia and, 709–714  
during off-pump CABG surgery, 492–493, 704–708
- Atrial natriuretic peptide, measurement of, in valve replacement, 30–33
- Atrial septal defect  
brain natriuretic peptide and, 783  
emphysematous bullae with, simultaneous repair of, 632–636  
vena cava anomalies with, 332–335
- Atrial septectomy, caval inflow occlusion in, in pediatric patients, 144–147
- Atrial tachyarrhythmias, after CABG surgery, magnesium and sotalol effects on, 309–312
- Atrium  
left, pseudoaneurysm behind, 671–672  
myxoma of, cardiac arrest in, resuscitation in, 769–771
- Auditory-evoked potentials, midlatency, in sedation assessment, 559–562

- Autologous blood donation, preoperative, 6S–11S  
 cell-washing systems for, continuous vs. discontinuous, 210–217  
 for orthopedic surgery, 29S
- Awareness, anesthetic, pharmacogenetics and, 361
- Beta-adrenergic receptor, genetic variations in, 359–361
- Beta-blockers, perioperative  
 for aortic aneurysm surgery, cost-effectiveness of, 7–13  
 for noncardiac surgery, physician attitudes toward, 14–24
- Bicaval venous cannulation, obstruction detection in, in pediatric patients, 293–303
- Biomarkers  
 in CABG surgery, 34–37  
 in cerebral function monitoring, in pediatric cardiac surgery, 650–652
- Biopsy, lung, video-assisted thoracoscopy in, intercostal nerve block for, 317–321
- Bisoprolol, perioperative, in aortic aneurysm surgery, cost-effectiveness of, 7–13
- Bispectral index value  
 in cerebral air embolism, 82–84  
 in pediatric cardiac surgery, 647
- Biventricular assist support, for dilated cardiomyopathy, 772–774
- Blalock-Taussig operation, modified, end-tidal carbon dioxide monitoring during, 152–155
- Bleeding, from coronary arteriotomy, in off-pump CABG surgery, 488
- Blood, for autologous use, 8S–10S
- Blood conservation  
 aprotinin in, 24S–28S  
 heparin and, 38S–42S  
 in liver transplantation, 31S–37S  
 in orthopedic surgery, 29S–30S, 40S–41S  
 business plan for, 12S–14S  
 in cardiac surgery, 18S–23S  
 CEO perspective on, 15S–17S  
 cost-effectiveness of, 26S–27S  
 costs of, 22S  
 in liver transplantation, 31S–37S  
 national perspective on, 6S–11S  
 in orthopedic surgery, 29S–30S, 40S–41S  
 reasons for, 1S–5S  
 risk-benefit analysis for, 14S  
 staff acceptance of, 16S–17S
- Blood flow  
 cerebral, in cardiopulmonary bypass, hemodilution and phenylephrine effects on, 423–428  
 mesenteric, after cardiopulmonary bypass, Doppler studies of, 731–733  
 pulmonary, measurement of, indirect Fick method for, 47–52  
 velocity of, in left ventricular pressure-relationship indices measurement, 415–422
- Blood gas, monitoring of, during thoracoscopic surgery, 587–591
- Blood glucose control, in intensive care, 690–697
- Blood loss  
 in off-pump CABG surgery, 492  
 prevention of, epsilon aminocaproic acid vs. tranexamic acid in, in pediatric patients, 141–143
- Blood volume, in left ventricular pressure-relationship indices measurement, 415–422
- Brain  
 air embolism in, bispectral index value in, 82–84  
 blood flow in, in cardiopulmonary bypass, hemodilution and phenylephrine effects on, 423–428  
 functional monitoring of, in pediatric surgery, 645–656
- Brain (*Continued*)  
 ischemia of, in bicaval venous obstruction, detection of, 293–303  
 oxygen saturation in, neuropsychologic dysfunction and, 552–558  
 transfusion effects on, 235
- Brain natriuretic peptide  
 in coronary angioplasty, 782–783  
 in heart failure  
 diastolic, 780–782  
 therapeutic, 784–785  
 in heart transplantation candidates, 782  
 in left ventricular volume overload, 784  
 in right ventricular pressure overload, 783  
 in valve surgery, 783
- Bronchi, rupture of, in endotracheal intubation, 228–233
- Bupivacaine  
 for intercostal nerve block, for video-assisted thoracoscopy, 317–321  
 interpleural, for post-thoracotomy pain, 461–465
- Business plan, for blood conservation, 12S–14S
- CABG surgery. *See* Coronary artery bypass graft (CABG) surgery
- Calcium, ionized, concentration of, coagulation and, 581–586
- Cancer  
 cells of, removal of, from cell-washing autotransfusion systems, 214–215  
 heparin action in, 96
- Candida*, in central venous catheter-related infections, 304–308
- Capillary leak syndrome, postoperative, 222–223
- Carbon dioxide  
 in cardiac output measurement, before and after cardiopulmonary bypass, 563–572  
 for deairing, in air embolism prevention, 180–184  
 end-tidal, monitoring of, in systemic-to-pulmonary artery shunt insertion, 152–155
- Cardiac arrest  
 in left atrial myxoma, resuscitation in, 769–771  
 in prosthetic aortic valve dehiscence, 812–813
- Cardiac output, measurement of  
 in CABG surgery, off-pump, 43–46  
 before and after cardiopulmonary bypass, methods for, 563–572  
 continuous, methods for, 185–189  
 indirect Fick method for, 47–52  
 in one-lung ventilation, 190–193
- Cardiomyopathy, dilated, treatment of, in pediatric patients, 772–774
- Cardioplegia, blood for, salvage application of, in intraoperative ventricular fibrillation, 641–644
- Cardiopulmonary bypass  
 activated coagulation time in, arterial vs. venous, 573–580  
 air embolism in, 112–113  
 aortic interruption during, 469–471  
 aprotinin in, stroke after, in pediatric patients, 194–196  
 atrial natriuretic peptide levels in, 30–33  
 bicaval venous cannulation in, obstruction detection in, 293–303  
 cardiac biomarker release in, 34–37  
 cardiac output measurement before and after, 563–572  
 cerebral blood flow and metabolism in, hemodilution and phenylephrine effects on, 423–428  
 cerebral function monitoring in, 645–656  
 C-reactive protein kinetics after, in pediatric patients, 160–165  
 deep hypothermic circulatory arrest in, femoral-to-radial arterial pressure gradient in, 175–179  
 duration of, kidney failure risk and, 442–445  
 extubation timing after, pain assessment in, 64–67  
 heparin in, resistance development in, 131–135  
 isoflurane in, minimum alveolar concentration and, 620–623

- Cardiopulmonary bypass (*Continued*)
- kidney failure after, 442–445
  - mesenteric blood flow after, Doppler studies of, 731–733
  - for mitral valve replacement, in pregnancy, 339–343
  - multisystem organ failure after, 366–374
  - myocardial protection in. *See* Myocardial protection
  - in narcolepsy, anesthesia safety in, 201–203
  - platelet dysfunction in, heparin and, 93–94
  - procalcitonin kinetics after, in pediatric patients, 160–165
  - pulmonary artery perfusion during, importance of, 166–174
  - pulmonary artery pressures after, milrinone and nitric oxide effects on, in pediatric patients, 156–159
  - pump priming in, albumin vs. crystalloid for, 429–437
  - quinaprilat in, hemodynamic results of, 47–52
  - rewarming garment for, in pediatric patients, 148–151
  - separation from
    - inotrope use in, 401–403, 404–408
    - unsuccessful, 242–244
  - in sickle cell trait, exchange transfusion requirements in, 663–667
  - for simultaneous atrial septal defect and emphysematous bullae repair, 632–636
  - superior vena caval partial occlusion during, 472–474
- CardioThoracic Systems coronary stabilizer, 487
- Carotid artery(ies)
- endarterectomy of
    - myocardial ischemia in, risk factors for, 288–292
    - shunting in, 375–383
  - thrombosis of, after cardiopulmonary bypass, with aprotinin, in pediatric patients, 194–196
- Catheter(s)
- central venous. *See* Central venous catheter
  - cerebrospinal fluid, in transmediastinal repair of aortic aneurysm, 624–627
  - epidural, hematoma from, 759–762
  - multiperforated, for carbon dioxide deairing, 180–184
  - pulmonary artery. *See* Pulmonary artery catheter
- Caval inflow occlusion technique, for congenital heart disease repair, 144–147
- Cavopulmonary anastomosis, superior vena caval partial occlusion during, 472–474
- Cell-washing systems, for autotransfusion, continuous vs. discontinuous, 210–217
- Central venous catheter
- infections due to, prospective study of, 304–308
  - mediastinal misplacement of, 75–77
  - in persistent left superior vena cava, 529–531
  - placement of, measurement differences and, 446–450
- Central venous pressure
- in fluid management, in liver transplantation, 438–441
  - vs. peripheral venous pressure, patient position and, 446–450
- Cervical necrotizing fasciitis, vacuum-assisted closure for, 336–338
- Chest radiography, exterior epidural connector rings showing on, 673–674
- Chest trauma, pumpless extracorporeal lung assist in, 777–779
- Children. *See* Pediatric patients
- Circulating-water rewarming, for pediatric patients, 148–151
- Circulatory arrest, deep hypothermic, femoral-to-radial arterial pressure gradient in, 175–179
- Cisapride, postoperative, for pediatric patients, 222–223
- Coagulation
- intraoperative assessment of, 573–580
  - ionized calcium concentration and, 581–586
  - in off-pump CABG surgery, 492
- Coarctation, aortic
- coronary artery anomalous origin and, 327–331
  - jet ventilation during, 207–209
- Cognitive dysfunction, in off-pump CABG surgery, 489–491
- Cohn Stabilizer, for off-pump CABG surgery, 487–488
- Color Doppler studies, in patent foramen ovale, 725–730
- Complement, release of, from cell-washing autotransfusion systems, 214
- Complications, perioperative, NHBLI recommendations on, 1–6
- Computed tomography, in pseudoaneurysm from pulmonary artery catheter, 668–670
- Congenital anomalies, difficult intubation in, 610–612
- Congenital heart disease
- bicaval venous cannulation in, obstruction detection in, 293–303
  - caval inflow occlusion technique for, in pediatric patients, 144–147
  - heart block, pacemaker implantation for, 628–629
  - postoperative intensive care for, in pediatric patients, 218–227
  - pulmonary artery shunt insertion in, end-tidal carbon dioxide monitoring during, 152–155
  - pulmonary hypertension in, brain natriuretic peptide and, 783, 784
  - repair of
    - cerebral function monitoring in, 645–656
    - ketorolac for, 454–457
    - superior vena caval partial occlusion during, 472–474
- Congestive heart failure
- brain natriuretic peptide and, 780–787
  - in pregnancy, mitral valve replacement for, 339–343
- Conservation, blood. *See* Blood conservation
- Constant-flow ventilation, in postoperative care, in pediatric patients, 221
- Continuous autotransfusion system, 210–217
- complement release in, 214
  - cytokine release in, 214
  - fat elimination in, 214
  - leukocyte quality in, 212–214
  - platelet quality in, 211–212
  - principles of, 210
  - red cell quality in, 211
  - tumor cell removal in, 214–215
- Contractility, left ventricular blood flow velocity and, 415–422
- Contusion, lung, pumpless extracorporeal lung assist in, 777–779
- Coronary artery(ies)
- angioplasty of, brain natriuretic peptide and, 782–783
  - anomalous origin of
    - with aortic coarctation, 327–331
    - in left sinus of Valsalva, 788–798
- Coronary artery bypass graft (CABG) surgery
- adhesion molecules after, simvastatin effects on, 605–609
  - anesthesia induction in, remifentanyl vs. fentanyl for, 263–268
  - aortic pseudoaneurysm after, 527–528
  - atrial fibrillation in, amiodarone and epidural analgesia and, 709–714
  - atrial tachyarrhythmias after, magnesium and sotalol effects on, 309–312
  - brain natriuretic peptide and, 782–783
  - cardiac biomarker release in, 34–37
  - cerebral oxygen saturation in, neurologic dysfunction and, 552–558
  - diclofenac after, 742–747
  - electrocardiographic changes after, 38–42
  - emboli in, 545–547, 548–551
  - endothelin production in, off-pump vs. conventional, 25–29
  - endotracheal intubation in, stress response to, 275–280
  - extubation after, vs. anesthesia type, 613–619
  - glycemic control after, 690–697

- Coronary artery bypass graft (CABG) surgery (*Continued*)  
 inflammatory response in, propofol effects on, 592–604  
 inotropic agents after, 401–403, 404–408  
 ischemic preconditioning for, 799–805  
 kidney dysfunction in, 492, 545–547, 548–551  
 multisystem organ failure after, 366–374  
 myocardial protection in. *See* Myocardial protection  
 in narcolepsy, anesthesia safety in, 201–203  
 off-pump, 486–505  
   anastomosis quality in, 493–494  
   anesthesia for, 488–489  
   for anomalous coronary artery correction, 788–798  
   approach to, 487  
   arteriotomy bleeding in, 488  
   atrial fibrillation in, 492–493, 704–708  
   blood loss in, 492  
   cardiac biomarker release in, 34–37  
   cardiac output monitoring during, 43–46  
   cardiovascular dysfunction in, 491–492  
   coagulation disorders in, 492  
   cost savings with, 493  
   duration of stay and, 493  
   electrocardiographic changes after, 38–42  
   endothelin production in, 25–29  
   exposure for, 487  
   future of, 496  
   hemodynamic instability in, 488  
   high thoracic epidural anesthesia for, 256–262  
   in high-risk patients, 494  
   historic background of, 486–487  
   indications for, 488  
   intrathecal morphine for, 451–453  
   kidney dysfunction in, 492  
   limitations of, 495–496  
   mortality in, 493  
   myocardial protection in, 488  
   neurocognitive effects of, 489–491  
   outcome of, 495  
   patient-controlled analgesia for, 755–758  
   pulmonary dysfunction in, 491, 698–703  
   quality of life considerations in, 494–495  
   reappraisal of, 253–255  
   stabilization in, 487–488  
   stroke in, 489–491  
   systemic impact of, 489–495  
   systemic inflammatory response in, 489  
   transfusion requirements for, 492  
 oxycodone during and after, 748–754  
 patent foramen ovale diagnosis during, 101–106  
 perioperative enoximone in, 409–414  
 pulmonary artery pseudoaneurysm after, computed tomography in, 668–670  
 pulmonary dysfunction in, 698–703  
 sheared catheter wire in, 765–766  
 ventricular pseudoaneurysm after, 671–672  
 in women, 344–352  
   long-term mortality in, 347  
   morbidity in, 347–348  
   research on, 348–350  
   short-term mortality in, 345–347  
 Corticosteroids, preoperative, delayed-onset adrenal insufficiency despite, 80–81  
 Cost(s)  
   of blood conservation, 22S  
 Cost(s) (*Continued*)  
   of off-pump CABG surgery, 493  
   of perioperative complications, NHBLI recommendations on, 1–6  
   of transfusions, 2S–3S, 15S  
 Cost-effectiveness  
   of blood conservation, 26S–27S  
   of perioperative beta-blockade, 7–13  
 C-reactive protein, kinetics of, after cardiopulmonary bypass, in pediatric patients, 160–165  
 Creatinine, elevation of, after CABG surgery, vs. emboli count, 548–551  
 Creutzfeldt-Jacob disease organisms, new variant, in blood supply, 1S–2S  
 Crystalloid, for pump priming, vs. albumin, 429–437  
 Cyanosis, pulmonary artery shunt insertion in, end-tidal carbon dioxide monitoring during, 152–155  
 Cytochrome P450 family, pharmacogenetics of, 357–359  
 Cytokines, release of  
   in CABG surgery, propofol effects on, 592–604  
   from cell-washing autotransfusion systems, 214  
   intraoperative, quinaprilat effects on, 47–52  
   in myocardium, sevoflurane effects on, 269–274  
 Danaparoid, for heparin-induced thrombocytopenia, 95  
 Deairing, carbon dioxide for, 180–184  
 Decision modeling, of perioperative beta-blockade strategies, in aortic aneurysm surgery, 7–13  
 Desflurane  
   for cardiac pheochromocytoma resection, 630–631  
   for ischemic preconditioning, 803  
 Diabetes mellitus  
   CABG surgery in, glycemic control in, 690–697  
   cardiac surgery in, kidney failure risk in, 442–445  
 Diastolic dysfunction, brain natriuretic peptide and, 780–782  
 Diclofenac, after CABG surgery, 742–747  
 Dilated cardiomyopathy, treatment of, in pediatric patients, 772–774  
 Discontinuous autotransfusion system, 210–217  
   complement release in, 214  
   cytokine release in, 214  
   fat elimination in, 214  
   leukocyte quality in, 212–214  
   platelet quality in, 211–212  
   principles of, 210  
   red cell quality in, 211  
   tumor cell removal in, 214–215  
 Diuretics  
   for liver transplantation, 438–441  
   postoperative, for pediatric patients, 222  
 DNA analysis, in pharmacogenetics, 354–355  
 Dobutamine  
   for cardiopulmonary bypass separation, 404–408  
   left ventricular pressure-relationship indices and, 415–422  
 Dopamine  
   for cardiopulmonary bypass separation, 404–408  
   for postoperative care, in pediatric patients, 220  
 Doppler studies  
   esophageal, in continuous cardiac output measurement, 185–189  
   in mesenteric blood flow alterations, after cardiopulmonary bypass, 731–733  
   in patent foramen ovale, 725–730  
   transcranial, in cerebral function monitoring, in pediatric cardiac surgery, 647–648  
 Down syndrome, difficult intubation in, 610–612  
 Ductus arteriosus, patent, interruption of, in cardiopulmonary bypass, 469–471

- Echocardiography  
 in aortic pseudoaneurysm, 527–528  
 in left ventricular pressure-relationship indices measurement, 415–422  
 transesophageal. *See* Transesophageal echocardiography
- Edema  
 postoperative, in pediatric patients, 222  
 pulmonary, in off-pump CABG surgery, 491
- Editorials  
 blood glucose control, 687–689  
 cardiopulmonary bypass separation, 401–403
- Electrocardiography  
 after CABG surgery, 38–42  
 in general anesthesia, with cardiovascular disease, 281–287
- Electroencephalography, in cerebral function monitoring, in pediatric cardiac surgery, 647
- Embolectomy, femoral, atrial appendage thrombosis dislodgment in, transesophageal echocardiography in, 809–811
- Embolism  
 air. *See* Air embolism  
 in CABG surgery, 545–547, 548–551  
 iliac artery, from sinus of Valsalva thrombus, 384–386  
 pulmonary, removal of, deep hypothermic circulatory arrest in, femoral-to-radial arterial pressure gradient in, 175–179
- Emergency surgery, bloodstream infections in, 304–308
- Emphysematous bullae, atrial septal defect with, simultaneous repair of, 632–636
- Enderectomy, carotid  
 myocardial ischemia in, risk factors for, 288–292  
 shunting in, 375–383
- Endothelin, in CABG surgery, off-pump vs. conventional, 25–29
- Endotracheal intubation  
 difficult, in pediatric patients, 610–612  
 double-lumen, tracheobronchial rupture from, 228–233  
 extubation in  
 vs. anesthesia type, 613–619  
 timing of, postoperative pain and, 64–67  
 stress in, laryngoscopy vs. laryngeal mask airway for, 275–280
- Enflurane, for ischemic preconditioning, 803
- Enoximone, for CABG surgery, 409–414
- Enterobacter*, in central venous catheter-related infections, 304–308
- Epidural analgesia and anesthesia  
 for atrial fibrillation, after CABG surgery, 709–714  
 hematoma from, 759–762  
 for off-pump CABG surgery, 256–262  
 thoracic, neurologic deficits due to, 512–520  
 during and after thoracotomy, 317–321, 458–460
- Epinephrine  
 for cardiopulmonary bypass separation, 404–408  
 for postoperative care, in pediatric patients, 220
- Epsilon aminocaproic acid, for blood conservation, 38S  
 in pediatric patients, 141–143
- Erythropoietin  
 for postoperative anemia, 59–63  
 for preoperative autologous blood donation, 6S–7S, 29S
- Escherichia coli*, in central venous catheter-related infections, 304–308
- Esmolol, perioperative, in aortic aneurysm surgery, cost-effectiveness of, 7–13
- Esophageal Doppler studies, for continuous cardiac output measurement, 185–189
- Esophagectomy, left lower lobe collapse after, ventilation for, 482–485
- Esophagus, tear of, in transesophageal echocardiography, 78–79
- Exchange transfusions, in sickle cell trait, for cardiopulmonary bypass, 663–667
- Extracorporeal lung assist, for blunt trauma, 777–779
- Extubation  
 in off-pump CABG surgery, intrathecal morphine effects on, 451–453  
 timing of  
 vs. anesthesia type, 613–619  
 postoperative pain and, 64–67
- Fasciitis, necrotizing, vacuum-assisted closure for, 336–338
- Fat, removal of, from cell-washing autotransfusion systems, 214
- Femoral embolectomy, atrial appendage thrombus dislodgment in, transesophageal echocardiography in, 809–811
- Femoral-to-radial arterial pressure gradient, in deep hypothermic circulatory arrest, 175–179
- Fentanyl  
 interpleural, for post-thoracotomy pain, 461–465  
 patient-controlled, for off-pump CABG surgery, 755–758  
 vs. remifentanyl, for induction, in cardiac disease, 263–268
- Fibroelastoma, papillary, of aortic valve, 114
- Fick technique, indirect, for pulmonary blood flow measurement, 47–52
- Fluid management  
 in liver transplantation, 438–441  
 postoperative, in pediatric patients, 222–223
- Fontan-type operations, postoperative care in, 224–225
- Foramen ovale, patent  
 intraoperative diagnosis of, 101–106  
 pulmonary artery catheter for, 763–764  
 transesophageal echocardiography in, 725–730
- Foreign objects, left in valve replacement, 242–244
- Free radicals, in CABG surgery, propofol effects on, 592–604
- Furosemide, postoperative, for pediatric patients, 222
- Gas, blood, monitoring of, during thoracoscopic surgery, 587–591
- Gas-diffuser device, for carbon dioxide deairing, 180–184
- Gender differences. *See also* Women  
 in kidney failure, after cardiac surgery, 442–445  
 in pharmacogenetics, 355
- General anesthesia  
 for off-pump CABG surgery, 488–489  
 high thoracic epidural anesthesia with, 256–262  
 QT dispersion in, in peripheral vascular disease and silent myocardial ischemia, 281–287
- Genetic factors, in anesthesia. *See* Pharmacogenetics
- Glucose control, in intensive care, 690–697
- Glycocalicin, heparin and, 93–94
- Glycoproteins, platelet, heparin and, 93–94
- Granuloma, subglottic, after lung resection, 479–481
- Halothane  
 arrhythmias due to, in magnesium deficiency, 313–316  
 left ventricular pressure-relationship indices and, 415–422
- Heart  
 myxoma of, atrial, cardiac arrest in, resuscitation in, 769–771  
 pheochromocytoma of, resection of, 630–631  
 rupture of, in anesthesia induction, 767–768  
 transplantation of, brain natriuretic peptide and, 782  
 univentricular, brain natriuretic peptide and, 783, 784
- Heart block, congenital, pacemaker implantation for, 628–629
- Heart failure  
 brain natriuretic peptide and, 780–787  
 in pregnancy, mitral valve replacement for, 339–343
- Hematocrit, in transfusion decision making, 234–241
- Hematoma, from thoracic epidural analgesia, 759–762

- Hemodilution  
in cardiopulmonary bypass, cerebral blood flow and, 423–428  
for preoperative autologous blood donation, 7S–8S
- Hemodynamics  
in endotracheal intubation, induction effects on, 263–268  
intraoperative, quinaprilat effects on, 47–52  
in liver transplantation, 438–441  
in off-pump CABG surgery, 488  
support of, for postoperative care, in pediatric patients, 219–221
- Heparin, 93–100  
activated coagulation time measurement with, 573–580  
aldosterone and, 96–97  
antithrombin III and, 93  
chemical structure of, 93  
inflammation and, 95–96  
lipid metabolism and, 97  
malignancy and, 96  
metabolism and, 97  
osteoporosis and, 97  
platelets and, 93–94  
preoperative, resistance development in, 131–135  
resistance to, 129–135  
thrombin and, 93  
thrombocytopenia induced by, 94–95  
thyroid hormones and, 97
- Histamine, release of, in CABG surgery, oxycodone and, 748–754
- Hyperglycemia, control of, in intensive care, 690–697
- Hyperkalemia, heparin-induced, 96–97
- Hypertension, pulmonary  
brain natriuretic peptide and, 783  
lung transplantation for, severe intraoperative hypoxemia in, 204–206  
milrinone and nitric oxide effects on, after cardiopulmonary bypass, in pediatric patients, 156–159  
sildenafil with nitric oxide for, 775–776
- Hyperthermia, malignant, pharmacogenetics of, 355–356
- Hyperventilation therapy, in postoperative care, in pediatric patients, 221
- Hypothermia, cerebral, monitoring of, in pediatric cardiac surgery, 646–647
- Hypothermic cardiopulmonary bypass, circulating-water rewarming garment for, in pediatric patients, 148–151
- Hypoxemia, in single-lung transplantation, 204–206
- Iliac artery, embolism in, from sinus of Valsalva thrombus, 384–386
- Indirect Fick technique, for pulmonary blood flow measurement, 47–52
- Indomethacin, after thoracotomy, 458–460
- Induction, in cardiac disease, remifentanyl vs. fentanyl for, 263–268
- Infants. *See* Pediatric patients
- Infarction  
mesenteric, surgery for, prognostic factors in, 734–741  
myocardial. *See* Myocardial infarction
- Infections, central venous catheter-related, prospective study of, 304–308
- Inflammation  
in CABG surgery  
off-pump vs. conventional, 25–29  
propofol effects on, 592–604  
heparin and, 95–96  
intraoperative quinaprilat for, hemodynamic results of, 47–52
- Inflow occlusion, for congenital heart disease repair, 144–147
- Inotropic agents  
for CABG surgery, 409–414  
for cardiopulmonary bypass separation, 401–403, 404–408
- Insufflation, carbon dioxide, in air embolism prevention, 180–184
- Insulin, for glycemic control, in intensive care, 690–697
- Integrins, heparin and, 95
- Intensive care  
glycemic control in, 690–697  
for pediatric patients, 218–227  
antidiuretics in, 222  
anti-inflammatory drugs in, 222  
fluid management, 222–223  
gastrointestinal interventions in, 222–223  
inhaled nitric oxide in, 218–219  
mechanical ventilation in, 221  
pharmacologic hemodynamic support in, 219–221  
thyroid hormones in, 222  
in univentricular physiology, 223–224  
personnel directing, 521–526
- Intercellular adhesion molecule, after CABG surgery, simvastatin effects on, 605–609
- Intercostal nerve block, for video-assisted thoracoscopy, 317–321
- Interleukin(s), in CABG surgery, propofol effects on, 592–604
- Interleukin-6, release of, in myocardium, sevoflurane effects on, 269–274
- Intra-aortic balloon counterpulsation  
bloodstream infections in, 304–308  
kidney failure risk with, 442–445
- Intrapulmonary shunt, measurement of, indirect Fick method for, 47–52
- Intubation, endotracheal. *See* Endotracheal intubation
- Iron hydroxide sucrose complex, intravenous, for postoperative anemia, 59–63
- Ischemia  
mesenteric, surgery for, prognostic factors in, 734–741  
myocardial. *See* Myocardial ischemia
- Isoflurane  
for CABG surgery, extubation time with, 613–619  
for ischemic preconditioning, 803  
minimum alveolar concentration of, in cardiopulmonary bypass, 620–623
- Jugular bulb oximetry, in cerebral function monitoring, in pediatric cardiac surgery, 648
- Jugular vein, left internal, pseudoaneurysm of, from venipuncture, 806–808
- Ketobemidone, for postoperative pain, extubation timing and, 64–67
- Ketorolac, for postoperative pain, in congenital heart surgery, 454–457
- Kidney, dysfunction/failure of  
in CABG surgery, 366–374, 545–547, 548–551  
off-pump, 492  
after cardiac surgery, incidence and risk of, 442–445
- Laryngeal mask airway and laryngoscopy, stress response in, 275–280
- Laser revascularization  
percutaneous, vs. transmyocardial laser revascularization, 87  
transmyocardial, review of, 85–92
- Lepirudin, for heparin-induced thrombocytopenia, 95
- Letters to the editor  
abdominal wall mucormycosis after heart transplantation, 822–823  
atrial septal defect vs. patent foramen ovale, 824  
combined transesophageal echocardiography and laryngeal mask airway, 395–396  
correlation of bispectral index with tet spells, 682–683  
defective triple-lumen catheter, 680–681

- Letters to the editor (*Continued*)
- endothelin release after off-pump coronary artery bypass graft surgery, 680
  - esophageal Doppler monitoring in off-pump surgery, 539–541
  - heparin reversal in coronary artery bypass surgery, 394–395
  - high thoracic epidural anesthesia for coronary artery bypass surgery, 392–394
  - inadvertent high dose epinephrine administration, 121
  - liver transplantation anesthesia, 825–826
  - massive epistaxis from nasopharyngeal probe, 123–124
  - nasopharyngeal angiofibroma with hemophilia A, 819
  - paravalvular abscess after aortic valve replacement, 820–821
  - propofol infusion into cardiopulmonary bypass pump, 122–123
  - pyruvate-fortified cardioplegia, 537–539
  - recurrent laryngeal nerve palsy after esophagectomy, 822
  - venous air embolism, 681–682
- Leukocytes, quality of, from cell-washing autotransfusion systems, 212–214
- Levosimendan, for dilated cardiomyopathy, 772–774
- Lidocaine, epidural, after thoracotomy, 458–460
- Lipid metabolism, heparin and, 97
- Lipid peroxidation, in CABG surgery, propofol effects on, 592–604
- Lipoproteins, heparin and, 97
- Lithotomy position, vs. peripheral and central venous pressures, 446–450
- Liver
- dysfunction of, after CABG surgery, 366–374
  - transplantation of
    - blood conservation in, 31S–37S
    - fluid management in, 438–441
- Lobectomy, for living-related lung transplantation, 506–511
- Lung
- biopsy of, video-assisted thoracoscopy in, intercostal nerve block for, 317–321
  - blood flow in, measurement of, indirect Fick method for, 47–52
  - dysfunction of, in CABG surgery, 491, 698–703
  - emphysematous bullae in, atrial septal defect with, simultaneous repair of, 632–636
  - extracorporeal assist for, in blunt trauma, 777–779
  - perfusion of, importance of, 166–174
  - resection of, subglottic granuloma after, 479–481
  - transplantation of
    - intraoperative hypoxemia during, 204–206
    - living-related, 506–511
- Mac-1, heparin and, 95
- Mad cow disease organisms, in blood supply, 1S–2S
- Magnesium
- for atrial tachyarrhythmias, after CABG surgery, 309–312
  - deficiency of, arrhythmias in, during anesthesia, 313–316
- Malignant hyperthermia, pharmacogenetics of, 355–356
- Malondialdehyde, in CABG surgery, propofol effects on, 592–604
- Mediastinum
- central venous catheter placement in, 75–77
  - necrotizing fasciitis of, vacuum-assisted closure for, 336–338
- Medivent RTX respirator, for left lower lobe collapse, after esophagectomy, 482–485
- Mesenteric blood flow, after cardiopulmonary bypass, Doppler studies of, 731–733
- Mesenteric infarction, surgery for, prognostic factors in, 734–741
- Metabolism, drug, genetic variations in. *See* Pharmacogenetics
- Metastasis, heparin action in, 96
- Methylprednisolone, postoperative, for pediatric patients, 222
- Metoprolol, perioperative, in aortic aneurysm surgery, cost-effectiveness of, 7–13
- Milrinone
- for cardiopulmonary bypass separation, 404–408
  - for pediatric patients, 220
  - pulmonary artery pressure effects of, after cardiopulmonary bypass, 156–159
- Mitral valve
- regurgitation of, brain natriuretic peptide and, 783
  - replacement of
    - atrial natriuretic peptide levels in, 30–33
    - cardiopulmonary bypass separation after, 404–408
    - gauze object left during, 242–244
    - left atrial thrombus formation after, 637–640
    - in pregnancy, 339–343
    - surgery on, kidney failure risk in, 442–445
- Monitoring, in carotid endarterectomy, 375–383
- Morphine, for off-pump CABG surgery
- intrathecal, 451–453
  - patient-controlled, 755–758
- Mortality
- in CABG surgery, in women, 345–348
  - in mesenteric infarction, 734–741
  - in off-pump CABG surgery, 493
- Multidrug resistance transporter, genetic variations in, 359
- Multiperforated catheter, for carbon dioxide deairing, 180–184
- Multisystem organ failure, after CABG surgery, 366–374
- Myocardial infarction
- in aortic aneurysm surgery, beta-blockers for, 7–13
  - in noncardiac surgery, beta-blockers for, 14–24
  - ventricular rupture in, in anesthesia induction, 767–768
- Myocardial ischemia
- in CABG surgery
    - enoximone for, 409–414
    - off-pump, 491
    - for preconditioning, 799–805
  - silent, general anesthesia in, QT dispersion in, 281–287
- Myocardial protection
- enoximone for, 409–414
  - ischemic preconditioning for, 799–805
  - in off-pump CABG surgery, 488
  - sevoflurane for, 269–274
- Myocarditis, treatment of, in pediatric patients, 772–774
- Myxoma, atrial, cardiac arrest in, resuscitation in, 769–771
- Nadoparin, for heparin-induced thrombocytopenia, 95
- Narcolepsy, cardiac surgery in, anesthesia safety in, 201–203
- National Heart, Blood and Lung Institute (NHLBI), perioperative medicine recommendations of, 1–6
- Natriuretic peptide, brain. *See* Brain natriuretic peptide
- Nausea and vomiting, postoperative, acupressure wristbands for, 68–71
- Near-infrared spectroscopy, in cerebral function monitoring, in pediatric cardiac surgery, 648–649
- Necrotizing fasciitis, vacuum-assisted closure for, 336–338
- Nesiritide, for heart failure, 784–785
- Neurologic complications
- in CABG surgery
    - off-pump, 489–491
    - prevention of, aortic atherosclerosis detection for, 715–718
  - in cerebral oxygen desaturation, 552–558
  - in thoracic epidural supplementation, 512–520
- Neurologic function, monitoring of, in pediatric patients, 645–656
- biochemical markers in, 650–652
  - bispectral index monitor in, 647
  - electroencephalography in, 647



- Neurologic function, monitoring of, in pediatric patients, 645–656  
(*Continued*)  
jugular bulb oximetry in, 648  
multimodality, 652–653  
near-infrared spectroscopy in, 648–649  
temperature measurement, 646–647  
transcranial Doppler studies in, 647–648
- NHLBI (National Heart, Blood and Lung Institute), perioperative medicine recommendations of, 1–6
- Nitric oxide  
inhaled, with sildenafil, for pulmonary hypertension, 775–776  
for postoperative care, in pediatric patients, 218–219  
pulmonary artery pressure effects of, after cardiopulmonary bypass, in pediatric patients, 156–159
- Nitroglycerin, for liver transplantation, 438–441
- Nonsteroidal anti-inflammatory drugs, after CABG surgery, 742–747
- Norepinephrine, for liver transplantation, 438–441
- Norwood operation, postoperative care in, 223–224
- Octopus Stabilization System, for off-pump CABG surgery, 487
- Off-pump CABG surgery. *See* Coronary artery bypass graft (CABG) surgery, off-pump
- One-lung ventilation, cardiac output measurement in, 190–193
- OPCAB. *See* Coronary artery bypass graft (CABG) surgery, off-pump
- Orthopedic surgery  
blood conservation in, 29S–30S, 40S–41S  
with cardiovascular disease, general anesthesia in, 281–287
- Osteoporosis, heparin-induced, 97
- Outcomes, of perioperative beta-blockade, in aortic aneurysm surgery, 7–13
- Oximetry, jugular bulb, in cerebral function monitoring, in pediatric cardiac surgery, 648
- Oxycodone, during and after CABG surgery, 748–754
- Oxygen saturation  
cerebral, neuropsychologic dysfunction and, 552–558  
monitoring of, in systemic-to-pulmonary artery shunt insertion, 152–155
- Pacemaker, for congenital heart block, 628–629
- Pain, postoperative. *See* Postoperative pain
- Papillary fibroelastoma, of aortic valve, 114
- Paracetamol, after CABG surgery, 742–747
- Paratrend blood gas monitoring system, during thoroscopic surgery, 587–591
- Partial rebreathing, in cardiac output measurement, before and after cardiopulmonary bypass, 563–572
- Patent ductus arteriosus, interruption of, in cardiopulmonary bypass, 469–471
- Patent foramen ovale  
intraoperative diagnosis of, 101–106  
pulmonary artery catheter for, 763–764  
transesophageal echocardiography in, 725–730
- Patient-controlled analgesia, for off-pump CABG surgery, 755–758
- Pediatric patients  
absent right superior vena cava with persistent left superior vena cava in, 332–335  
aortic coarctation repair in  
with coronary artery anomalous origin, 327–331  
jet ventilation during, 207–209  
bicaval venous cannulation in, obstruction detection in, 293–303  
blood loss prevention in, epsilon aminocaproic acid vs. tranexamic acid for, 141–143  
cardiopulmonary bypass in  
aortic interruption during, 469–471
- Pediatric patients (*Continued*)  
procalcitonin and C-reactive protein kinetics after, 160–165  
pulmonary artery pressures after, milrinone and nitric oxide effects on, 156–159  
cavopulmonary anastomosis in, superior vena caval partial occlusion during, 472–474  
congenital heart disease in  
caval inflow occlusion technique for, 144–147  
repair of, postoperative ketorolac for, 454–457  
dilated cardiomyopathy in, treatment of, 772–774  
heart block in, pacemaker implantation for, 628–629  
hypothermic cardiopulmonary bypass in, rewarming garment for, 148–151  
intubation of, difficult, 610–612  
postoperative intensive care for, 218–227  
pulmonary hypertension in, sildenafil with nitric oxide for, 775–776  
stroke in, after cardiopulmonary bypass with aprotinin, 194–196
- Perioperative medicine, NHLBI recommendations on, 1–6
- Peripheral vascular disease, general anesthesia in, QT dispersion in, 281–287
- Peripheral venous pressure, vs. central venous pressure, patient position and, 446–450
- Peroxidation, lipid, in CABG surgery, propofol effects on, 592–604
- Persistent left superior vena cava, central venous catheter in, 529–531
- Pharmacogenetics, 353–365  
allele frequencies in, 355  
of beta-adrenergic receptor, 359–360  
of cytochrome P450 family, 357–359  
gender differences in, 355  
gene structure and, 353–354  
genetic variation and, 354  
in malignant hyperthermia, 355–356  
of multidrug resistance transporter, 359  
nomenclature of, 355–357  
polymorphism-drug metabolism phenotype and, 355  
in pseudocholinesterase deficiency, 356–357  
testing in, 354–355
- Phenylephrine  
in cardiopulmonary bypass, cerebral blood flow and, 423–428  
for liver transplantation, 438–441
- Pheochromocytoma, cardiac, resection of, 630–631
- Phosphodiesterase inhibitors  
for CABG surgery, 409–414  
for postoperative care, in pediatric patients, 220
- Platelets  
aggregometry of, for platelet dysfunction assessment, 136–140  
cardiopulmonary bypass effects on, pump priming fluid and, 429–437  
deficiency of, heparin-induced, 94–95  
dysfunction of  
aspirin-induced, 136–140  
heparin-induced, 93–94  
quality of, from cell-washing autotransfusion systems, 211–212
- Plateletworks analyzer, for platelet dysfunction assessment, 136–140
- Pleura, biopsy of, video-assisted thoracoscopy in, intercostal nerve block for, 317–321
- Pleural space, anesthetic administration into, for post-thoracotomy pain, 461–465
- Pleurectomy, video-assisted thoracoscopy in, intercostal nerve block for, 317–321
- Position, patient, vs. peripheral and central venous pressures, 446–450
- Post cardiopulmonary bypass myocardial dysfunction, enoximone for, 409–414

- Postoperative pain  
 in CABG surgery  
   diclofenac and paracetamol for, 742–747  
   off-pump, patient-controlled analgesia for, 755–758  
   oxycodone for, 748–754  
 epidural analgesia for, neurologic deficits due to, 512–520  
 extubation timing and, 64–67  
 thoracic epidural analgesia for, atrial fibrillation and, 709–714  
 in thoracotomy  
   epidural local anesthetics for, 458–460  
   intercostal nerve block vs. morphine for, 317–321  
   interpleural bupivacaine with fentanyl for, 461–465  
 in video-assisted thoracoscopy, 317–322  
 Prednisolone, postoperative, for pediatric patients, 222  
 Prednisone, preoperative, delayed-onset adrenal insufficiency despite, 80–81  
 Procalcitonin, kinetics of, after cardiopulmonary bypass, in pediatric patients, 160–165  
 Prone position, vs. peripheral and central venous pressures, 446–450  
 Propofol  
   for CABG surgery  
     extubation time with, 613–619  
     lipid peroxidation and inflammatory response effects of, 592–604  
     safety of, in narcolepsy, 201–203  
*Proteus*, in central venous catheter-related infections, 304–308  
 Pseudoaneurysm  
   aortic, echocardiography of, 527–528  
   left internal jugular vein, from venipuncture, 806–808  
   pulmonary artery, from catheter, computed tomography in, 668–670  
   ventricular, behind left atrium, 671–672  
 Pseudocholinesterase deficiency, 356–357  
 Pulmonary artery  
   perfusion of, importance of, 166–174  
   shunt insertion in, end-tidal carbon dioxide monitoring during, 152–155  
 Pulmonary artery catheter  
   gravity-driven fluid infusions through, interruption of, 72–74  
   guided insertion of, transesophageal echocardiography in, 658–662  
   for patent foramen ovale reduction, 763–764  
   pseudoaneurysm due to, computed tomography in, 668–670  
 Pulmonary artery pressure, milrinone and nitric oxide effects on, after cardiopulmonary bypass, in pediatric patients, 156–159  
 Pulmonary artery thermodilution, for cardiac output measurement  
   continuous, 185–189  
   in one-lung ventilation, 190–193  
 Pulmonary blood flow, measurement of, indirect Fick method for, 47–52  
 Pulmonary edema, in off-pump CABG surgery, 491  
 Pulmonary embolism, removal of, deep hypothermic circulatory arrest  
   in, femoral-to-radial arterial pressure gradient in, 175–179  
 Pulmonary fibrosis, living-related lung transplantation in, 506–511  
 Pulmonary hypertension  
   brain natriuretic peptide and, 783, 784  
   lung transplantation for, severe intraoperative hypoxemia in, 204–206  
   postoperative, in pediatric patients, 218–219  
   sildenafil with nitric oxide for, 775–776  
 Pulse contour analysis, for continuous cardiac output measurement, 185–189  
 QT dispersion, in general anesthesia, with cardiovascular disease, 281–287  
 Quality of life, in off-pump CABG surgery, 494–495  
 Quinaprilat, intraoperative, hemodynamic results of, 47–52  
 Rebreathing, partial, in cardiac output measurement, before and after cardiopulmonary bypass, 563–572  
 Red blood cells, quality of, from cell-washing autotransfusion systems, 211  
 Remifentanyl  
   for cardiac pheochromocytoma resection, 630–631  
   vs. fentanyl, for induction, in cardiac disease, 263–268  
   patient-controlled, for off-pump CABG surgery, 755–758  
   safety of, in narcolepsy, 201–203  
 Resistance, heparin, 129–130  
 Respiratory disorders, in off-pump CABG surgery, 451–453, 491  
 Resuscitation  
   in atrial myxoma, 769–771  
   in cardiac rupture, 767–768  
 Revascularization. *See also* Coronary artery bypass graft (CABG) surgery  
   laser, transmyocardial, review of, 85–92  
   percutaneous, vs. transmyocardial laser revascularization, 87  
 Rewarming, circulating-water garment for, in pediatric patients, 148–151  
 Risk assessment, perioperative, NHBLI recommendations on, 1–6  
 S-100 protein, in cerebral function monitoring, in pediatric cardiac surgery, 650–652  
 Sedation, assessment of, midlatency auditory-evoked potentials in, 559–562  
 Selectins  
   after CABG surgery, simvastatin effects on, 605–609  
   heparin and, 95–96  
 Sepsis, after cardiopulmonary bypass, in pediatric patients, 160–165  
 Septectomy, atrial, caval inflow occlusion in, in pediatric patients, 144–147  
 Sevoflurane  
   arrhythmias due to, in magnesium deficiency, 313–316  
   for CABG surgery, extubation time with, 613–619  
   for ischemic preconditioning, 803–804  
   for myocardial protection, 269–274  
 Shoulder pain, after thoracotomy, epidural analgesia for, 458–460  
 Shunt  
   brain natriuretic peptide and, 784  
   in carotid endarterectomy, 375–383  
   intrapulmonary, measurement of, indirect Fick method for, 47–52  
 Sick cell trait, cardiopulmonary bypass in, exchange transfusion requirements in, 663–667  
 Sildenafil, with nitric oxide, for pulmonary hypertension, 775–776  
 Simvastatin, adhesion molecule effects of, after CABG surgery, 605–609  
 Sinus of Valsalva  
   right anomalous coronary artery originating in, 788–798  
   thrombosis of, in bicuspid aortic valve, 384–386  
 SonoSite ultrasound device, in aortic atherosclerosis, 715–718  
 Sotalol, for atrial tachyarrhythmias, after CABG surgery, 309–312  
 Spectroscopy, near-infrared  
   in cerebral function monitoring, in pediatric cardiac surgery, 648–649  
   for obstruction detection, in bicaval venous cannulation, 293–303  
 Stabilization, for off-pump CABG surgery, 487–488  
*Staphylococcus*, in central venous catheter-related infections, 304–308  
 Stent, aortic, placement of, transesophageal echocardiography in, 466–468  
 Stress, in endotracheal intubation, laryngoscopy vs. laryngeal mask airway for, 275–280  
 Stroke  
   in carotid artery thrombosis, after cardiopulmonary bypass, with aprotinin, in pediatric patients, 194–196

- Stroke (*Continued*)  
 in off-pump CABG surgery, 489–491  
 prevention of, in pediatric cardiac surgery, neuromonitoring for, 645–656
- Stroke work, left ventricular blood flow velocity and, 415–422
- Subglottic granuloma, after lung resection, 479–481
- Superior vena cava, obstruction of, in cannulation, detection of, 293–303
- Supine position, vs. peripheral and central venous pressures, 446–450
- Systemic inflammatory response syndrome  
 after CABG surgery, 366–374  
 after cardiopulmonary bypass, in pediatric patients, 160–165  
 intraoperative, quinaprilat effects on, 47–52  
 in off-pump CABG surgery, 489
- Systemic vascular resistance, intraoperative, quinaprilat effects on, 47–52
- Temperature, cerebral, monitoring of, in pediatric cardiac surgery, 646–647
- Thermodilution  
 in cardiac output measurement, before and after cardiopulmonary bypass, 563–572  
 pulmonary artery, for cardiac output measurement  
 continuous, 185–189  
 in one-lung ventilation, 190–193
- Thoracic epidural anesthesia  
 for atrial fibrillation, after CABG surgery, 709–714  
 hematoma from, 759–762  
 high, for off-pump CABG surgery, 256–262
- Thoracoscopy  
 blood gas monitoring during, 587–591  
 video-assisted, intercostal nerve block for, 317–321
- Thoracotomy, pain after  
 epidural local anesthetics for, 458–460  
 intercostal nerve block vs. morphine for, 317–321  
 interpleural bupivacaine with fentanyl for, 461–465
- Thrombin, heparin binding to, 93
- Thrombocytopenia, heparin-induced, 94–95
- Thromboelastography, in ionized calcium vs. coagulation studies, 581–586
- Thromboendarterectomy, pulmonary, deep hypothermic circulatory arrest in, femoral-to-radial arterial pressure gradient in, 175–179
- Thrombosis  
 atrial appendage, dislodgment of, in femoral embolectomy, 809–811  
 carotid artery, after cardiopulmonary bypass, with aprotinin, in pediatric patients, 194–196  
 left atrial, after mitral valve replacement, 637–640  
 sinus of Valsalva, in bicuspid aortic valve, 384–386
- Thyroid hormones  
 heparin and, 97  
 postoperative, for pediatric patients, 222
- Trachea  
 intubation of. *See* Endotracheal intubation  
 nonpenetrating traumatic transection of, 475–478  
 rupture of, in endotracheal intubation, 228–233
- Tranexamic acid, for blood loss prevention, in pediatric patients, 141–143
- Transcranial Doppler studies, in cerebral function monitoring, in pediatric cardiac surgery, 647–648
- Transesophageal echocardiography  
 in anomalous coronary artery correction, 788–798  
 in aortic stent placement, 466–468  
 in aortic valve mass, 114
- Transesophageal echocardiography (*Continued*)  
 in atrial appendage thrombus dislodgment, 809–811  
 in CABG surgery  
 in cardiac output monitoring, 43–46  
 sheared catheter wire detection by, 765–766  
 in cardiopulmonary bypass separation, 401–403, 404–408  
 intraoperative  
 gauze object found in, 242–244  
 in noncardiac surgery, utility of, 107–111  
 in left atrial thrombus, 637–640  
 in patent foramen ovale, 101–106, 725–730  
 probe of, esophageal tear from, 78–79  
 in pulmonary artery catheter guided insertion, 658–662  
 in tricuspid regurgitation, 719–724  
 in ventricular pseudoaneurysm, 671–672
- Transfusion(s)  
 blood conservation and. *See* Blood conservation  
 blood shortages for, 1S–2S  
 cell-washing autotransfusion systems for, continuous vs. discontinuous, 210–217  
 costs of, 2S–3S, 15S  
 demand for, 2S  
 efficacy of, 3S–4S  
 errors in, 15S  
 exchange, in sickle cell trait, for cardiopulmonary bypass, 663–667  
 hematocrit criteria for, 234–241  
 in liver transplantation, 438–441  
 for off-pump CABG surgery, 492  
 pathogens in, 1S  
 refusal of, by Jehovah's Witness patient, 765–766  
 risks of, 234–235
- Transmyocardial laser revascularization, 85–92  
 angiogenesis in, 86  
 background of, 85–86  
 candidates for, 87  
 clinical trials of, 87–90  
 lasers for, 86  
 myocardial necrosis in, 86  
 vs. percutaneous revascularization, 87  
 placebo effect in, 86  
 techniques for, 86–87
- Transplantation  
 heart, brain natriuretic peptide and, 782  
 liver  
 blood conservation in, 31S–37S  
 fluid management in, 438–441  
 lung  
 intraoperative hypoxemia during, 204–206  
 living-related, 506–511
- Transpulmonary aortic thermodilution, for cardiac output measurement, in one-lung ventilation, 190–193
- Trauma  
 chest, pumpless extracorporeal lung assist in, 777–779  
 tracheal, management of, 475–478
- Trendelenburg position, vs. peripheral and central venous pressures, 446–450
- Tricuspid valve, regurgitation of, transesophageal echocardiography in, 719–724
- Triiodothyronine, postoperative, for pediatric patients, 222
- Troponin I, release of  
 in CABG surgery, 34–37, 409–414  
 off-pump, 491  
 in ischemic preconditioning, 803
- Tube(s), endotracheal. *See* Endotracheal intubation

- Tumor cells, removal of, from cell-washing autotransfusion systems, 214
- Tumor necrosis factor- $\alpha$ , release of, in myocardium, sevoflurane effects on, 269–274
- Turbidometric platelet aggregometry, for platelet dysfunction assessment, 136–140
- Ultrasonography  
 in aortic atherosclerosis, 715–718  
 in cardiac output measurement, before and after cardiopulmonary bypass, 563–572  
 in mesenteric blood flow alterations, after cardiopulmonary bypass, 731–733
- Univentricular heart  
 brain natriuretic peptide and, 783, 784  
 surgery for, postoperative care in, 223–225
- Vacuum-assisted closure, for necrotizing fasciitis treatment, 336–338
- Valve  
 aortic. *See* Aortic valve  
 mitral. *See* Mitral valve
- Vascular cell adhesion molecule, after CABG surgery, simvastatin effects on, 605–609
- Vasopressin, for postoperative care, in pediatric patients, 220
- Vena cava  
 anomalies of, absent right superior vena cava with persistent left superior vena cava in, 332–335  
 obstruction of, detection of, 293–303  
 occlusion of, in atrial septectomy, 144–147  
 superior  
 persistent, central venous catheter in, 529–531  
 unintentional partial occlusion of, in cavopulmonary anastomosis, 472–474
- Venipuncture, of left internal jugular vein, pseudoaneurysm formation in, 806–808
- Venous infusion port, for pulmonary artery catheter, gravity-driven fluid infusions through, interruption of, 72–74
- Venous pressure, peripheral, vs. central venous pressure, patient position and, 446–450
- Ventilation  
 constant-flow, in pediatric patients, 221  
 after esophagectomy, for left lower lobe collapse, 482–485  
 jet, during aortic coarctation repair, 207–209  
 mechanical, in postoperative care, in pediatric patients, 221  
 one-lung, cardiac output measurement in, 190–193  
 with pumpless extracorporeal device, for blunt trauma, 777–779  
 volume-targeted, in pediatric patients, 221
- Ventricle(s)  
 left  
 pressure overload of, brain natriuretic peptide and, 784  
 pressure-area relationship indices of, 415–422  
 pseudoaneurysm of, behind left atrium, 671–672  
 right, pressure overload of, brain natriuretic peptide and, 783  
 rupture of, in anesthesia induction, 767–768
- Ventricular arrhythmias, after aortic valve replacement, controlled aortic root perfusion for, 197–200
- Ventricular assist support, for dilated cardiomyopathy, 772–774
- Ventricular fibrillation  
 after CABG surgery, 366  
 intraoperative, salvage with secondary blood cardioplegia, 641–644
- Ventricular septal defect  
 brain natriuretic peptide and, 783, 784  
 pulmonary hypertension in, sildenafil with nitric oxide for, 775–776  
 repair of, aortic interruption during, 469–471
- Video-assisted thoracoscopy, intercostal nerve block for, 317–321
- Virginia Commonwealth University, blood conservation program of, 18S–23S
- Visual analog scale, for pain assessment, extubation timing and, 64–67
- Volume-targeted ventilation, in pediatric patients, 221
- Vomiting, postoperative, acupressure wristbands for, 68–71
- Warming garment, circulating-water, for pediatric patients, 148–151
- Women, CABG surgery in, 344–352  
 long-term mortality in, 347  
 morbidity in, 347–348  
 preoperative presentation of, 344–345  
 research on, 348–350  
 short-term mortality in, 345–347
- Wristbands, acupressure, for postoperative nausea and vomiting, 68–71