

- Varma, P.K., 136, 594  
 Veal, M., 149  
 Vega, R., 768  
 Vender, J.S., 156, 295, 469, 493  
 Vento, A.E., 526  
 Venugopal, P., 80  
 Vijayakumar, A., 126, 627  
 Vinatier, I., 668  
 Virmani, S., 128  
 Vislocky, I., 353  
 Vlahakes, G.J., 772  
 Vretzakis, G., 68  
 Vuylsteke, A., 748
- Walker, J., 834  
 Warters, R.D., 232  
 Wasnick, J.D., 826  
 Weber, T., 51
- Wei, J., 503  
 Weidemann, F., 340  
 Weiner, J., 8, 673  
 Weiss, S.J., 765  
 Weitzberg, E., 311  
 Wells, F.C., 748  
 Welsby, I.J., 531  
 Welters, I., 509  
 Wernet, A., 26  
 Westphal, B., 320  
 Wheatley, D.J., 541  
 White, S.A., 659  
 Williams, B.A., 554  
 Willigers, H.M., 364  
 Wiltink, E.H.H., 573  
 Winterhalter, M., 716  
 Woo, A., 759  
 Wouters, P.F., 466
- Wyatt, R., 762  
 Wynn, M.M., 273
- Yamak, B., 90  
 Yamamoto, K., 656  
 Yang, P., 905  
 Yazigi, A., 760, 803  
 Ye, L., 905  
 Yeatman, D., 106  
 Yildirim, V., 48  
 Yıldız, Y., 709  
 Yudin, M., 307
- Zangrillo, A., 31, 477, 788  
 Zeeni, C., 760  
 Zuber, K., 291  
 Zuk, J., 716  
 Zvara, D.A., 149

## SUBJECT INDEX

- Abdominal compartment syndrome, after congenital heart disease repair, 71–75  
 Abscess, aortic root, 280–283  
 Adhesion molecules, phosphodiesterase inhibitors and, 162–166  
 Adrenergic alpha-agonists, for sympathetic inhibition, myocardial oxygen consumption and, 364–370  
 Adrenergic beta-antagonists, for sympathetic inhibition, myocardial oxygen consumption and, 364–370  
 Aggregometry, for platelet dysfunction, in cardiopulmonary bypass, 536–540, 554–561  
 Air embolism  
 after congenital heart disease repair, 80–81  
 transesophageal echocardiography in, 387–389  
 Ajmaline test, for Brugada syndrome, 407–413  
 Albumin  
 in normovolemic hemodilution, hemodynamic changes in, 812–818  
 for volume replacement, 99  
 Allon Thermo-Wrapping Thermoregulation system, in off-pump CABG surgery, 788–792  
 Aminocaproic acid, for liver transplantation, 888–893  
 Amplatzer Asymmetric VSD Occluder, 202–208  
 Anesthesia. *See* Thoracic epidural anesthesia; Preconditioning; *specific agents*  
 Aneurysm  
 aortic, repair of. *See* Aorta, repair of  
 coronary artery, 119–120  
 left sinus of Valsalva, 276–279  
 pulmonary artery, 562–564  
 subaortic, 276–279  
 Animal studies  
 fenoldopam in ischemic kidney failure, 179–186  
 hemodynamic effects of sevoflurane, 209–216  
 intra-aortic balloon pumping effects on cardiac output determination, 320–324  
 left ventricular function evaluation, 340–346  
 lower limb ischemia-reperfusion injury, 187–195  
 meperidine effects on pulmonary vascular bed, 691–695  
 myocardial oxygen consumption in sympathetic stimulation, 364–370  
 sevoflurane-induced left ventricular depression, 358–363  
 sildenafil in pulmonary hypertension, 725–726  
 Anomalous pulmonary venous drainage, partial, 416  
 Anticoagulants  
 for CABG surgery, in prekallikrein deficiency, 580–582  
 thrombocytopenia induced by, 106–111  
 Antidiuretic hormone, in volume control, 98  
 Antifibrinolytics, for liver transplantation, 888–893  
 Aorta  
 atherosclerosis of, cardiopulmonary bypass outcome and, 149–155  
 coarctation of, 429  
 dissection of, in cocaine abuse, 458–460  
 repair of  
 bispectral index inaccuracy in, 68–70  
 bleeding control in, 570–572, 573–575, 736–741  
 bloody cerebrospinal fluid and, 269–275  
 cardiac output monitoring in, 26–30  
 with contained rupture and hemothorax, 401–404  
 deep hypothermic circulatory arrest for, 8–13, 673–677  
 with retrograde cerebral perfusion, 3–7  
 dilutional hypokalemia—induced arrhythmias in, 568–569  
 femoral or axillary artery cannulation complications in, 20–25  
 hemothorax after, TEE in, 229–231  
 left ventricular perforation in, 247–250  
 statins in, 251–258  
 tracheostomy after, 14–19  
 transesophageal echocardiography, 772–776  
 Aortic arch, anomalies of, 429  
 Aortic root abscess, 280–283  
 Aortic valve  
 bicuspid, 428  
 repair of  
 bleeding control in, 573–575  
 persistent ST elevation after, 397–400  
 replacement of  
 intraoperative pacing in, 117–118  
 in Parkinson's disease with deep-brain stimulation, 707–708  
 transesophageal echocardiography during, 847–849  
 stenosis of, dobutamine stress echocardiography in, 862–866  
 Arginine vasopressin  
 in hemorrhagic shock, 167–172  
 for hypotension, after pheochromocytoma resection, 394–396  
 Arrhythmias, in dilutional hypokalemia, in autologous transfusion, 568–569  
 Arterial switch operation, for transposition of great arteries, 428

- Arterial-to-end-tidal carbon dioxide tension difference, in congenital heart disease, 196–201
- Arteries. *See specific arteries*
- Atherosclerosis, aortic, cardiopulmonary bypass outcome and, 149–155
- Atrial appendage, left, thrombosis of, TEE in, 96–97
- Atrial fibrillation, prediction of, with white blood cell count, 51–56
- Atrial septal defect  
in adults, 416–420  
Gore-Tex patch for, echocardiographic visualization of, 751–752  
repair of, TEE in, 90–93
- Atrioventricular canal, 420
- Atrium  
left  
catheter malposition in, in inferior vena cava thrombectomy, 405–406  
enlargement, TEE in, 772–776  
errant catheter in, 565–567  
right, perforation of, from pulmonary artery catheter, 88–89
- Automatic external defibrillator, in air travel, 438–444
- Axillary artery, cannulation of, for aortic dissection repair, complications of, 20–25
- Bair Hugger forced-air system, in off-pump CABG surgery, 788–792
- Balloon occlusion, for pulmonary artery rupture, 376–378
- Bispectral index  
in anesthetic preconditioning evaluation, 484–492  
cardiopulmonary bypass machine effects on, in aortic repair, 68–70
- Bivalirudin, thrombocytopenia induced by, 106–111
- Blast injury, 872–880  
anesthesia management in, 876–878  
initial management in, 873–874  
patterns of, 874–876  
physics of explosions and, 872–873  
triage in, 873–874
- Bleeding  
in aortic aneurysm repair, control of, factor VII in, 570–572  
in aortic dissection repair, control of, factor VIIa in, 736–741  
in CABG surgery, autologous transfusion and, 541–547  
in cardiac surgery  
factor VIIa in, 573–575  
prediction of, Thromboelastograph for, 531–535  
in cardiopulmonary bypass, evaluation of, 536–540  
monitoring of, in liver transplantation, 548–553
- Blood, in cerebrospinal fluid, aortic aneurysm surgery and, 269–275
- Blood flow  
carotid artery, left ventricular elastance determination from, 340–346  
cerebral, in veno-veno bypass, 744–745
- Blood transfusions. *See Transfusion(s)*
- Bombing injury, 872–880
- Bosentan, for pulmonary hypertension, vs. sildenafil, 730
- Brain  
oxygenation monitoring in  
in cardiac surgery, 445–455, 606–615  
continuous jugular bulb venous saturation in, 826–833  
in hypothermic hemodilution, 819–825  
near-infrared spectroscopy in, 826–833  
protection of, for cardiac surgery, 520–525  
retrograde perfusion of  
air embolism in, 80–81  
with deep hypothermic circulatory arrest, for aortic arch repair, 3–7
- Bronchial occlusion, with secretions, 620–621
- Bronchial stenting, with laryngeal mask airway, 227–228
- Brugada syndrome, 407–413
- Bulla formation, in lung transplantation, 748–750
- Bundle branch block, right, in Brugada syndrome, 407–413
- Bupivacaine, for paravertebral block  
for thoracoscopic surgery, 639–643  
for thoracotomy, 644–647
- CABG surgery. *See Coronary artery bypass graft (CABG) surgery*
- Calcium, in intravenous fluids, coagulation profiles in, 807–811
- Cancer, lung, resection of, thoracic epidural anesthesia for, in multiple sclerosis, 704–706
- Carbon dioxide, absorption of, in saphenous vein graft harvesting, 347–352
- Carbon dioxide tension, arterial-to-end-tidal, in congenital heart disease, 196–201
- Cardiac arrest, in air travel, 438–444
- Cardiac arrhythmias, in blast injury, 875
- Cardiac catheterization, sedation for, in pediatric patients, 515–519
- Cardiac massage, in hemodynamic collapse, after CABG, 143–148
- Cardiac output, monitoring of  
in aortic repair, 26–30  
intra-aortic balloon pump effects on, 320–324  
after off-pump CABG, Doppler device for, 315–319  
pulmonary artery catheter in, complications of, 85–87
- Cardiac resynchronization therapy, tissue Doppler echocardiography in, 589
- Cardiac surgery. *See also specific procedures*  
anti-inflammatory effects of ketamine in, 217–222  
atrial fibrillation prediction after, white blood cell count and, 51–56  
bleeding after  
factor VIIa in, 573–575  
prediction of, 531–535  
cerebral oxygenation monitoring in, 445–455  
with coexistent hypertrophic cardiomyopathy, mitral stenosis, and coronary artery fistula, 594–605  
neurologic outcome after, electroencephalography in, 520–525  
neutrophil activity after, sevoflurane cardioplegia effects on, 57–62  
outcome of, aortic atherosclerosis effects on, 149–155  
preconditioning for. *See Preconditioning*  
pulmonary artery catheter in, wedging effects on, 311–314  
pulmonary hypertension after, sildenafil for, 731  
relative pulmonary hypertension in, 331–339  
repeat, percutaneous superior vena cava cannulation for, 881–887  
risk evaluation for, in octogenarians, 526–530  
statins in, 251–258  
volume therapy in, 98–105  
in von Willebrand's disease, in neonates, 576–579
- Cardiomyopathy  
hypertrophic obstructive, with mitral stenosis and coronary artery fistula, 594–605  
restrictive, tissue Doppler echocardiography in, 589
- Cardioplegia, sevoflurane in, neutrophil activity and, 57–62
- Cardioprotection, anesthetic preconditioning for. *See Preconditioning*
- Cardiopulmonary bypass  
air embolism in, 80–81  
autologous blood transfusions for, 541–547  
bispectral index inaccuracy in, in aortic repair, 68–70  
bleeding after, prediction of, 531–535  
hemodilution in, moderate vs. tepid, oxygen saturation in, 819–825  
heparinized oxygenators in, 554–561  
inflammatory effects of, on internal mammary artery, 63–67  
kidney dysfunction after, 684–690  
liver dysfunction after, 684–690

- Cardiopulmonary bypass (*Continued*)  
 for lung transplantation, 668–672  
 neurologic outcome after  
 electroencephalography in, 520–525  
 near-infrared spectroscopy monitoring for, 606–615  
 neutrophil activity after, sevoflurane cardioplegia effects on, 57–62  
 outcome of, aortic atherosclerosis effects on, 149–155  
 oxygenation in, near-infrared spectroscopy in, 606–615  
 oxygenation index after, vs. gas composition, 136–142  
 platelet dysfunction in, aggregometry in, 536–540  
 preconditioning for, 473–476  
 retrograde autologous priming in, 156–161  
 risk evaluation for, in octogenarians, 526–530  
 in von Willebrand's disease, in neonates, 576–579
- Cardiopulmonary resuscitation, in air travel, 438–444
- Cardiotomy suction, in CABG surgery, platelet function and, 554–561
- Carotid artery(ies), blood flow in, left ventricular elastance determination from, 340–346
- Carotid endarterectomy  
 CABG surgery combined with  
 risk in, 796–802  
 staged anesthesia for, 803–806  
 thoracic epidural anesthesia for, 850–852  
 cardiovascular collapse during, 259–268
- Catheter(ization)  
 cardiac, sedation for, in pediatric patients, 515–519  
 central venous, positioning, 777–780  
 femoral vein, retroperitoneal hematoma in, 859–861  
 internal jugular vein, innominate artery pseudoaneurysm in, 853–855  
 malposition of, in inferior vena cava thrombectomy, 405–406  
 pulmonary artery. *See* Pulmonary artery catheter  
 superior vena cava, in repeat sternotomy, 881–887  
 for ventricular septal defect closure, 202–208
- CD11b, phosphodiesterase inhibitors and, 162–166
- Central venous catheter, positioning of, 777–780
- Cerebrospinal fluid, bloody, aortic aneurysm surgery and, 269–275
- Chest radiography, in pulmonary artery location, 456–457
- Circulating-water garment, in off-pump CABG surgery, 788–792
- Circulatory arrest, hypothermic  
 for aortic arch repair, 3–7  
 for aortic repair, 673–677  
 for thoracic artery repair, 8–13  
 vascular response to thiopental in, 678–683
- Coagulation disorders  
 cardiac surgery in, in neonates, 576–579  
 in congenital heart disease, activated protein C and, 38–41  
 detection of, in liver transplantation, 548–553  
 evaluation of, in cardiopulmonary bypass, 536–540  
 in fluid therapy, calcium supplementation effects on, 807–811  
 hetastarch-induced, 102–103  
 in liver transplantation, antifibrinolytic agents for, 888–893  
 prediction of, after cardiac surgery, Thromboelastograph for, 531–535
- Coarctation, aortic, 429
- Cocaine abuse, aortic dissection in, 458–460
- Colloids, in normovolemic hemodilution, hemodynamic changes in, 812–818
- Compartment syndrome, abdominal, after congenital heart disease repair, 71–75
- Complement activation, in veno-veno bypass, 745
- Congenital heart disease  
 activated protein C in, 38–41  
 in adults, 414–437
- Congenital heart disease (*Continued*)  
 anatomy of, 414–415  
 anesthesia for, 429–434  
 categories of, 414  
 pathology of, 416–429  
 prognosis for, 414  
 shunts in, 415  
 arterial-to-end-tidal carbon dioxide tension difference in, 196–201  
 lobar emphysema with, 239–241  
 radial artery cannulation in, 48–50  
 repair of  
 air embolism after, 80–81  
 lactate level after, 43–47  
 phrenic nerve palsy after, 76–79  
 sepsis after, 71–75  
 in von Willebrand's disease, 576–579
- Contractility, estimation of, from carotid blood flow, 340–346
- Coronary artery(ies)  
 aneurysm of, 119–120  
 anomalous origin on, 429  
 dissection of, peripartum, 837–841  
 fistula of, with hypertrophic obstructive cardiomyopathy and mitral stenosis, 594–605  
 hemodynamics of, sevoflurane effects on, 209–216  
 occlusion of, during carotid endarterectomy, 259–268  
 spasm of, medical vs. surgical management of, 834–836
- Coronary artery bypass graft (CABG) surgery  
 autologous transfusions for, storage conditions and, 541–547  
 bleeding in  
 factor VIIa in, 573–575  
 prediction of, 531–535  
 carotid endarterectomy combined with  
 risk in, 796–802  
 staged anesthesia for, 803–806  
 thoracic epidural anesthesia for, 850–852  
 in diastolic dysfunction, isoflurane effects on, 509–514  
 in dissection, 837–841  
 endovascular vein harvesting in, 347–352  
 hemodynamic collapse after, prediction of, 143–148  
 heparinized oxygenators in, 554–561  
 kidney dysfunction after, 684–690  
 left atrial appendage thrombosis in, 96–97  
 for left main artery spasm, 834–836  
 after left pneumonectomy, 709–711  
 liver dysfunction after, 684–690  
 in low ejection fraction, drugs for, 353–357  
 myocardial infarction after, 781–787  
 normovolemic hemodilution in, hemodynamics in, 812–818  
 off-pump  
 anticoagulants for, thrombocytopenia due to, 106–111  
 cardiac output monitoring after, 315–319  
 carotid endarterectomy combined with, 850–852  
 preconditioning for, 477–483, 484–492  
 in prekallikrein deficiency, anticoagulation management in, 580–582  
 pulmonary artery catheter in, 300–306  
 temperature management during, 788–792  
 thoracic epidural anesthesia for, 656–658, 850–852  
 oxygenation index after, oxygen vs. nitrous oxide-oxygen mixture in, 136–142  
 preconditioning for, 473–476  
 opioids in, 493–502  
 sevoflurane in, 503–508

- Coronary artery bypass graft (CABG) surgery (*Continued*)  
 pulmonary artery catheter in, 88–89, 300–306, 307–310  
 statins in, 251–258
- C-reactive protein, in cardiac surgery, ketamine effects on, 217–222
- Creatine kinase-MB, as myocardial ischemia marker, in preconditioning, 503–508
- Creatinine, after CABG surgery, 684–690
- Crush injury, in explosions, 876
- Crystalloids  
 calcium supplementation in, coagulation profiles in, 807–811  
 in normovolemic hemodilution, hemodynamic changes in, 812–818  
 for volume replacement, 99
- Cyanosis, in congenital heart disease  
 activated protein C and, 38–41  
 adult, 414–437
- Cystic fibrosis, bronchial stenting in, with laryngeal mask airway, 227–228
- Cytokines, in CABG surgery, preconditioning effects on, 503–508
- Deep-brain stimulation, for Parkinson's disease, cardiothoracic surgery with, 707–708
- Desflurane, for preconditioning, 477–483
- Dexmedetomidine  
 for cardiac catheterization, in pediatric patients, 515–519  
 for sympathetic inhibition, myocardial oxygen consumption and, 364–370
- Dextrans, for volume replacement, 99
- Dextrocardia, pulmonary artery catheter position in, 456–457
- Diabetes mellitus, CABG surgery in, with in low ejection fraction, drugs for, 353–357
- Disasters, blast injury in, 872–880
- Dissection  
 aortic  
 in cocaine abuse, 458–460  
 repair of  
 bispectral index inaccuracy in, 68–70  
 bleeding control in, 736–741  
 femoral or axillary artery cannulation complications in, 20–25  
 coronary artery, peripartum, 837–841
- Diverticulum, Zenker's, transesophageal echocardiography in, 385–386
- Dobutamine stress echocardiography  
 in aortic valve stenosis, 862–866  
 in mitral valve assessment, 867–871
- Dopamine  
 for kidney failure, in septic shock, 173–178  
 for sevoflurane-induced left ventricular depression, 358–363
- Dopexamine, for kidney failure, in septic shock, 173–178
- Doppler studies  
 in cardiac output monitoring  
 in aortic repair, 26–30  
 after off-pump CABG, 315–319  
 in left ventricular function evaluation, 340–346
- Dyspnea, in pulmonary embolism, after heart transplantation, 236–238
- Ebstein's anomaly, 420–421
- Echocardiography  
 in diastolic dysfunction, isoflurane effects on, 509–514  
 in mitral valve repair, with left ventricular outflow tract obstruction, 842–846  
 stress, in aortic valve stenosis, 862–866  
 tissue Doppler, 583–593  
 transesophageal. *See* Transesophageal echocardiography
- Echo-esophageal Doppler studies, in cardiac output monitoring, in aortic repair, 26–30
- Edema, pulmonary, in Brugada syndrome, 407–413
- Editorials  
 Anesthetic Preconditioning: How Important Is It in Today's Cardiac Anesthesia?, 473–476  
 Is Supplemental Oxygen Necessary?, 133–135  
 Pulmonary Artery Catheter Utilization: The Use, Misuse, or Abuse, 295–299  
 The Time is Ripe, 765–767  
 Underutilization of Paravertebral Block in Thoracic Surgery, 635–638  
 The Value of Teamwork, 1–2
- Education, on thoracic epidural anesthesia, 652–655
- Elderly persons, cardiac surgery in, risk evaluation for, 526–530
- Electrocardiography, in Brugada syndrome, 407–413
- Electroencephalography, after cardiopulmonary bypass, 520–525
- Embolism  
 air  
 after congenital heart disease repair, 80–81  
 transesophageal echocardiography in, 387–389  
 cardiac, in liver transplantation, 232–235  
 pulmonary  
 after heart transplantation, 236–238  
 persistent, 616–619
- Emergencies, in air travel, 438–444
- Emphysema, lobar, in congenital heart disease, 239–241
- Enderectomy, carotid  
 CABG surgery combined with  
 risk in, 796–802  
 staged anesthesia for, 803–806  
 thoracic epidural anesthesia for, 850–852  
 cardiovascular collapse during, 259–268
- Endocardial cushion defect, 420
- Endocarditis, aortic valve repair in, persistent ST elevation after, 397–400
- Endothelial function, in internal mammary artery, in cardiopulmonary bypass, 63–67
- Endotracheal intubation  
 with tracheoesophageal fistula, 223–224  
 vocal cord paralysis due to, 225–226
- Endovascular repair, of aorta, left ventricular perforation in, 247–250
- Endovascular vein harvesting, for CABG surgery, 347–352
- Enflurane, for preconditioning, 473–476
- Epoprostenol, for pulmonary hypertension, vs. sildenafil, 730
- Esmolol, for sympathetic inhibition, myocardial oxygen consumption and, 364–370
- Esophageal Doppler studies, in cardiac output monitoring, in aortic repair, 26–30
- Esophagus  
 fistula of, endotracheal tube placement with, 223–224  
 injury of, in TEE, 379–384  
 perforation of, perioperative, TEE in, 896–897  
 Zenker's diverticulum of, TEE in, 385–386
- EuroSCORE model, for risk evaluation, of cardiac surgery, in octogenarians, 526–530
- Explosions, injury from, 872–880
- Factor VII, for bleeding control, in aneurysm repair, 570–572
- Factor VIIa, for bleeding control  
 in aortic repair, 736–741  
 in cardiac surgery, 573–575
- Femoral artery  
 cannulation of, for aortic dissection repair, complications of, 20–25

- Femoral artery (*Continued*)  
 pressure measurement in, for left ventricular function evaluation, 325–330
- Femoral vein, catheterization of, retroperitoneal hematoma in, 859–861
- Femorofemoral bypass, hemothorax after, TEE in, 229–231
- Fenoldopam, for ischemic kidney failure, 179–186
- Fentanyl  
 local anesthetics with, for paravertebral block, 644–647  
 for preconditioning, 493–502
- Fistula  
 coronary artery, with hypertrophic obstructive cardiomyopathy and mitral stenosis, 594–605  
 tracheoesophageal, endotracheal tube placement with, 223–224
- Fletcher factor (prekallikrein) deficiency, CABG surgery in, anticoagulation management in, 580–582
- Fluid therapy  
 for blast injury, 877  
 calcium supplementation in, coagulation profiles in, 807–811  
 normovolemic hemodilution in, hemodynamic changes in, 812–818
- Fogarty catheter, malposition of, in inferior vena cava thrombectomy, 405–406
- Fontan operation, 423–425
- Foramen ovale, patent, 419–420
- Forced-air system, in off-pump CABG surgery, 788–792
- Fractional oxygen concentration, after cardiopulmonary bypass, vs. gas composition, 136–142
- Gastrointestinal system, injury of, in TEE, 379–384
- General anesthesia  
 thoracic epidural anesthesia with, assessment of, 664–667  
 volatile agents for. *See* Preconditioning; *specific agents*
- Glutamate pyruvate transaminase, after CABG surgery, 684–690
- Glutamic oxaloacetic transaminase, after CABG surgery, 684–690
- Gore-Tex patch, for atrial septal defect repair, echocardiographic visualization of, 751–752
- Grafts, CABG surgery, endovascular harvesting of, 347–352
- Groin incision, lower limb ischemia due to, 187–195
- Heart  
 blast injury of, 875  
 leiomyomatosis of, excision of, TEE in, 94–95  
 swinging, pericardial effusion with, 242–244  
 transplantation of, pulmonary embolism after, 236–238
- Hematoma, retroperitoneal, in femoral vein cannulation, 859–861
- Hemodilution  
 in cardiopulmonary bypass, 156–161  
 moderate vs. tepid, oxygen saturation in, 819–825  
 normovolemic, hemodynamics in, 812–818
- Hemodynamics  
 of accessory tricuspid valve, in tetralogy of Fallot, 856–858  
 after CABG, prediction of, 143–148  
 of hemorrhagic shock, vasopressin effects on, 167–172  
 of lidocaine, in paravertebral block, 648–651  
 monitoring of  
 after off-pump CABG, 315–319  
 transesophageal echocardiography in, 768–771  
 in normovolemic hemodilution, 812–818  
 in percutaneous transcatheter closure of ventricular septal defect, 202–208  
 of sevoflurane, 209–216
- Hemoglobin levels, in congenital heart disease, 196–201
- Hemorrhage. *See* Bleeding
- Hemorrhagic shock, vasopressin effects on, 167–172
- Hemostatic material, spinal cord compression from, 696–699
- Hemothorax  
 in aneurysm, aortic repair with, 401–404  
 transesophageal echocardiography in, 229–231
- Heparin  
 in autologous blood transfusion, platelet dysfunction due to, 541–547  
 in CABG surgery, in prekallikrein deficiency, 580–582  
 in cardiopulmonary bypass, platelet function and, 554–561  
 thrombocytopenia induced by, 106–111
- Hepatic veins, air embolism in, TEE in, 387–389
- Hernia, hiatal, repair of, pericardial tamponade after, 245–246
- Hetastarch, for volume replacement, 99–102
- Hiatal hernia, repair of, pericardial tamponade after, 245–246
- Hydroxyethyl starches  
 calcium supplementation in, coagulation profiles in, 807–811  
 for volume replacement, 99–103
- Hypertension, pulmonary. *See* Pulmonary hypertension
- Hypertonic solutions, for volume replacement, 99
- Hypertrophic cardiomyopathy, obstructive, with mitral stenosis and coronary artery fistula, 594–605
- Hypokalemia, dilutional, in autologous transfusion, 568–569
- Hypotension, after pheochromocytoma resection, 394–396
- Hypothermia  
 correction of, in off-pump CABG surgery, 788–792  
 intentional  
 for aortic arch repair, 3–7  
 for aortic repair, kidney dysfunction after, 673–677  
 cerebral oxygen monitoring in, 826–833  
 moderate vs. tepid, oxygen saturation in, 819–825  
 for thoracic artery repair, 8–13  
 vascular response to thiopental in, 678–683
- Hypotonic solutions, for volume replacement, 99
- Hypovolemia, replacement therapy for, 98–105
- Hypoxia, cerebral, monitoring for, in cardiac surgery, 445–455, 606–615
- Iloprost, for pulmonary hypertension  
 in lung transplantation, 716–718  
 vs. sildenafil, 730
- Implantable cardioverter defibrillator, for Brugada syndrome, 407–413  
 in internal mammary artery, in cardiopulmonary bypass, 63–67
- Infants, congenital heart disease in  
 lobar emphysema with, 239–241  
 repair of  
 phrenic nerve palsy after, 76–79  
 sepsis after, 71–75
- Infections  
 after congenital heart disease repair, 71–75  
 methicillin-*Staphylococcus aureus*, in intensive care unit, 31–37
- Inflammation  
 in cardiac surgery, ketamine effects on, 217–222  
 of internal mammary artery, in cardiopulmonary bypass, 63–67
- Inflammatory mediators, in veno-veno bypass, 745
- Injury, blast, 872–880
- Innominate artery, pseudoaneurysm of, after internal jugular vein catheterization, 853–855
- Intensive care unit  
 methicillin-*Staphylococcus aureus* infections in, 31–37  
 prolonged stay in, after deep hypothermic circulatory arrest for thoracic artery repair, 8–13
- Interleukin(s)  
 in CABG surgery, preconditioning effects on, 503–508  
 in cardiac surgery, ketamine effects on, 217–222

- Intermittent positive-pressure ventilation, in tetralogy of Fallot, hemodynamics in, accessory tricuspid valve effects on, 856–858
- Intra-aortic balloon pump, cardiac output determination with, 320–324
- Ischemia  
 lower limb, in groin incision, 187–195  
 myocardial, in hemodynamic collapse, after CABG, 143–148  
 renal, fenoldopam for, 179–186
- Isoflurane  
 in diastolic dysfunction, 509–514  
 for preconditioning, 473–476
- Isotonic solutions, for volume replacement, 99
- Jatene operation, for transposition of great arteries, 428
- Jugular bulb venous saturation, in cerebral oxygen monitoring, 826–833
- Jugular vein, internal, catheterization of, innominate artery pseudoaneurysm in, 853–855
- Ketamine  
 anti-inflammatory effects of, in cardiac surgery, 217–222  
 for cardiac catheterization, in pediatric patients, 515–519  
 for cardiac surgery, neurologic outcome with, 520–525
- Kidney, dysfunction/failure of  
 after CABG surgery, 684–690  
 in deep hypothermic circulatory arrest  
 for aortic repair, 673–677  
 for thoracic artery repair, 8–13  
 fenoldopam for, 179–186  
 heparin-induced, 106–111  
 in septic shock, 173–178  
 in veno-veno bypass, 744
- Kyphoscoliosis, thoracic epidural anesthesia in, with multiple sclerosis, 704–706
- Lactate, prognostic significance of, after pediatric cardiac surgery, 43–47
- Lactate dehydrogenase, after CABG surgery, 684–690
- Landirolol, for tachycardia, 793–795
- Laryngeal mask airway, bronchial stenting with, 227–228
- Left atrial pressure  
 estimation of, tissue Doppler echocardiography in, 588–589  
 pulmonary artery catheter for, in cardiac surgery, 311–314
- Left sinus of Valsalva, aneurysm of, 276–279
- Legal issues, in air travel emergency care, 438–444
- Leiomyomatosis, intravenous, excision of, TEE in, 94–95
- Letters to the Editor  
 Addendum to cardiac arrest at 30,000 feet, 626–627  
 An unusual cause of high airway pressure and inadequate ventilation because of a defective connector despite accurate placement of a double-lumen tube, 627–630  
 Bispectral index in total circulatory arrest, 292–293  
 Cardiac output monitoring with brachial arterial catheter, 465–467  
 Combination of low-dose phenoxybenzamine and sodium nitropruside in children undergoing cardiac surgery, 291–292  
 Comparison of lumbar epidural tramadol and lumbar epidural morphine for pain relief after thoracotomy: a repeated-dose study, 757–759  
 Difficult intubation in pediatric cardiac surgery, 290–291  
 Echocardiography in suspected acute type A aortic dissection: detection and definitive management of a false-positive presentation, 912–914  
 Endoscopic saphenous vein harvest: severe hypercarbia, 759–760
- Letters to the Editor (*Continued*)  
 Evaluation of the supraglottic retractor for guided intraoperative transesophageal echocardiography probe insertion, 914–915  
 Fluid management in cardiac surgery: is one hydroxyethyl starch solution safer than another?, 916–917  
 High-frequency jet ventilation: experienced operators only, 763  
 Immediate extubation after cardiac surgery, 467–469  
 Intraoperative transesophageal ultrasonography can measure the left renal blood flow, 905–910  
 Management of major hemorrhage during mediastinoscopy, 762–763  
 Mediastinoscopy requiring a blood transfusion: unusual but significant, 469–471  
 Multimodal analgesia for chest tube removal after cardiac surgery, 760–761  
 Not all glucose clamps are equal, 630–631  
 Possible explanation for why blue blushed, 471–472  
 Pseudoaneurysm after attempted left internal venous cannulation, 290  
 Recombinant factor VIIa in intractable bleeding due to severe mediastinitis, 761–762  
 Resection of the Arndt bronchial blocker during stapler resection of the left lower lobe, 131–132  
 Resource utilization in on-and off-pump coronary artery surgery: factors influencing postoperative length of stay, 128–129  
 Severe acute tricuspid regurgitation after the withdrawal of an entrapped pulmonary artery catheter, 910–911  
 Silbronco double-lumen tube, 129–131  
 Total anomalous pulmonary venous connection in patient with situs inversus and dextrocardia: which internal jugular vein to cannulate, right or left?, 632–633  
 Use of cerebrospinal fluid drainage catheters during single-stage transmediastinal repair of ascending and descending aortic aneurysms, 126–127  
 Use of ketamine and midazolam cocktail for pericardiectomy at the University of Nigeria teaching hospital, 631–632
- Levosimendan  
 for low ejection fraction, in CABG surgery, 353–357  
 for right ventricular failure, after mitral valve replacement, 82–84
- Lidocaine, for paravertebral block, in one-lung ventilation, 648–651
- Liver  
 dysfunction of, after CABG surgery, 684–690  
 transplantation of  
 antifibrinolytics for, 888–893  
 cardiac thromboembolism in, 232–235  
 transfusion triggers in, 548–553  
 veno-veno bypass in, 742–747
- Low output syndrome, in diabetes mellitus, CABG surgery in, 353–357
- Lumbar epidural anesthesia, teaching practices for, 652–655
- Lung  
 blast injury of, 874–875  
 collapse of, 620–621  
 emphysema of, in congenital heart disease, 239–241  
 function of, in veno-veno bypass, 745  
 nodule resection from, thoracic epidural anesthesia for, subarachnoid injection of, 700–703  
 resection of  
 CABG surgery after, 709–711  
 paravertebral block for, 648–651  
 thoracic epidural anesthesia for, in multiple sclerosis, 704–706  
 vocal cord paralysis after, 225–226

- Lung (*Continued*)  
 transplantation of  
   bullae formation in, 748–750  
   cardiopulmonary bypass for, 668–672  
   pulmonary hypertension in, inhaled iloprost for, 716–718  
   pulmonary vein thrombosis in, 712–715  
   vascular bed of, meperidine effects on, 691–695  
 Lymphangioleiomyomatosis, lung transplantation for, bullae formation in, 748–750
- Mammary artery, internal, inflammation of, in cardiopulmonary bypass, 63–67
- Mass casualties, from blast injury, 872–880
- Mean systemic-to-pulmonary artery pressure ratio and, in cardiac surgery, 331–339
- Meperidine, pulmonary vascular effects of, 691–695
- Methicillin, *Staphylococcus aureus* resistant to, in intensive care unit, 31–37
- Midazolam, for cardiac surgery, neurologic outcome with, 520–525
- Milrinone  
   for low ejection fraction, in CABG surgery, 353–357  
   platelet-monocyte aggregation and, 162–166
- Mitral valve  
   assessment of, dobutamine stress echocardiography in, 867–871  
   repair of, sliding leaflet technique for, 842–846  
   replacement of, right ventricular failure after, levosimendan for, 82–84  
   stenosis of, with hypertrophic obstructive cardiomyopathy and coronary artery fistula, 594–605
- Monocytes, phosphodiesterase inhibitor effects on, 162–166
- Montreal Convention, on air travel emergency care, 442
- Morphine, for preconditioning, 493–502
- Mortality  
   in aortic repair, tracheostomy and, 14–19  
   in Brugada syndrome, 407–413  
   in cardiopulmonary bypass, for lung transplantation, 668–672  
   in combined CABG surgery and carotid endarterectomy, 796–802  
   in deep hypothermic circulatory arrest, for thoracic artery repair, 8–13  
   in hemodynamic collapse, after CABG, 143–148  
   in off-pump CABG surgery, 300–306  
   in pediatric cardiac surgery, lactate levels and, 43–47  
   prediction of, in cardiac surgery, in octogenarians, 526–530  
   in pulmonary artery catheter entrapment, 371–375
- Multiple sclerosis, lung resection in, thoracic epidural anesthesia for, 704–706
- Myocardial infarction  
   in carotid endarterectomy, 259–268  
   in combined CABG surgery and carotid endarterectomy, 796–802  
   perioperative, troponin detection of, 781–787
- Myocardial ischemia  
   in anesthetic preconditioning. *See* Preconditioning  
   after aortic valve repair, 397–400  
   in coronary artery dissection, 837–841  
   in hemodynamic collapse, after CABG, 143–148  
   isoflurane effects on, 509–514
- Myocardium  
   function of, tissue Doppler echocardiography for, 583–593  
   oxygen consumption of, in sympathetic stimulation, 364–370
- Near-infrared spectroscopy, in cerebral oxygen monitoring, 826–833
- Neonates  
   cardiopulmonary bypass in, in von Willebrand's disease, 576–579  
   congenital heart disease repair in, phrenic nerve palsy after, 76–79
- Neurologic disorders  
   after cardiopulmonary bypass  
     electroencephalography in, 520–525  
     near-infrared spectroscopy monitoring for, 606–615  
   in combined CABG surgery and carotid endarterectomy, 796–802, 803–806  
   after thoracotomy, in hemostatic material compression, 696–699  
 Neurostimulation, for Parkinson's disease, cardiothoracic surgery with, 707–708
- Neutrophils, in cardiopulmonary bypass, sevoflurane cardioplegia effects on, 57–62
- Nitric oxide, for pulmonary hypertension, vs. sildenafil, 730
- Nitric oxide synthase, in internal mammary artery, in cardiopulmonary bypass, 63–67
- Nitroglycerin, for coronary vasospasm, 834–836
- Nitrous oxide, after cardiopulmonary bypass, oxygenation index and, 136–142
- Norepinephrine, for septic shock, kidney failure treatment and, 173–178
- Normovolemic hemodilution, hemodynamic changes in, 812–818
- Nuclear factor kappa B, inhibition of, in kidney failure, 179–186
- Off-pump CABG. *See* Coronary artery bypass graft (CABG) surgery, off-pump
- Olprinone, for sevoflurane-induced left ventricular depression, 358–363
- Opioids. *See also* Morphine  
   for preconditioning, 493–502
- Optic neuropathy, nonarteritic ischemic, due to sildenafil, 733
- Oropharynx, injury of, in TEE, 379–384
- Outcomes  
   of aortic arch repair, with deep hypothermic circulatory arrest and retrograde cerebral perfusion, 3–7  
   of CABG surgery  
     with carotid endarterectomy, 796–802  
     in perioperative myocardial infarction, 781–787  
     pulmonary artery catheter effects on, 300–306, 307–310  
   of cardiopulmonary bypass  
     aortic atherosclerosis effects on, 149–155  
     for lung transplantation, 668–672  
     with retrograde autologous priming, 156–161  
   of surgery, statins effects on, 251–258
- Oximetry  
   cerebral  
     in cardiac surgery, 445–455, 606–615  
     continuous jugular bulb venous saturation in, 826–833  
     near-infrared spectroscopy in, 826–833  
     pulse, in congenital heart disease, 432
- Oxygen consumption, myocardial, in sympathetic stimulation, 364–370
- Oxygen tension, in congenital heart disease, 196–201
- Oxygen therapy  
   necessity of, 133–135  
   vs. nitrous oxide—oxygen mixture, oxygenation index and, 136–142
- Oxygenation  
   in cardiopulmonary bypass  
     in hypothermic hemodilution, 819–825  
     near-infrared spectroscopy monitoring of, 606–615  
     platelet function and, 554–561  
   cerebral, monitoring of  
     in cardiac surgery, 445–455, 606–615  
     continuous jugular bulb venous saturation in, 826–833  
     near-infrared spectroscopy in, 826–833
- Oxygenation index, after cardiopulmonary bypass, vs. gas composition, 136–142

- Pacing, intraoperative, 117–118
- Paralysis, vocal cord, after lung resection, 225–226
- Paraplegia, after thoracotomy, 696–699
- Paravertebral block
  - agents for, 644–647
  - anesthetics for, 637
  - multiple-injection, for video-assisted surgery, 639–643
  - in one-lung ventilation, hemodynamic effects of, 648–651
  - technique for, 635–636
  - underutilization of, 635–638
- Parkinson's disease, deep-brain stimulation for, cardiothoracic surgery with, 707–708
- Partial anomalous pulmonary venous drainage, 416
- Patent foramen ovale, 419–420
- Pediatric patients
  - cardiac catheterization in, sedation for, 515–519
  - congenital heart disease in
    - activated protein C in, 38–41
    - arterial-to-end-tidal carbon dioxide tension difference in, 196–201
    - lobar emphysema with, 239–241
    - radial artery cannulation in, 48–50
    - repair of
      - air embolism after, 80–81
      - lactate level after, 43–47
      - phrenic nerve palsy after, 76–79
      - sepsis after, 71–75
    - hypotension in, after pheochromocytoma resection, 394–396
    - pulmonary hypertension in
      - due to protamine, nitric oxide for, 719–721
      - sildenafil for, 731
- Percutaneous transcatheter closure, of ventricular septal defect, 202–208
- Perfusion, cerebral, with deep hypothermic circulatory arrest, for aortic arch repair, 3–7
- Pericardial effusion, with swinging heart, 242–244
- Pericardial reflection, central venous catheter placement and, 777–780
- Pericardial tamponade
  - in central venous catheter placement, 777–780
  - after hiatal hernia repair, 245–246
- Pericarditis, tissue Doppler echocardiography in, 589
- Pharynx, injury of, in TEE, 379–384
- Pheochromocytoma
  - intraoperative presentation of, 390–393
  - resection of, hypotension after, 394–396
- Phosphodiesterase inhibitors, platelet-monocyte aggregation and, 162–166
- Phrenic nerve palsy, after congenital heart disease repair, 76–79
- Plasma substitutes, 99–103
- Platelet(s)
  - dysfunction of
    - in autologous blood transfusion, 541–547
    - after cardiac surgery, prediction of, Thromboelastograph for, 531–535
    - in cardiopulmonary bypass, 554–561
      - aggregometry in, 536–540
    - phosphodiesterase inhibitor effects on, 162–166
- Plateletworks platelet-function analyzer, in cardiopulmonary bypass, 554–561
- Pleural cavity, catheter in, malposition into left atrium, 565–567
- Pneumectomy
  - left, CABG surgery after, 709–711
  - pleural drainage after, catheter malposition in, 565–567
- Positioning, for thoracic epidural anesthesia, 656–658
- Positive-pressure ventilation, intermittent, in tetralogy of Fallot, hemodynamics in, accessory tricuspid valve effects on, 856–858
- Postoperative pain
  - paravertebral block for
    - agents for, 644–647
    - in lung resection, 648–651
    - multiple-injection, 639–643
  - thoracic epidural anesthesia for
    - catheter position confirmation in, 659–663
    - in multiple sclerosis with kyphoscoliosis, 704–706
- Preconditioning, 473–476
  - isoflurane for, 509–514
  - multicenter study on, 477–483
  - for off-pump CABG surgery, 477–483, 484–492
  - opioids with, 493–502
  - sevoflurane for, 503–508
- Pregnancy
  - in congenital heart disease, 433
  - coronary artery dissection after, 837–841
- Prekallikrein deficiency, CABG surgery in, anticoagulation management in, 580–582
- Propofol
  - for cardiac catheterization, in pediatric patients, 515–519
  - with remifentanyl, for preconditioning, 484–492
- Prostaglandins, for lower limb ischemia prevention, 187–195
- Protamine, pulmonary hypertension due to, nitric oxide for, in pediatric patients, 719–721
- Protein C, activated, in congenital heart disease, 38–41
- Pseudoaneurysm, innominate artery, after internal jugular vein catheterization, 853–855
- Pulmonary artery
  - aneurysm of, transesophageal echocardiography in, 562–564
  - thromboendarterectomy of, thiopental for, 678–683
  - thrombosis of, persistent, 616–619
- Pulmonary artery catheter
  - arterial rupture due to, 376–378
  - in CABG surgery, 307–310
    - off-pump, 300–306
  - for cardiac output determination, intra-aortic balloon pump effects on, 320–324
  - in cardiac surgery
    - mean systemic-to-pulmonary artery pressure ratio and, 331–339
    - perforation by, 311–314
    - wedging effects on, 311–314
  - in congenital heart disease, 432
  - in dextrocardia, 456–457
  - entrapment of, morbidity and mortality in, 371–375
  - mechanical complications of, 85–87
  - right atrial perforation due to, 88–89
  - utilization of, 295–299
- Pulmonary artery pressure, meperidine effects on, 691–695
- Pulmonary edema, in Brugada syndrome, 407–413
- Pulmonary embolism, after heart transplantation, 236–238
- Pulmonary hypertension
  - in adult congenital heart disease, 419
  - in cardiac surgery, 331–339
  - due to protamine, nitric oxide for, in pediatric patients, 719–721
  - treatment of, 431, 722–735
- Pulmonary vascular resistance, sevoflurane effects on, 209–216
- Pulmonary vein, thrombosis of, in lung transplantation, 712–715
- Pulsatile pressure waveform, for catheter position confirmation, in thoracic epidural anesthesia, 659–663
- Pulse oximetry, in congenital heart disease, 432



- Pupillary reflex dilation, in thoracic epidural analgesia assessment, in general anesthesia, 664–667
- Pyopneumothorax, thoracotomy for, 696–699
- Radial artery cannulation, direct vs. guidewire-assisted, 48–50
- Remifentanyl, with sevoflurane or propofol, for preconditioning, 484–492
- Renin-aldosterone-angiotensin system, in volume control, 98
- Respiratory system, blast injury of, 874–875
- Resuscitation, in hemodynamic collapse, after CABG, 143–148
- Retrograde autologous priming, in cardiopulmonary bypass, 156–161
- Retroperitoneal hematoma, in femoral vein cannulation, 859–861
- Ringer's lactate
  - in normovolemic hemodilution, hemodynamic changes in, 812–818
  - for volume replacement, 99
- Risk analysis, for cardiac surgery, in octogenarians, 526–530
- Ropivacaine, for paravertebral block, for thoracotomy, 644–647
- ROTEM thromboelastometry analyzer, in liver transplantation, 548–553
- Saline solutions, for volume replacement, 99
- Saphenous vein, endovascular harvesting of, 347–352
- Sarcoma, pulmonary artery, 616–619
- Sedation, for cardiac catheterization, in pediatric patients, 515–519
- Selectins, phosphodiesterase inhibitors and, 162–166
- Sepsis
  - after aortic repair, 14–19
  - after congenital heart disease repair, 71–75
- Septic shock, treatment of, 173–178
- Sevoflurane
  - hemodynamic effects of, 209–216
  - kidney dysfunction due to, 684–690
  - left ventricular depression due to, 358–363
  - liver dysfunction due to, 684–690
  - for preconditioning, 473–476
    - cytokine levels and, 503–508
    - remifentanyl with, 484–492
- Shock
  - hemorrhagic, vasopressin effects on, 167–172
  - septic, treatment of, 173–178
- Sigmoidoscopy, air embolism from, TEE in, 387–389
- Sildenafil, for pulmonary hypertension, 722–735
  - animal models for, 725–726
  - vs. bosentan, 730
  - chemical structure of, 722–723
  - clinical efficacy of, 726–731
  - clinical trials of, 726–731
  - current status of, 733
  - in drug combinations, 727, 729–730
  - vs. epoprostenol, 730
  - hemodynamic studies of, 726
  - mechanism of action of, 722–724
  - vs. nitric oxide, 730
  - in pediatric patients, 731
  - pharmacodynamics of, 724
  - pharmacokinetics of, 724
  - postoperative, 731
  - rationale for, 724–725
  - safety of, 731–733
  - selective nature of, 722
- Sinus of Valsalva, ruptured, intraoperative detection of, 847–849
- Sliding leaflet technique, for mitral valve repair, with left ventricular outflow tract obstruction, 842–846
- Sonoclot analysis, for platelet function, in cardiopulmonary bypass, 536–540
- Spasm, left main coronary artery, medical vs. surgical management of, 834–836
- Spinal cord
  - compression of, from hemostatic material, 696–699
  - transesophageal echocardiography of, mistaken for aorta, 894–895
- Staphylococcus aureus* infections, methicillin-resistant, in intensive care unit, 31–37
- Starches, for volume replacement, 99–103
- Statins, effects on perioperative outcomes, 251–258
- Stenosis
  - aortic, dobutamine stress echocardiography in, 862–866
  - mitral, with hypertrophic obstructive cardiomyopathy and coronary artery fistula, 594–605
- Stents, bronchial, with laryngeal mask airway, 227–228
- Sternotomy, repeat, percutaneous superior vena cava cannulation for, 881–887
- Stomach, injury of, in TEE, 379–384
- Strain and strain rate, in tissue Doppler echocardiography, 584–586
- Stress echocardiography, dobutamine
  - in aortic valve stenosis, 862–866
  - in mitral valve assessment, 867–871
- Stroke
  - in combined CABG surgery and carotid endarterectomy, 796–802, 803–806
  - in deep hypothermic circulatory arrest
    - for aortic arch repair, 3–7
    - for thoracic artery repair, 8–13
- Subaortic aneurysm, 276–279
- Sudden death, in Brugada syndrome, 407–413
- Sufentanil, for cardiac surgery, neurologic outcome with, 520–525
- SURGICEL hemostatic material, spinal cord compression from, 696–699
- Swan-Ganz catheter, mechanical complications of, 85–87
- Swinging heart, pericardial effusion with, 242–244
- Sympathetic nervous system, myocardial oxygen consumption and, 364–370
- Syncope, in Brugada syndrome, 407–413
- Systemic-to-pulmonary artery pressure ratio and, in cardiac surgery, 331–339
- Tachycardia, landiolol for, 793–795
- Teaching, of thoracic epidural anesthesia, 652–655
- Teamwork, value of, 1–2
- TEE. *See* Transesophageal echocardiography
- Temperature management. *See also* Hypothermia, intentional
  - in off-pump CABG surgery, 788–792
- Terrorism, blast injury in, 872–880
- Tetralogy of Fallot, 425–427
  - accessory tricuspid valve in, hemodynamics of, in intermittent positive-pressure ventilation, 856–858
- Thermodilution catheter
  - for cardiac output determination, intra-aortic balloon pump effects on, 320–324
  - in cardiac output monitoring, in aortic repair, 26–30
- Thiopental, vascular response to, in deep hypothermia, 678–683
- Thoracic epidural anesthesia
  - catheter placement for, pulsatile pressure waveform confirmation of, 659–663
  - for combined CABG and carotid endarterectomy, 850–852
  - general anesthesia with, assessment of, 664–667
  - lumbar test dose for, 700–703
  - for lung surgery, in multiple sclerosis, 704–706

- Thoracic epidural anesthesia (*Continued*)  
 patient position for, 656–658  
 teaching practices for, 652–655
- Thoracic paravertebral block. *See* Paravertebral block
- Thoracoscopic surgery  
 paravertebral block for, multiple-injection, 639–643  
 for pulmonary nodule, thoracic epidural anesthesia for, subarachnoid injection of, 700–703
- Thoracotomy  
 paraplegia after, 696–699  
 paravertebral block for. *See* Paravertebral block  
 thoracic epidural anesthesia for  
 catheter position confirmation in, 659–663  
 general anesthesia with, assessment of, 664–667
- Thrombectomy, of inferior vena cava, left atrial catheter malposition in, 405–406
- Thrombocytopenia  
 aspirin for, 112–116  
 heparin-induced, 106–111
- Thromboelastograph  
 kaolin-activated, for bleeding prediction, after cardiac surgery, 531–535  
 for liver transplantation, 548–553
- Thrombosis  
 cardiac, in liver transplantation, 232–235  
 in congenital heart disease, activated protein C and, 38–41  
 coronary artery aneurysm, 119–120  
 left atrial appendage, TEE in, 96–97  
 in liver transplantation, antifibrinolytic agents for, 888–893  
 pulmonary artery, persistent, 616–619  
 pulmonary vein, in lung transplantation, echocardiography in, 712–715
- Tissue Doppler echocardiography, 583–593  
 clinical applications of, 587–589  
 image acquisition in, 586–587  
 principles of, 583–587  
 strain and strain rate in, 584–586
- Tracheobronchial stenting, with laryngeal mask airway, 227–228
- Tracheoesophageal fistula, endotracheal tube placement with, 223–224
- Tracheostomy, after aortic repair, 14–19
- Tranexamic acid, for liver transplantation, 888–893
- Transesophageal echocardiography  
 in aortic dissection  
 in cocaine abuse, 458–460  
 in repair, 20–25  
 in aortic root abscess, 280–283  
 in atrial septal defect repair, 90–93  
 benefits of, 765–767  
 in catheter malposition in left atrium, 405–406  
 in catheter removal from left atrium, 565–567  
 in esophageal perforation, 896–897  
 Gore-Tex patch visualized in, 751–752  
 in hemothorax, 229–231  
 of inferior vena cava, 284–285  
 intraoperative  
 in aortic valve replacement, 847–849  
 in noncardiac surgery, 768–771  
 in intravenous leiomyomatosis excision, 94–95  
 in left atrial appendage thrombosis detection, 96–97  
 in left atrial enlargement, 772–776  
 in left sinus of Valsalva aneurysm, 276–279  
 in pericardial effusion, 242–244  
 in pulmonary artery aneurysm, 562–564  
 in pulmonary artery thromboembolism, 616–619
- Transesophageal echocardiography (*Continued*)  
 in pulmonary vein thrombosis, 712–715  
 spinal cord misinterpretation in, 894–895  
 upper gastrointestinal injuries related to, 379–384  
 in valvular surgery, 772–776  
 in Zenker's diverticulum, 385–386
- Transfusion(s)  
 autologous  
 in CABG surgery, platelet function and, 554–561  
 for cardiopulmonary bypass, storage conditions and, 541–547  
 dilutional hypokalemia in, 568–569  
 blood substitutes for, 99–103  
 triggers for, in liver transplantation, 548–553  
 in veno-veno bypass, 745
- Transplantation  
 heart, pulmonary embolism after, 236–238  
 liver  
 antifibrinolytics for, 888–893  
 cardiac thromboembolism in, 232–235  
 transfusion triggers in, 548–553  
 veno-veno bypass in, 742–747  
 lung  
 bulla formation in, 748–750  
 cardiopulmonary bypass for, 668–672  
 pulmonary hypertension in, inhaled iloprost for, 716–718  
 pulmonary vein thrombosis in, 712–715  
 statins in, 251–258
- Transposition of great arteries, 428
- Transpulmonary thermodilution technique  
 for cardiac output determination, intra-aortic balloon pump effects on, 320–324  
 in hemodynamic evaluation, in paravertebral block, 648–651
- Transthoracic hiatal hernia repair, pericardial tamponade after, 245–246
- Tricuspid valve  
 accessory, in tetralogy of Fallot, hemodynamic changes due to, 856–858  
 defects of, in Ebstein's anomaly, 420–421
- Troponins  
 in CABG surgery  
 myocardial infarction and, 781–787  
 preconditioning effects on, 503–508  
 in preconditioning, 477–483
- Tubal ligation, coronary artery dissection during, 837–841
- Tuberculosis, pyopneumothorax in, thoracotomy for, 696–699
- Ultrasonic Cardiac Output Monitor, after off-pump CABG, 315–319
- Valvular heart disease. *See specific valves*
- Vascular endothelial growth factor, in internal mammary artery, in cardiopulmonary bypass, 63–67
- Vasoconstriction, from thiopental, in deep hypothermia, 678–683
- Vasopressin  
 for hemorrhagic shock, 167–172  
 for hypotension, after pheochromocytoma resection, 394–396
- Vasopressor, dependence on, in deep hypothermic circulatory arrest, for thoracic artery repair, 8–13
- Vasospasm, left main coronary artery, medical vs. surgical management of, 834–836
- Vena cava  
 inferior  
 thrombectomy of, left atrial catheter malposition in, 405–406  
 transesophageal echocardiography of, 284–285

Vena cava (*Continued*)

## superior

- catheterization of, in repeat sternotomy, 881–887
- central venous catheter placement in, 777–780
- persistence of, 416

Veno-veno bypass, in liver transplantation, 742–747

Ventilation, one-lung, paravertebral block in, hemodynamic effects of, 648–651

## Ventricle(s)

## left

- abnormal wall motion of, 397–400, 587
- after aortic valve repair, 397–400
- depression of, sevoflurane-induced, 358–363
- diastolic function of, tissue Doppler echocardiography in, 588
- elastance of, from carotid blood flow, 340–346
- function of, femoral artery pressure tracing for, 325–330
- hypertrophy of, isoflurane and, 509–514
- outflow tract obstruction, 427–428, 842–846
- perforation of, in aortic repair, 247–250
- relaxation indices of, isoflurane effects on, 509–514
- systolic function of, tissue Doppler echocardiography in, 587–588
- tissue Doppler echocardiography in, 587

Ventricle(s) (*Continued*)

## right

- failure of, after mitral valve replacement, levosimendan for, 82–84
- sevoflurane effects on, 209–216

Ventricular septal defect, 420–423

lobar emphysema with, 239–241

percutaneous transcatheter closure of, anesthesia for, 202–208

Video-assisted surgery, multiple-injection paravertebral block for, 639–643

Vocal cord paralysis, after lung resection, 225–226

Volatile anesthetics. *See* Preconditioning; *specific agents*

Volume therapy, in cardiac surgery, 98–105

blood substitutes for, in Europe vs. U.S., 99–103

principles of, 98–99

von Willebrand's disease, cardiopulmonary bypass in, in neonates, 576–579

White blood cell count, in atrial fibrillation prediction, 51–56

Zenker's diverticulum, transesophageal echocardiography in, 385–386