

- Ugurlucan, M., 816
 Uludag, E., 827
 Ulusoy, H.B., 696
 Unnikrishnan, M., 250
 Urovi, S., 847
- Valliattu, J., 906
 van der Linden, J., 602
 van der Maaten, J.M.A.A., 357
 van der Starre, P.J.A., 561, 584
 Vandrovec, S.M., 610
 van Hout, N., 351
 Van Wagoner, D.R., 68
 van Wezel, H.B., 406, 917
 Varma, P.K., 288
 Varshney, S.M., 51
 Veglia, F., 810
 Vellutini, R., 847
 Vender, J.S., 147, 475
 Vento, A.E., 224, 232, 529
 Verbrugge, L.B., 406
- Vetrugno, L., 96
 Vijayakumar, A., 478, 724, 876
- Wagener, G., 253
 Waikar, H.D., 625
 Wajima, Z., 793
 Walesby, R., 805
 Wang, H., 432
 Wasler, A., 384
 Wass, C.T., 720
 Watcha, S.M., 208
 Weiner, J., 388
 Weisenberg, M., 497
 Weiss, M., 535
 Weng, H., 774
 Whalen, F.X., 832
 Whitlock, R., 103
 Wiesner, G., 162
 Wolf, S.E., 593
 Wong, D.T., 774
- Woo, Y.J., 212
 Wu, C.L., 1
- Yaacoub, C., 432
 Yadav, A., 99
 Yared, J.-P., 68
 Yasny, J.S., 587
 Yildirim, K., 41
 Yildirim, V., 696
 Yildiz, O., 696
 Yoon, S.M., 678
 Yoon, S.-Z., 715
- Zaccaria, F., 174
 Zacharias, A., 820
 Zakhouri, M., 683
 Zangrillo, A., 502, 672, 761
 Zestos, M.M., 243
 Zingone, B., 166
 Zollinger, A., 524
 Zuchner, M.B., 618
 Zvara, D.A., 133, 257, 570

SUBJECT INDEX

- Abdominal pain, from migrating pacemaker, 906–907
 Activated coagulation time measurement, cardiopulmonary bypass, 524–528
 Acute lung injury, after pneumonectomy, 748–750
 Acute Physiology and Chronic Health Evaluation III, for pneumonectomy, 832–837
 Acute respiratory distress syndrome
 extracorporeal membrane oxygenation for, 384–387
 mechanical ventilation and, 279–284
 Adrenergic activity, of dexmedetomidine, on internal mammary artery, 696–700
 Aintree intubation catheter, for mediastinal mass resection with cardiac surgery, 259–261
 Air embolism, after cardiopulmonary bypass, 288–297
 Air-leak test, predictive value of, 197–202
 Airway abnormalities, undiagnosed, with tetralogy of Fallot, 876–879
 Airway exchange catheter, tracheal laceration due to, 718–719
 Airway obstruction, in facial trauma repair, 903–905
 Albumin, extravasation of, after cardiopulmonary bypass, 174–178
 Alfentanil, in cardiac surgery, inflammatory mediators and, 512–518
 Allergy, milk, aprotinin anaphylactic reaction in, 243–244
 Alveolar recruitment, after cardiac surgery, 375–378
 Amiodarone, for ventricular fibrillation, after aortic valve replacement, 414–416
 Anaphylaxis
 in aprotinin test dose, 243–244
 after negative aprotinin test dose, 597–601
 Ancrod, in history of heparin-induced thrombocytopenia, 114–115
 Anesthesia. *See also* Epidural anesthesia; Fast-track anesthesia
 general
 for carotid endarterectomy, vs. regional anesthesia, 127–132
 combined with spinal anesthesia, for anorectoplasty, 873–875
 high thoracic epidural anesthesia with, for CABG, 810–815
 regional, for carotid endarterectomy, vs. general anesthesia, 127–132
 spinal, combined with general anesthesia, for anorectoplasty, 873–875
 total intravenous, for CABG, 810–815
 AneuRx device, for aneurysm repair, 730, 732
- Aneurysm
 aortic
 repair of
 complications of, 763–765
 diastolic function assessment in, 486–491
 elephant trunk procedure, 584–586
 endovascular, 730–742
 subdural hematoma repair after, 704–708
 right atrial compression and displacement in, 152–154
 tracheal and bronchial compression with, 88–90
 sinus of Valsalva, 615–617
 Animal studies
 cardiac output methods, 659–663
 fluid responsiveness predicted by global end-diastolic volume, 650–654
 ischemic left ventricular dysfunction treatment with saline, 400–405
 lidocaine effects on hemodynamics, 393–399
 myocardial ischemia in increased contractility and end-diastolic wall tension, 8–17
 propofol effects in cold cardioplegia and reperfusion, 678–682
 thermodilution technique validity, 644–649
 thrombin inhibition during reperfusion, 224–231
 verapamil for reperfusion injury prevention, 337–343
 Anorectoplasty, combined general and spinal anesthesia for, 873–875
 Anterior spinal artery syndrome, after off-pump CABG surgery, 709–711
 Anticoagulants
 for heparin-induced thrombocytopenia, 113–126
 measurement of, in cardiopulmonary bypass, 524–528
 reversal of, after heart transplantation, 269–272
 thrombocytopenia induced by, after cardiac surgery, 18–22
 Antiplatelet agents, for percutaneous coronary interventions, 436–442
 Aorta
 aneurysm of
 repair of
 complications of, 763–765
 diastolic function assessment in, 486–491
 elephant trunk procedure, 584–586

- Aorta (*Continued*)
 endovascular, 730–742
 subdural hematoma repair after, 704–708
 right atrial compression and displacement in, 152–154
 tracheal and bronchial compression with, 88–90
 atherosclerosis of, intraoperative assessment of, 805–809
 compression of, from echocardiographic probe, 724–726
 dissection of
 aortic regurgitation with, echocardiography in, 85–87
 in cocaine abuse, 462–464
 repair of
 brachiocephalic malperfusion during, 81–84
 innominate artery cannulation in, 727–729
 surgery on, transesophageal echocardiography during, 800–804
 triple-lumen, in cocaine abuse, 462–464
- Aortic arch
 pseudoaneurysm of, in cocaine abuse, 462–464
 repair of
 deep hypothermic circulatory arrest for, atrial fibrillation after, 388–392
 intraoperative thrombosis in, 897–902
- Aortic valve
 regurgitation of
 in aortic dissection, echocardiography in, 85–87
 mitral valve prosthesis obscuring, 610–611
 after mitral valve replacement, 276–278
 pressure half-time alternans in, 150–151
 replacement of
 enoximone after, 357–366
 mediastinal mass resection with, 259–261
 mitral valve anterior motion after, 574–576
 myocardial function after, echocardiography in, 367–370
 rectus abdominis sheath hematoma after, 612–614
 transesophageal echocardiography during, 800–804
 ventricular fibrillation after, amiodarone for, 414–416
 stenosis of, surgery for, 197–202, 357–366, 367–370
- APACHE (Acute Physiology and Chronic Health Evaluation) III, for pneumonectomy, 832–837
- Apoptosis, cardiomyocyte, prevention of, in reperfusion, 224–231
- Aprotinin
 anaphylactic reaction to, 243–244, 597–601
 in cardiac surgery
 coronary thrombosis with, 558–560
 outcomes of, 23–27
 in coronary artery bypass graft surgery
 evidence against, 302–304
 evidence for, 298–301
 measurement of, in cardiopulmonary bypass, 524–528
- Argatroban, in history of heparin-induced thrombocytopenia, 115
- Arterial pressure, cardiac output measurement based on, 3–7
- Arterial pressure waveform analysis, for cardiac output, 629–631, 632–635, 636–643
- Arterial switch operation, albumin extravasation after, 174–178
- Arteries. *See specific arteries*
- Aspirin
 after cardiac surgery, ketorolac with, 820–826
 continuation of, for cardiac surgery
 evidence against, 606–609
 evidence for, 602–605
 for percutaneous coronary interventions, 436–442
- Atelectasis, after cardiac surgery, 375–378
- Atheroma, aortic, with dissection, 727–729
- Atherosclerosis, aortic, intraoperative assessment of, 805–809
- Atrial fibrillation
 after aortic arch repair, 388–392
 after cardiac surgery, dexamethasone effects on, 68–75
- Atrial septal defect, surgery for, air-leak test in, 197–202
- Atrium
 myxoma of, mitral valve regurgitation with, 103–105
 right
 avoidance of, in internal jugular vein catheterization, 497–501
 compression of, in aortic aneurysm, 152–154
 sinus of Valsalva aneurysm in, 615–617
 thrombus in, after off-pump CABG surgery, 564–566
- Axillary artery catheterization, for mitral valve replacement, pectoral muscle hematoma due to, 96–98
- Beta blockers, perioperative
 patient understanding of, 325–329
 survey on, 330–336
- Biopsy
 liver, cardiac tamponade in, 712–714
 mediastinal lymph node, ultrasound-guided, 892–896
- Bispectral index, in deep hypothermic circulatory arrest with retrograde cerebral perfusion, 61–67
- Bivalirudin, in history of heparin-induced thrombocytopenia, 118–120
- Biventricular assist device, implantation of, right ventricular drainage cannula obstruction in, 262–264
- Biventricular pacing, after cardiopulmonary bypass, 570–571
- Blalock-Taussig shunt, modified, hemodilution in, 179–183
- Bland-Altman analysis, in cardiac output measurement, 3–7
- Bleeding
 global end-diastolic volume and, 650–654
 in liver transplantation, 410–413
 prevention of, in emergency carotid endarterectomy, 851–854
- Blood flow
 cerebral, during hyperoxia, 212–217
 in saphenous vein grafts, vasoactive agents for, 344–350
- Blood loss, prevention of, with hemodilution, 179–183
- Blood patch, for spinal subdural hematoma, 704–708
- Blood pressure
 cardiac output measurement based on, 3–7
 in extubation, nicardipine vs. esmolol effects on, 45–50
 in undiagnosed pheochromocytoma, 858–862
- Brachial plexus block, interscalene, for post-thoracotomy shoulder pain, 554–557
- Brachiocephalic malperfusion, during aortic dissection repair, 81–84
- Brain, blood flow in, during hyperoxia, 212–217
- Bronchogenic cyst, perioperative risk with, 720–722
- Bronchoscopy, in mediastinal lymph node fine needle aspiration, 892–896
- Bronchus, compression of, in aortic aneurysm, 88–90
- Bupivacaine
 for interscalene brachial plexus block, for post-thoracotomy shoulder pain, 554–557
 intrapleural, for CABG, in chronic obstructive pulmonary disease, 816–819
- CABG surgery. *See* Coronary artery bypass graft (CABG) surgery
- Calcium channel blockers
 for hemodynamic control, in emergence and extubation, 45–50
 overdose of, pathophysiology of, 406–409
 for reperfusion injury prevention, 337–343
- Capillary leak syndrome, after cardiopulmonary bypass, in newborns, 174–178
- Carbon dioxide
 elevated, after cardiac surgery, 371–374

- Carbon dioxide (*Continued*)
 embolism of, in off-pump CABG surgery, 715–717
 mean expired, vs. arterial carbon dioxide levels, in cardiopulmonary bypass, temperature-related differences in, 57–60
 in partial carbon dioxide rebreathing method, 655–658
- Carcinoid syndrome, valvular heart disease caused by, 855–857
- Cardiac arrest, after cardiac surgery, due to pericardial effusion, 91–92
- Cardiac arrhythmias, intraoperative, 908–909
- Cardiac index, after cardiopulmonary bypass, biventricular pacing effects on, 570–571
- Cardiac output
 measurement of
 arterial pressure-based, 3–7
 FloTrac system for, 629–631, 632–635, 636–643
 partial carbon dioxide rebreathing method for, 655–658
 pulmonary artery vs. transpulmonary aortic methods for, 659–663
 promotion of, after cardiac surgery, 351–356
- Cardiac surgery. *See also specific procedures*
 air embolism after, 288–297
 air-leak test in, in pediatric patients, 197–202
 airway abnormalities in, 876–879
 albumin extravasation after, 174–178
 alveolar recruitment after, 375–378
 aortic compression in, from echocardiographic probe, 724–726
 aprotinin test dose for, 243–244, 597–601
 aspirin continuation for
 evidence against, 606–609
 evidence for, 602–605
 atrial fibrillation after, dexamethasone effects on, 68–75
 beta blocker importance in, 325–329
 cardiac output measurement after, FloTrac system for, 629–631, 632–635, 636–643
 cardiac output promotion after, 351–356
 clopidogrel continuation for
 evidence against, 606–609
 evidence for, 602–605
 colloids in
 evidence against, 457–459
 evidence for, 453–456
 coronary thrombosis in, with aprotinin, 558–560
 dental health management before, 587–591
 desflurane vs. propofol for, 672–677
 epiaortic echocardiography before, 425–427
 fast-track anesthesia for, 35–40
 hemodilution in, 179–183
 heparin-induced thrombocytopenia after
 alternative anticoagulants for, 113–126
 outcome of, 18–22
 hypercapnia after, 371–374
 hyperglycemia during, in nondiabetic patients, 843–846
 inferior vena cava diameter vs. central venous pressure in, 492–496
 isoflurane for, 683–689
 kallikrein inhibition in, outcomes of, 23–27
 kidney protection during, fenoldopam for, 847–850
 mechanical ventilation in, open-lung concept and, 279–284
 mediastinal mass resection with, 259–261
 in morbidly obese persons, 133–143
 myocardial function after, echocardiography in, 367–370
 neurologic disorders after, prevention of, evidence-based guidelines for, 529–534
 in Niemann-Pick type B disease, 428–431
 in osteogenesis imperfecta, 245–249
 oxygenation monitoring after, near-infrared spectroscopy in, 535–539
- Cardiac surgery. *See also specific procedures (Continued)*
 pain management after, parasternal intercostal block for, 547–553
 pericardial effusion after, cardiac arrest due to, 91–92
 perioperative interleukin levels in, in newborns, 184–190
 propofol for
 in extended cardioplegia and reperfusion, 678–682
 vs. sevoflurane, 664–671
 in pulmonary cripple, 443–452
 sevoflurane for, vs. propofol, 664–671
 stroke after, magnetic resonance imaging in, 106–109
 substernal epicardial echocardiography in, 237–242
 superior vena cava syndrome after, 417–419
 third-nerve palsy after, 110–112
 universal leukocyte reduction in, 232–236
 volatile anesthetics for, 502–511
- Cardiac tamponade, acute, unusual presentation of, 712–714
- Cardiomyopathy, dilated, pulmonary hypertension in, detection of, 51–56
- Cardiopulmonary bypass
 activated coagulation time measurement in, 524–528
 air embolism after, 288–297
 albumin extravasation after, in newborns, 174–178
 alveolar recruitment after, 375–378
 aprotinin for
 evidence against, 302–304
 evidence for, 298–301
 biventricular pacing after, 570–571
 electrocardiographic artifacts in, 572–573
 enflurane levels in, 218–223
 fast-track anesthesia in, 512–518
 for heart transplantation, anticoagulant reversal after, 269–272
 hyperoxia in, cerebral blood-flow response to, 212–217
 hyponatremia in, 273–275
 inflammatory mediator release in
 cell saver transfusions and, 519–523
 vs. temperature, 191–196
 kidney protection during, fenoldopam for, 847–850
 for lung transplantation, in cystic fibrosis, 208–211
 mean expired carbon dioxide in, vs. arterial carbon dioxide levels, temperature-related differences in, 57–60
 mechanical ventilation in, open-lung concept and, 279–284
 neurologic disorders after, prevention of, evidence-based guidelines for, 529–534
 perioperative interleukin levels in, in newborns, 184–190
 polymer-coated oxygenator in, for pediatric patients, 28–34
 reperfusion in, thrombin inhibition in, 224–231
 for thymoma, 867–869
- Cardiovascular collapse, due to contaminated red blood cell transfusion, 870–872
- Carotid artery
 intraoperative ultrasonography of, during aortic dissection repair, 81–84
 pseudoaneurysm of, after central venous access, endovascular covered stent for, 99–102
- Carotid endarterectomy
 coronary artery surgery with, wake-up test after, 540–546
 emergency, platelet inhibition reversal for, 851–854
 regional vs. general anesthesia for, 127–132
- Catheterization
 jugular vein
 external landmarks for, 497–501
 inferior thyroid artery injury in, 422–424
 malposition of, 465–467
 lumbar subarachnoid, complications of, 763–765

- Central venous access
 internal jugular vein, external landmarks for, 497–501
 pseudoaneurysm after, endovascular covered stent for, 99–102
- Central venous pressure, vs. inferior vena cava diameter, in cardiac surgery, 492–496
- Cerebrospinal fluid, drainage of, for aneurysm repair
 catheter retention in, 763–765
 subdural hematoma after, 704–708
- Charcot-Marie-Tooth disease, CABG surgery in, 567–569
- Chemotherapy, intrathoracic, for pseudomyxoma peritonei extension, 265–268
- Chronic obstructive pulmonary disease, cardiac surgery in, 443–452, 816–819
- Chylothorax, after pneumonectomy, 743–751
- Circulatory arrest, deep hypothermic. *See* Deep hypothermic circulatory arrest
- Clopidogrel
 continuation of, for cardiac surgery
 evidence against, 606–609
 evidence for, 602–605
 for percutaneous coronary interventions, 436–442
 reversal of, for emergency carotid endarterectomy, 851–854
- Coagulation, in cardiopulmonary bypass, polymer-coated oxygenator effects on, 28–34
- Cocaine abuse, aortic pathology in, 462–464
- Cognitive function
 after cardiopulmonary bypass, hyperoxia response and, 212–217
 after off-pump CABG, sevoflurane effects on, 690–695
- Colloids, for cardiac surgery
 evidence against, 457–459
 evidence for, 453–456
- Color-flow Doppler studies, of carotid artery, during aortic dissection repair, 81–84
- Complement, after cardiac surgery, dexamethasone effects on, 68–75
- Computed tomography, in rectus abdominis sheath hematoma, after aortic valve replacement, 612–614
- Congenital heart disease
 pulmonary hypertension in, sildenafil for, 203–207
 surgery for
 air-leak test in, 197–202
 albumin extravasation after, 174–178
 aortic compression during, 724–726
 aprotinin test dose for, anaphylactic reaction to, 243–244
 hemodilution in, 179–183
 polymer-coated oxygenator in, 28–34
 resistant hypoxemia in, 880–882
 undiagnosed airway abnormalities with, 876–879
- Continuous arterial spin labeling perfusion magnetic resonance imaging, in stroke, after cardiac surgery, 106–109
- Contractility, increased, myocardial ischemia in, 8–17
- Cooling phase, of cardiopulmonary bypass, carbon dioxide levels during, 57–60
- Coronary artery(ies), stents in, antiplatelet agents for, 436–442
- Coronary artery bypass graft (CABG) surgery
 aprotinin for
 evidence against, 302–304
 evidence for, 298–301
 carotid endarterectomy with, wake-up test after, 540–546
 in Charcot-Marie-Tooth disease, 567–569
 in chronic obstructive pulmonary disease, intrapleural analgesia for, 816–819
 coronary thrombosis in, with aprotinin, 558–560
 for elderly persons, 781–783, 784–792
 heparin-induced thrombocytopenia after, outcome of, 18–22
- Coronary artery bypass graft (CABG) surgery (*Continued*)
 high thoracic epidural anesthesia for, 810–815
 hyponatremia after, 273–275
 inflammatory mediator release in, vs. temperature, 191–196
 mediastinal mass resection with, 259–261
 in morbidly obese persons, 133–143
 off-pump
 carbon dioxide embolism in, 715–717
 in elderly persons, 784–792
 inflammatory mediator release in, cell saver transfusions and, 519–523
 intraoperative conversion to on-pump, 793–799
 isoflurane for, 690–695
 right atrial thrombus after, 564–566
 spinal artery syndrome after, 709–711
 transesophageal echocardiography during, 800–804
 pericardial effusion after, spinal cord ischemia due to, 91–92
 postoperative ketorolac in, mortality rate and, 820–826
 in pulmonary cripple, 443–452
 radial artery graft for, intravenous anesthetic effects on, 41–44
 third-nerve palsy after, 110–112
 transesophageal echocardiography during, 800–804
- Coronary sinus, dilated, with persistent left superior vena cava, 761–762
- Coronary vein, repair of, in off-pump CABG surgery, 715–717
- Cost analysis, of cardiac surgery, with off-pump to on-pump intraoperative conversion, 793–799
- C-reactive protein, after cardiac surgery, dexamethasone effects on, 68–75
- Cyst(s), bronchogenic, perioperative risk with, 720–722
- Cystic fibrosis, lung transplantation for, cardiopulmonary bypass in, 208–211
- Danaparoid, in history of heparin-induced thrombocytopenia, 115–116
- Deep hypothermic circulatory arrest
 for aortic arch repair, atrial fibrillation after, 388–392
 in osteogenesis imperfecta, 245–249
 with retrograde cerebral perfusion, electroencephalographic bispectral index and suppression ration during, 61–67
- Dental health management, before cardiac surgery, 587–591
- Desflurane
 for cardiac surgery, 502–511
 for mitral valve surgery, 672–677
- Desmopressin, for platelet inhibitor reversal, for emergency carotid endarterectomy, 851–854
- Dexamethasone
 for atrial fibrillation, after cardiac surgery, 68–75
 for mediastinal lymph node fine needle aspiration, 895
- Dexmedetomidine, adrenergic activity of, on internal mammary artery, 696–700
- Dialysis, venovenous, after cardiac surgery, 379–383
- Diaphragm, herniation of, 758–760
- Diastolic function, measurement of, transmitral flow velocity propagation velocity in, 483–485, 486–491
- Diclofenac, for post-thoracotomy shoulder pain, 554–557
- Dissection, aortic
 aortic regurgitation with, echocardiography in, 85–87
 in cocaine abuse, 462–464
 repair of
 brachiocephalic malperfusion during, 81–84
 innominate artery cannulation in, 727–729
- Dobutamine
 myocardial ischemia induced by, 8–17
 requirements for, isoflurane effects on, 683–689

- Double-lumen tube placement
 distal cuff inflation effects on, 838–842
 tracheal laceration in, 718–719
- Dutch Randomized Endovascular Aneurysm Management (DREAM) trial, 732–733
- Dyspnea, in reexpansion pulmonary edema, 887
- Echocardiography
 epiaortic, 425–427
 substernal epicardial, approach to, 237–242
 transesophageal. *See* Transesophageal echocardiography
- Economic issues, in cardiac surgery, with off-pump to on-pump intra-operative conversion, 793–799
- Edema, pulmonary, reexpansion, 887–891
- Editorials
 An Orphan Subspecialty, or Whither Pediatric Cardiac Anesthesia?, 171–173
 Cardiac Output From the Arterial Catheter: Deceptively Simple, 629–631
 Coronary Artery Bypass Surgery in the Very Old: Light at the End of a Tunnel or a Dead-End Road?, 781–783
 Measuring Diastolic Function With Transmitral Flow Propagation Velocity: More Than Just a Pretty Picture?, 483–485
 The Preoperative Evaluation: An Opportunity for Education and Other Strategies to Improve Care, 323–324
 Robotics and Telepresence—The Future Is Arriving Ahead of Schedule, 592
 Strengths and Limitations of Meta-analysis, 1–2
- Ejection fraction, thermodilution measurement of, 644–649
- Elastase, in transfusions, with cell saver vs. systemic blood, 519–523
- Elderly persons, CABG in, outcome of, 781–783, 784–792
- Electrocardiography, in cardiopulmonary bypass, artifacts in, 572–573
- Electroencephalography, in deep hypothermic circulatory arrest with retrograde cerebral perfusion, 61–67
- Elephant trunk procedure, for aortic aneurysm, 584–586
- Embolism
 air, after cardiopulmonary bypass, 288–297
 carbon dioxide, in off-pump CABG surgery, 715–717
- Emergence, hemodynamic response to, nicardipine vs. esmolol effects on, 45–50
- Endarterectomy, carotid
 coronary artery surgery with, wake-up test after, 540–546
 emergency, platelet inhibition reversal for, 851–854
 regional vs. general anesthesia for, 127–132
- End-diastolic wall tension, elevated, myocardial ischemia in, 8–17
- EndoFlex endotracheal tube, airway obstruction from, in facial trauma repair, 903–905
- EndoSure Wireless AAA Pressure Sensor, 735–736
- Endothelin-1, after cardiac surgery, dexamethasone effects on, 68–75
- Endotracheal tube
 air-leak test of, predictive value of, 197–202
 airway obstruction from, in facial trauma repair, 903–905
 double-lumen, tracheal laceration from, 718–719
 for mediastinoscopy, cuff perforation during, 723
 for one-lung ventilation, in pneumonectomy, 883–886
- Endovascular covered stent, for pseudoaneurysm, 99–102
- Endovascular techniques
 for aortic aneurysm, 730–742
 for carotid pseudoaneurysm, 99–102
- Enflurane, in hypothermic cardiopulmonary bypass, 218–223
- Enoximone, left ventricular diastolic function effects of, after aortic valve replacement, 357–366
- Epiaortic echocardiography, 425–427
- Epiaortic ultrasonography, in aortic atherosclerosis assessment, 805–809
- Epicardial echocardiography
 in left ventricular thrombus, 305–306
 substernal, approach to, 237–242
- Epidural anesthesia
 acute spinal artery syndrome in, 709–711
 in cardiac surgery, inflammatory mediators and, 512–518
 drug-eluting stents and, 701–703
 for post-thoracotomy shoulder pain, 554–557
- Epidural blood patch, for spinal subdural hematoma, 704–708
- Epinephrine
 for cardiac output promotion, after cardiac surgery, 351–356
 for saphenous vein graft blood flow, 344–350
- Epoprostenol-heparin, in history of heparin-induced thrombocytopenia, 121
- Esmolol
 for hemodynamic control, in emergence and extubation, 45–50
 myocardial ischemia induced by, 8–17
- Etomidate
 for mediastinal lymph node fine needle aspiration, 894
 for radial artery graft spasm prevention, 41–44
- Excluder device, for aneurysm repair, 730, 732
- Extracorporeal membrane oxygenation, outcomes of, 384–387
- Extubation
 air-leak test before, predictive value of, 197–202
 in fast-track anesthesia, 35–40
 hemodynamic response to, nicardipine vs. esmolol effects on, 45–50
 after liver transplantation, timing of, 752–757
 after off-pump CABG, sevoflurane effects on, 690–695
- Facial trauma, subtotal airway obstruction in, 903–905
- Factor VII, activated, intraoperative thrombosis due to, 897–902
- Factor VIII inhibitor bypass activity administration, thromboses after, 561–563
- Fast-track anesthesia
 acute spinal artery syndrome in, 709–711
 for cardiac surgery
 inflammatory mediators and, 512–518
 sevoflurane and, 690–695
 for liver transplantation, 752–757
- Fenoldopam, for kidney protection, in cardiac surgery, 847–850
- Fentanyl, pericardial lidocaine with, 393–399
- Fibrinolysis
 in cardiopulmonary bypass, polymer-coated oxygenator effects on, 28–34
 in liver transplantation, 410–413
- Fine needle aspiration, of mediastinal lymph nodes, ultrasound-guided, 892–896
- FloTrac system, for cardiac output measurement, 629–631, 632–635, 636–643
- Fluid responsiveness, global end-diastolic volume and, 650–654
- Fortron device, for aneurysm repair, 730, 732
- General anesthesia
 for carotid endarterectomy, vs. regional anesthesia, 127–132
 combined with spinal anesthesia, for anorectoplasty, 873–875
 high thoracic epidural anesthesia with, for CABG, 810–815
- Global end-diastolic volume, fluid responsiveness and, 650–654
- Glucose elevation, in cardiac surgery, in nondiabetic patients, 843–846
- Guidewires, misplaced, transesophageal echocardiography of, 420–421
- Heart. *See also* Cardiac surgery; *specific procedures*
 hypokinesia of, due to undiagnosed pheochromocytoma, 863–866
 metastasis to, leiomyosarcoma, 432–433
 transplantation of

- Heart. *See also* Cardiac surgery; *specific procedures (Continued)*
 heparin-induced thrombocytopenia in, 269–272
 propofol for, in extended cardioplegia and reperfusion, 678–682
 pulmonary hypertension detection before, 51–56
 transesophageal echocardiography during, 800–804
- Heart rate
 in extubation, nicardipine vs. esmolol effects on, 45–50
 pericardial lidocaine effects on, 393–399
- Heart-liver transplantation, 253–256
- Hematoma
 mammary artery, in stab injury, 307–309
 pectoral muscle, after axillary artery catheterization, for mitral valve replacement, 96–98
 rectus abdominis sheath, after aortic valve replacement, 612–614
 subdural, after spinal catheter drainage, 704–708
- Hemodialysis, venovenous, after cardiac surgery, 379–383
- Hemodilution, in modified Blalock-Taussig shunt operation, 179–183
- Hemodynamics
 after aortic valve replacement, 357–366
 after cardiac surgery, hypercapnia and, 371–374
 after cardiopulmonary bypass, biventricular pacing effects on, 570–571
 of extubation, nicardipine vs. esmolol effects on, 45–50
 pericardial lidocaine effects on, 393–399
- Hemofiltration, venovenous, for anticoagulant reversal, after heart transplantation, 269–272
- Hemolysis, in transfusions, with cell saver vs. systemic blood, 519–523
- Hemorrhage
 global end-diastolic volume and, 650–654
 in liver transplantation, 410–413
 prevention of, in emergency carotid endarterectomy, 851–854
- Heparin
 low-molecular weight, in history of heparin-induced thrombocytopenia, 114
 thrombocytopenia induced by
 acute, cardiac surgery with, 122–123
 after cardiac surgery, 18–22
 after heart transplantation, 269–272
 history of, alternative anticoagulants for, 113–126
 subacute, cardiac surgery with, 123
- Hereditary motor and sensory neuropathy (Charcot-Marie-Tooth disease), CABG surgery in, 567–569
- Herniation, intestinal, into mediastinum, 758–760
- Hirudin, for thrombin inhibition, during reperfusion, 224–231
- Horseshoe lung, in pneumonectomy, 250–252
- Hydropneumopericardium, in pediatric patients, 758–760
- Hydroxyethyl starch
 for cardiac surgery
 evidence against, 457–459
 evidence for, 453–456
 hemodilution with, in modified Blalock-Taussig shunt operation, 179–183
- Hypercapnia, after cardiac surgery, 371–374
- Hyper eosinophilic syndrome, prosthetic valve thrombus in, 434–435
- Hyperglycemia, in cardiac surgery, in nondiabetic patients, 843–846
- Hyperoxia, in cardiopulmonary bypass, cerebral blood-flow response to, 212–217
- Hypertension, pulmonary
 in cardiac surgery, on obese persons, 137–140
 detection of, before heart transplantation, 51–56
 sildenafil for, 203–207
- Hypertensive crisis, due to undiagnosed pheochromocytoma, 858–862
- Hyperthermia, for pseudomyxoma peritonei extension chemotherapy, 265–268
- Hyponatremia, in cardiopulmonary bypass, 273–275
- Hypoplastic left-heart syndrome, correction of
 air-leak test in, 197–202
 perioperative interleukin levels in, 184–190
- Hypothermia. *See also* Deep hypothermic circulatory arrest
 in cardiopulmonary bypass
 enflurane levels and, 218–223
 inflammatory mediator release and, 191–196
 circulatory arrest in, with retrograde cerebral perfusion, electroencephalographic bispectral index and suppression ratio during, 61–67
- Hypoxemia
 prevention of, after cardiac surgery, 375–378
 with Sano shunt, 880–882
- Idiopathic hyper eosinophilic syndrome, prosthetic valve thrombus in, 434–435
- Iloprost-heparin, in history of heparin-induced thrombocytopenia, 120–121
- Infants. *See also* Newborns
 reexpansion pulmonary edema in, 889
 Sano shunt in, resistant hypoxemia in, 880–882
- Infections
 dental, management of, before cardiac surgery, 587–591
 reduction of, universal leukocyte reduction for, 232–236
- Inflammatory mediators
 in cardiac surgery, fast-track anesthesia and, 512–518
 release of, in cardiopulmonary bypass, vs. temperature, 191–196
 in transfusions, with cell saver vs. systemic blood, 519–523
- Innominate artery, cannulation of, in severe aortic dissection repair, 727–729
- Insulin, for verapamil overdose, 406–409
- Intensive care unit
 APACHE III evaluation use in, after pneumonectomy, 832–837
 electrocardiographic artifacts in, 572–573
 telemedicine application for, 592–596
 tracheostomy practice in, 76–80
- Intercostal block, parasternal, for postoperative pain, 547–553
- Interleukin(s)
 after cardiac surgery, dexamethasone effects on, 68–75
 in cardiopulmonary bypass, 184–190
 in transfusions, with cell saver vs. systemic blood, 519–523
- Interscalene brachial plexus block, for post-thoracotomy shoulder pain, 554–557
- Intestinal herniation, into mediastinum, 758–760
- Intra-aortic counterpulsation
 for aortic regurgitation, 150–151
 cardiac output measurement in, 636–643
- Intrapleural analgesia, for CABG, in chronic obstructive pulmonary disease, 816–819
- Intravenous anesthetics
 in cardiac surgery, inflammatory mediators and, 512–518
 in radial artery graft spasm prevention, 41–44
- Intubation. *See* Endotracheal tube
- Ischemia
 myocardial. *See* Myocardial ischemia
 spinal cord, due to pericardial effusion, 91–92
- Isoflurane
 for off-pump CABG surgery, 690–695
 for preconditioning, in cardiac surgery, 683–689
- Jugular vein, internal, catheterization of
 external landmarks for, 497–501
 inferior thyroid artery injury in, 422–424
 malposition of, 465–467

- Kallikrein inhibitors, in cardiac surgery, outcomes of, 23–27
- Kaolin-based activated coagulation time measurement, cardiopulmonary bypass, 524–528
- Ketamine, in radial artery graft spasm prevention, 41–44
- Ketorolac, after CABG, mortality rate and, 820–826
- Kidney, protection of, in cardiac surgery, fenoldopam for, 847–850
- Kirklin, John, Moffitt collaboration with, 285–287
- Klebsiella pneumoniae*, red blood cells contaminated with, cardiovascular collapse due to, 870–872
- Laryngeal mask airway, for mediastinal mass resection with cardiac surgery, 259–261
- Laryngomalacia, undiagnosed, tetralogy of Fallot with, 876–879
- Left ventricular end-diastolic area, fluid responsiveness and, 650–654
- Leiomyosarcoma, right ventricular, 432–433
- Lepirudin
 - in history of heparin-induced thrombocytopenia, 116–118
 - reversal of, after heart transplantation, 269–272
- Letters to the Editor
 - Acute Back Pain and Paresthesia After Femoral Venous Catheter Placement, 317–318
 - Additional Information on the Pulsatile Pressure Waveform, 920–921
 - An Aid to Awake Bronchoscopic Nasal Intubation, 476–477
 - Anesthesiologists' Placement of Cannulae for Cardiopulmonary Bypass, 919
 - Another Technique of Internal Jugular Vein Catheterization, 319–320
 - Aortic Valve Replacement in an Adult White Male With Moyamoya Disease and Coronary Artery Fistulas, 166–168
 - Bispectral Index Monitoring During Aortic Arch Repair, 479–480
 - Caution When Positioning Central Venous Catheters and Its Relationship to the Pericardial Reflection, 923–924
 - Central Venous Catheter: A Journey Through the Central Vein—Destination Pulmonary Vein, 625–626
 - Diaphragm Paralysis in Children after Cardiac Surgery, 164–165
 - Esophageal Placement of an Airway Exchange Catheter, 773–774
 - Fish Baffles Doctors: Scombroid Fish Poisoning of Fish Allergy?, 921–923
 - GlideScope-Assisted Endotracheal Tube Exchange, 777
 - GlideScope-Assisted Insertion of a Transesophageal Echocardiography Probe, 628
 - Identifying Imminent Displacement of a Double-Lumen Tube Caused by Surgical Traction at the Pulmonary Hilum, 776–777
 - Is Routine Placement of a Pulmonary Artery Catheter Essential?, 480–481
 - Myocardial Effects of Protamine, 779
 - Opioids and Cardioprotection: The Impact of Morphine and Fentanyl on Recovery of Ventricular Function After Cardiopulmonary Bypass, 474–476
 - Paraplegia Because of Hemostatic Agents in the Costovertebral Space: This Occurs Even in Thoracic Aorta Surgery, 624–625
 - Peripheral Nerve Blocks for High-Risk Care, 780
 - Positioning of Double-Lumen Tubes by a Lighted Stylet, 774–775
 - The Possible Mechanism of Apnea After Institution of Cardiopulmonary Bypass in Conscious Patients, 316–317
 - Rapid Recovery of Left-Ventricular Function After Intraoperative Conversion of Atrial Fibrillation, 160–164
 - Reverse Sellick's Maneuver for Transesophageal Echocardiographic Probe Placement, 626–627
 - Safety Concerns About Perisurgical Epoetin Treatment: Should We Add Antiplatelet Drugs to Perisurgical Epoetin Alfa Treatment?, 623–624
- Letters to the Editor (*Continued*)
 - Secondary Pericarditis Caused by Hydatid Cyst Rupture: Perioperative Transesophageal Echocardiography for the Assessment of Surgical Treatment, 777–778
 - Storage Lesion in Intraoperative Autologous Whole Blood, 473–474
 - Success Rate, Decubitus Position, and Vagal Reflex During a High Thoracic Epidural for Coronary Artery Surgery, 919–920
 - Transpharyngeal Imaging of the Carotid Artery and Internal Jugular Vein: Possible Roles in Cerebral Perfusion Management During Adult Aortic Arch Repair, 318–319
 - Unrecognized Nasal Foreign Body: Potential Consequences, 478–479
 - Unusual Embolism of Intracardiac Thrombus to a Femoral Artery, 168–169
 - The Use of Rigid Bronchoscopy for Bronchial Stenting in Patients With Tracheal Stenosis, 320–321
 - What Are the Clinical Questions for Optimal Conduct of Deep Hypothermic Circulatory Arrest for Adult Aortic Arch Repair?, 918–919
- Leukocytes, reduction of, in cardiac surgery, 232–236
- Lidocaine, pericardial, hemodynamic effects of, 393–399
- Liver
 - biopsy of, cardiac tamponade in, 712–714
 - function of, in cardiac output promotion, after cardiac surgery, 351–356
 - transplantation of
 - cardiovascular collapse in, contaminated red blood cell transfusion and, 870–872
 - extubation timing in, 752–757
 - fibrinolysis during, 410–413
 - left ventricular outflow tract obstruction during, 577–580
- Liver-heart transplantation, 253–256
- Lung
 - adenocarcinoma of, pneumonectomy for, complications of, 743–751
 - alveolar recruitment in, after cardiac surgery, 375–378
 - collapse of, reexpansion after, pulmonary edema in, 887–891
 - dysfunction of, cardiac surgery in, 443–452
 - horseshoe, in pneumonectomy, 250–252
 - inflammation of, in mechanical ventilation, 279–284
 - resection of. *See* Pneumonectomy
 - separation of, double-lumen tube placement for, 838–842
 - transplantation of, for cystic fibrosis, cardiopulmonary bypass in, 208–211
 - water in, after cardiac surgery, 379–383
- Lymph nodes, mediastinal, fine needle aspiration of, ultrasound-guided, 892–896
- Lymphoma, intraoperative discovery of, 257–258
- Magnesium sulfate, morphine with, after thoracotomy, 827–831
- Magnetic resonance imaging
 - in cerebral hyperoxia, in cardiopulmonary bypass, 212–217
 - in stroke, after cardiac surgery, 106–109
- Mammary artery
 - hematoma of, in stab injury, 307–309
 - internal, dexmedetomidine effects on, 696–700
- Mechanical ventilation
 - for cardiac surgery
 - kallikrein inhibitor effects on, 23–27
 - open-lung concept and, 279–284
 - one-lung, for pneumonectomy, 883–886
 - for reexpansion pulmonary edema, 890
 - Mediastinoscopy, endotracheal tube cuff perforation during, 723

- Mediastinum
 intestinal herniation into, 758–760
 lymph nodes of, fine needle aspiration of, ultrasound-guided, 892–896
 masses in
 intraoperative discovery of, 257–258
 resection of, with cardiac surgery, 259–261
 widened, in mammary artery hematoma, 307–309
- Meta-analysis, 1–2
- Midazolam, pericardial lidocaine with, 393–399
- Milk allergy, aprotinin anaphylactic reaction in, 243–244
- Milrinone, for saphenous vein graft blood flow, 344–350
- Mitral valve
 anterior motion of, after aortic valve replacement, 574–576
 flow measurement in, for diastolic function measurement, 483–485, 486–491
 prolapse of, regurgitant jet direction in, 581–583
 prosthesis for, obscuring aortic regurgitation, 610–611
 reconstruction of, desflurane vs. propofol for, 672–677
 replacement of
 air embolism after, 288–297
 aortic insufficiency after, 276–278
 axillary artery catheterization for, pectoral muscle hematoma due to, 96–98
 emergent reoperation after, 93–95
 transesophageal echocardiography during, 800–804
- Mixed venous oxygen saturation, monitoring of, near-infrared spectroscopy in, 535–539
- Moffitt, Emerson, cardiac anesthesia pioneer, 285–287
- Morphine, magnesium sulfate with, after thoracotomy, 827–831
- Mortality
 in aortic arch repair, due to thrombosis, 897–902
 in cardiac surgery
 in elderly persons, 781–783, 784–792
 ketorolac effects on, 820–826
 with propofol vs. sevoflurane, 664–671
 in carotid endarterectomy, regional vs. general anesthesia and, 127–132
 in intracardiac thrombus, 561–563
 in liver transplantation, due to contaminated red blood cell transfusion, 870–872
 in reexpansion pulmonary edema, 888–889
- Multiple organ failure, due to undiagnosed pheochromocytoma, 863–866
- Muscle relaxants, for mediastinal lymph node fine needle aspiration, 894–895
- Myocardial dyssynchrony, after cardiopulmonary bypass, biventricular pacing for, 570–571
- Myocardial infarction
 after carotid endarterectomy, regional vs. general anesthesia and, 127–132
 prevention of, beta blockers for, 325–329
- Myocardial ischemia
 in increased contractility, 8–17
 prevention of
 during reperfusion, 224–231
 in reperfusion injury, 337–343
 propofol effects on, 678–682
 saline solution for, 400–405
- Myocardial reperfusion, thrombin inhibition during, 224–231
- Myocardial stunning, saline solution for, 400–405
- Myocardium, function of, after aortic valve replacement, 367–370
- Myxoma, atrial, mitral valve regurgitation with, 103–105
- Near-infrared spectroscopy, for oxygenation monitoring, 535–539
- Neonates. *See* Newborns
- Neurologic disorders. *See also* Stroke
 after cardiac surgery
 oculomotor nerve, 110–112
 prevention of, evidence-based guidelines for, 529–534
 stroke, 106–109
 after cardiopulmonary bypass, hyperoxia response and, 212–217
- Neuron-specific enolase, in transfusions, with cell saver vs. systemic blood, 519–523
- Newborns
 anorectoplasty in, combined general and spinal anesthesia for, 873–875
 cardiac surgery in, perioperative interleukin levels in, 184–190
 cardiopulmonary bypass for, albumin extravasation after, 174–178
- Nicardipine, for hemodynamic control, in emergence and extubation, 45–50
- Niemann-Pick type B disease, cardiac surgery in, 428–431
- Nitric oxide
 inhaled, in pulmonary hypertension detection, before heart transplantation, 51–56
 for pulmonary hypertension, 203–207
- Nitrous oxide, double-lumen tube placement for, distal cuff inflation effects on, 838–842
- Norwood procedure, modified, resistant hypoxemia in, 880–882
- Obesity, morbid, cardiac surgery in, 133–143
- Octaplas, in liver transplantation, fibrinolysis due to, 410–413
- Octogenarians, CABG in, outcome of, 781–783, 784–792
- Oculomotor nerve palsy, after cardiac surgery, 110–112
- Off-pump CABG. *See* Coronary artery bypass graft (CABG) surgery, off-pump
- One-lung ventilation
 cardiac output measurement in, partial carbon dioxide rebreathing method for, 655–658
 tracheal laceration in, 718–719
- Open-lung concept, in mechanical ventilation, 279–284
- Oral health, optimizing, before cardiac surgery, 587–591
- Osteogenesis imperfecta, deep hypothermic circulatory arrest in, 245–249
- Outcomes
 of aortic arch repair, 388–392
 of cardiac surgery
 in elderly persons, 781–783, 784–792
 with hemodilution, 179–183
 with heparin-induced thrombocytopenia, 18–22
 with kallikrein inhibition, 23–27
 neurologic status, 529–534
 in off-pump to on-pump intraoperative conversion, 793–799
 with propofol vs. sevoflurane, 664–671
 with volatile anesthetics, 502–511
 of carotid endarterectomy, with regional vs. general anesthesia, 127–132
 of lung transplantation, for cystic fibrosis, 208–211
 of universal leukocyte reduction, 232–236
- Oxygen, elevated, in cardiopulmonary bypass, cerebral blood-flow response to, 212–217
- Oxygenation, monitoring of, near-infrared spectroscopy in, 535–539
- Oxygenator, for cardiopulmonary bypass, polymer-coated, for pediatric patients, 28–34
- Pacemaker, migrating to abdomen, 906–907
- Pacing, biventricular, after cardiopulmonary bypass, 570–571
- Paclitaxel, stents eluting, 436–442

- Palpation, digital, in aortic atherosclerosis assessment, 805–809
- Parasternal intercostal block, for postoperative pain, 547–553
- Partial carbon dioxide rebreathing method, for cardiac output measurement, 655–658
- Patient education, on beta blockers, 325–329
- Patient-controlled analgesia, magnesium sulfate with morphine in, 826–831
- Pectoral muscle hematoma, after axillary artery catheterization, for mitral valve replacement, 96–98
- Pediatric patients
 - as anesthesia specialty, 171–173
 - anorectoplasty in, combined general and spinal anesthesia for, 873–875
 - aprotinin anaphylactic reaction in, 243–244
 - cardiac surgery in
 - air-leak test in, 197–202
 - aortic compression in, from echocardiographic probe, 724–726
 - modified Blalock-Taussig shunt operation in, hemodilution in, 179–183
 - perioperative interleukin levels in, 184–190
 - resistant hypoxemia in, 880–882
 - cardiopulmonary bypass for
 - albumin extravasation after, 174–178
 - polymer-coated oxygenator in, 28–34
 - migrating pacemaker in, 906–907
 - pulmonary hypertension in, sildenafil for, 203–207
 - reexpansion pulmonary edema in, 889
 - tetralogy of Fallot in, undiagnosed airway abnormalities with, 876–879
- Percutaneous coronary interventions, antiplatelet agents for, 436–442
- Pericardial effusion
 - after cardiac surgery, cardiac arrest due to, 91–92
 - drainage of, mediastinal mass discovery during, 257–258
 - in liver biopsy, 712–714
- Periodontitis, management of, before cardiac surgery, 587–591
- Peripheral vascular surgery, saphenous vein graft for, vasoactive agent effects on, 344–350
- Phenylephrine
 - cardiac output measurement effects of, 636–643
 - for saphenous vein graft blood flow, 344–350
- Pheochromocytoma, undiagnosed
 - cardiac hypokinesia and multiple organ failure due to, 863–866
 - hypertensive crisis due to, 858–862
- Plasma, solvent/detergent treated, in liver transplantation, fibrinolysis due to, 410–413
- Platelet(s)
 - activation of, in transfusions, with cell saver vs. systemic blood, 519–523
 - inhibition of, reversal of, for emergency carotid endarterectomy, 851–854
- Platelet inhibitors, in history of heparin-induced thrombocytopenia, 120–121
- Pleural cavity, pseudomyxoma peritonei extension to, chemotherapy for, 265–268
- Pleural effusion, drainage of, reexpansion pulmonary edema after, 887–891
- Pneumectomy
 - APACHE III evaluation after, 832–837
 - complications of, 743–751
 - double-lumen tube placement in, tracheal laceration in, 718–719
 - with horseshoe lung, 250–252
 - single-lung ventilation for, 883–886
- Pneumopericardium, in pediatric patients, 758–760
- Pneumothorax, reexpansion pulmonary edema after, 887–891
- Poly 2-methoxyethylacrylate-coated oxygenator, for cardiopulmonary bypass, hemostatic effects of, 28–34
- Polymer-coated oxygenator, for cardiopulmonary bypass, hemostatic effects of, 28–34
- Polyneuropathy, in Charcot-Marie-Tooth disease, CABG surgery in, 567–569
- Postoperative pain
 - high thoracic epidural anesthesia for, 810–815
 - interscalene brachial plexus block for, 554–557
 - intrapleural analgesia for, in chronic obstructive pulmonary disease, 816–819
 - magnesium with morphine for, 827–831
 - parasternal intercostal block for, 547–553
- Powelink device, for aneurysm repair, 730, 732
- Preconditioning, for cardiac surgery, isoflurane for, 683–689
- Preoperative evaluation, 323–324
- Propofol
 - for cardiac surgery
 - CABG, 810–815
 - in extended cardioplegia and reperfusion, 678–682
 - inflammatory mediators and, 512–518
 - mitral valve, 672–677
 - vs. sevoflurane, 664–671
 - for mediastinal lymph node fine needle aspiration, 894–895
 - for radial artery graft spasm prevention, 41–44
- Prosthetic valve, thrombus of, in idiopathic hypereosinophilic syndrome, 434–435
- Prothrombin fragments, in transfusions, with cell saver vs. systemic blood, 519–523
- Pseudoaneurysms
 - aortic arch, in cocaine abuse, 462–464
 - after central venous access, endovascular covered stent for, 99–102
- Pseudomyxoma peritonei extension, chemotherapy for, 265–268
- Pulmonary artery catheter
 - in cardiac output measurement, 3–7
 - evidence against using, 144–146
 - evidence for using, 147–149
- Pulmonary edema, reexpansion, 887–891
- Pulmonary hypertension
 - in cardiac surgery, on obese persons, 137–140
 - detection of, before heart transplantation, 51–56
 - sildenafil for, 203–207
- Pulmonary vascular resistance, measurement of, before heart transplantation, 51–56
- Quality of life, after cardiac surgery, in off-pump to on-pump intraoperative conversion, 793–799
- Radial artery graft, for CABG, intravenous anesthetic effects on, 41–44
- Rectus abdominis sheath hematoma, after aortic valve replacement, 612–614
- Reexpansion pulmonary edema, 887–891
- Regional anesthesia, for carotid endarterectomy, vs. general anesthesia, 127–132
- Remifentanyl
 - in cardiac surgery, inflammatory mediators and, 512–518
 - for fast-track anesthesia, 35–40
- Remon Impression AAA Sac Pressure Transducer, 735–736
- Reperfusion
 - injury prevention in, verapamil for, 337–343
 - myocardial, thrombin inhibition during, 224–231
 - propofol effects on, 678–682

- Ring-Adair-Ewyn tube, for one-lung ventilation, in pneumonectomy, 883–886
- Risk analysis, in cardiac surgery, with off-pump to on-pump intraoperative conversion, 793–799
- Robotics, 592–596
- Rocuronium, hypertensive crisis due to, 858–862
- Ropivacaine, in postoperative parasternal intercostal block, 547–553
- Sac Pressure Transducer, 735–736
- Saline solution, for left ventricular dysfunction, 400–405
- Sano shunt, resistant hypoxemia in, 880–882
- Saphenous vein graft, for peripheral vascular surgery, vasoactive agent effects on, 344–350
- S-100 β protein, in transfusions, with cell saver vs. systemic blood, 519–523
- Sevoflurane, for cardiac surgery, 502–511, 664–671
- Shoulder pain, after thoracotomy, interscalene brachial plexus block for, 554–557
- Sildenafil, for pulmonary hypertension, 203–207
- Single-lung ventilation, for pneumonectomy, 883–886
- Sinus of Valsalva, aneurysm of, 615–617
- Sodium, abnormalities of, in cardiopulmonary bypass, 273–275
- Sonoclot Analyzer, for activated coagulation time, in cardiopulmonary bypass, 524–528
- Spasm, of radial artery graft, prevention of, 41–44
- Spectroscopy, near-infrared, for oxygenation monitoring, 535–539
- Specular reflection, in transesophageal echocardiography, 460–461
- Spinal anesthesia, combined with general anesthesia, for anorectoplasty, 873–875
- Spinal artery syndrome, after off-pump CABG surgery, 709–711
- Spinal cord ischemia, due to pericardial effusion, 91–92
- Splanchnic circulation, in cardiac output promotion, after cardiac surgery, 351–356
- Stab injury, mammary artery hematoma in, 307–309
- Stents
- aortic, for aneurysm repair, 730–742, 763–765
 - drug-eluting
 - antiplatelet agents with, 436–442
 - with epidural catheter, 701–703 - endovascular covered, for pseudoaneurysm, 99–102
 - for superior vena cava syndrome, 417–419
- Sternotomy, cardiac output measurement in, 636–643
- Stroke
- after cardiac surgery
 - magnetic resonance imaging in, 106–109
 - prevention of, evidence-based guidelines for, 529–534 - after carotid endarterectomy, 127–132, 540–546
- Stroke volume
- pericardial lidocaine effects on, 393–399
 - variation in, fluid responsiveness and, 650–654
- Subarachnoid drainage catheter, complications of, 763–765
- Subclavian artery, pseudoaneurysm of, after central venous access, endovascular covered stent for, 99–102
- Subdural hematoma, after spinal catheter drainage, 704–708
- Substernal epicardial echocardiography, approach to, 237–242
- Sufentanil
- for CABG, 810–815
 - for fast-track anesthesia, 35–40
- Superior vena cava, persistent left, 761–762
- Superior vena cava syndrome, after cardiac surgery, 417–419
- Suppression ratio, in deep hypothermic circulatory arrest with retrograde cerebral perfusion, 61–67
- Swan, Jeremy, Moffitt collaboration with, 285–287
- Tachypnea, in reexpansion pulmonary edema, 887
- Talent LPS device, for aneurysm repair, 730, 732
- Teeth, optimizing health of, before cardiac surgery, 587–591
- Telemedicine, 592–596
- Temperature, in cardiopulmonary bypass, inflammatory mediator release and, 191–196
- Tetralogy of Fallot
- with absent left pulmonary artery, undiagnosed airway abnormalities with, 876–879
 - surgery for, air-leak test in, 197–202
- Thermodilution technique
- for cardiac output
 - vs. arterial pressure-based method, 3–7, 629–631, 632–635, 636–643
 - pulmonary artery vs. transpulmonary aortic methods for, 659–663
 - for right ventricular ejection fraction, 644–649
 - for right ventricular volume, 644–649
- Thiopental, in radial artery graft spasm prevention, 41–44
- Thoracic epidural anesthesia, high, for CABG, 810–815
- Thoracotomy
- cardiac output measurement in, partial carbon dioxide rebreathing method for, 655–658
 - postoperative analgesia for, magnesium sulfate with morphine, 827–831
 - shoulder pain after, interscalene brachial plexus block for, 554–557
- Thrombin
- formation of, in cardiopulmonary bypass, polymer-coated oxygenator effects on, 28–34
 - inhibition of, during reperfusion, 224–231
 - in transfusions, with cell saver vs. systemic blood, 519–523
- Thrombocytopenia, heparin-induced
- acute, cardiac surgery with, 122–123
 - after cardiac surgery, 18–22
 - after heart transplantation, 269–272
 - history of, alternative anticoagulants for, 113–126
 - subacute, cardiac surgery with, 123
- Thromboelastography, platelet-mapping, for emergency carotid endarterectomy, 851–854
- Thromboglobulin, in transfusions, with cell saver vs. systemic blood, 519–523
- Thrombosis and thrombi
- in aortic arch repair, 897–902
 - coronary, with aprotinin, 558–560
 - internal jugular vein, 422–424
 - intracardiac, Factor VIII inhibitor bypass activity administration, thromboses and, 561–563
 - left ventricular, 305–306
 - prosthetic valve, in idiopathic hypereosinophilic syndrome, 434–435
 - right atrial, after off-pump CABG surgery, 564–566
- Thymoma, peripheral cardiopulmonary bypass for, 867–869
- Thyroid artery, inferior, injury of, in internal jugular vein catheterization, 422–424
- Ticlopidine, for percutaneous coronary interventions, 436–442
- Tirofiban-heparin, in history of heparin-induced thrombocytopenia, 121
- Total intravenous anesthesia, for CABG, 810–815
- Trachea
- bifurcation of, undiagnosed, tetralogy of Fallot with, 876–879
 - compression of, in aortic aneurysm, 88–90
 - laceration of, in double-lumen tube placement, 718–719
- Tracheostomy, retrospective review of, in intensive care unit, 76–80
- Transesophageal echocardiography
- in aortic atherosclerosis assessment, 805–809
 - in aortic dissection with aortic regurgitation, 85–87

- Transesophageal echocardiography (*Continued*)
- in aortic regurgitation, mitral valve prosthesis obscuring, 610–611
 - in aortic valve replacement, myocardial function after, 367–370
 - in carbon dioxide embolism, 715–717
 - in complex aortic pathology, 462–464
 - in elephant trunk procedure, 584–586
 - in inferior vena cava diameter measurement, 492–496
 - in intracardiac thrombosis, 561–563
 - in left ventricular outflow obstruction, during liver transplantation, 577–580
 - mediastinal mass discovery in, 257–258
 - in misplaced guidewire, 420–421
 - in mitral valve prolapse, 581–583
 - in persistent left superior vena cava, 761–762
 - probe from, aortic compression by, 724–726
 - in right ventricular leiomyosarcoma, 432–433
 - routine, in cardiac surgery, effects on surgical management, 800–804
 - in sinus of Valsalva aneurysm, 615–617
 - specular reflection in, 460–461
 - in transmitral flow velocity propagation, 483–485, 486–491
 - in valvular heart disease, 855–857
- Transfusions
- cell saver vs. systemic blood, inflammatory mediators and, 519–523
 - with contaminated red blood cells, cardiovascular collapse due to, 870–872
 - requirements for, in cardiac surgery, kallikrein inhibitor effects on, 23–27
 - universal leukocyte reduction in, 232–236
- Transmitral flow propagation velocity, for diastolic function measurement, 483–485, 486–491
- Transplantation
- heart
 - heparin-induced thrombocytopenia in, 269–272
 - propofol for, in extended cardioplegia and reperfusion, 678–682
 - pulmonary hypertension detection before, 51–56
 - transesophageal echocardiography during, 800–804
 - heart-liver, 253–256
 - liver
 - cardiovascular collapse in, contaminated red blood cell transfusion and, 870–872
 - extubation timing in, 752–757
 - fibrinolysis during, 410–413
 - left ventricular outflow tract obstruction during, 577–580
 - lung, for cystic fibrosis, cardiopulmonary bypass in, 208–211
- Transposition of great arteries, correction of
- air-leak test in, 197–202
 - perioperative interleukin levels in, 184–190
- Trauma, facial, repair of, subtotal airway obstruction in, 903–905
- Tricuspid valve, replacement of
- with carcinoid syndrome, 855–857
 - transesophageal echocardiography during, 800–804
- Tumor necrosis factor α , after cardiac surgery, dexamethasone effects on, 68–75
- Ultrasonography
- of carotid artery, during aortic dissection repair, 81–84
 - epiaortic, in aortic atherosclerosis assessment, 805–809
 - in mediastinal lymph node fine needle aspiration, 892–896
- Universal leukocyte reduction, in cardiac surgery, 232–236
- Valvular heart disease. *See also specific valves*
- in carcinoid syndrome, 855–857
 - surgery for, transesophageal echocardiography during, 800–804
- Vasorelaxants, for radial artery graft, 41–44
- Vena cava, inferior, diameter of, vs. central venous pressure, in cardiac surgery, 492–496
- Venovenous hemodialysis, after cardiac surgery, 379–383
- Venovenous hemofiltration, for anticoagulant reversal, after heart transplantation, 269–272
- Ventilation. *See Mechanical ventilation*
- Ventricle(s)
- left
 - dysfunction of, saline solution for, 400–405
 - outflow obstruction of, during liver transplantation, 577–580
 - thrombus in, 305–306
 - right
 - drainage obstruction from, after biventricular assist device implantation, 262–264
 - leiomyosarcoma of, 432–433
 - volume of, thermodilution measurement of, 644–649
- Ventricular fibrillation, after aortic valve replacement, amiodarone for, 414–416
- Ventricular septal defect, surgery for, air-leak test in, 197–202
- Verapamil
- overdose of, pathophysiology of, 406–409
 - for reperfusion injury prevention, 337–343
- Vigileo monitor, for cardiac output measurement, 629–631, 632–635, 636–643
- Visser, Cees A., memorial statement for, 917
- Warming phase, of cardiopulmonary bypass, carbon dioxide levels during, 57–60
- Zenith device, for aneurysm repair, 730, 732, 734