

SUBJECT INDEX

Age and outcome

- , Age is not a contraindication for repair of thoracic aortic aneurysm, Silvey, G., et al., O-43
- , Is heart surgery a possible option for octogenarians?, Corona, A., et al., P-27
- , A study of cardiac bypass surgery outcome related to patient's age, Massó, E., et al., P-26

Airway device, I-gel supraglottic airway for ventilation during on-pump coronary artery bypass grafting surgery, Samalavicius, R., et al., P-17

Airway pressure

- , Continuous positive airway pressure at 10 cm H₂O during cardiopulmonary bypass does not improve postoperative oxygenation in paediatric patients, Kim, J.-T., et al., P-37
- , Evaluating the “safe mean airways pressure” in patients with ARDS: an echocardiographic retrospective approach, Di Nardo, M., et al., P-38
- , Features of ventilator in surgical treatment of lung echinococcosis patients, Miyerbekov, Y., et al., P-35
- , The effects of pulmonary ventilation during cardio-pulmonary bypass on pulmonary function after open heart surgery, Gorgu, A., et al., O-39

Ambulatory ECG, Impact of the observance of the ACC/AHA guidelines for interpretability of ambulatory electrocardiography on the association of silent ischaemia with troponin release, Lurati Buse, G., et al., O-66

Anaemia, The influence of profound haemodilution on indocyanine green plasma disappearance rate and hepatic function after normothermic cardiopulmonary bypass, von Heymann, C., et al., O-36

Anaerobic metabolism, Anaerobic metabolism during thoracic anaesthesia for hyperthermic intrathoracic chemotherapy, Scolletta, S., et al., O-50

Anaesthesia techniques, Comparison of the accuracy of central venous catheter placements between the landmark-guided and electrocardiogram-guided techniques, Lee, J.-H., et al., P-04

Anaesthetic management

- , Anaesthetic management of endovascular repair of type B thoracic aortic dissection, Matute-Jimenez, P., et al., P-06
- , Perioperative management for complex interventions in patients with renal tumours and cavo-atrial thrombosis, Ghenu, O., et al., P-51

Anaesthetics, The effects of propofol and sevoflurane on modulation of inflammation and oxidative stress in the kidney following aortic cross clamping, Rodriguez-Lopez, J.M., et al., O-48

Analgesia, Does pregabalin improve the analgesia in cardiac surgery? A preliminary study, de Cosmo, D., et al., P-40

Angina pectoris

- , Perioperative predictors of postoperative pain at coronary artery bypass grafting, Siciliano, T., et al., O-57
- , Treatment of chronic refractory angina pectoris, Dobias, M., et al., P-46
- , Spinal cord stimulator implantation for treatment of angina pectoris gave dysaesthetic pain and burning skin sensations: case reports, Andersen, C., et al., P-59

Anti-arrhythmic, Sodium and calcium channel blockade by esmolol, Fallouh, H.B., et al., P-07

Aortic aneurysm

- , Intraoperative EEG and prognosis in aortic reconstruction surgery when using selective cerebral perfusion, Demir, A., et al., P-22
- , Age is not a contraindication for repair of thoracic aortic aneurysm, Silvey, G., et al., O-43

Aortic dissection

- , High preoperative central venous pressure is a risk factor for mortality and adverse neurological outcome after repair of acute type A aortic dissection, Basciani, R.M., et al.
- , Anaesthetic management of endovascular repair of type B thoracic aortic dissection, Matute-Jimenez, P., et al., P-06

Aortic stenosis

- , Effects of sildenafil on left ventricular hypertrophy in a porcine model, Dahl Christensen, S., et al., P-48
- , NT-proBNP as a marker of pulmonary hypertension in cardiac surgical patients with aortic stenosis, Kozlov, I., et al., P-58

Aortic valve replacement

- , Aortic valve replacement with a mini-extracorporeal circulation circuit, Flo, A., et al., O-35
- , Preoperative statin treatment is not associated with reduced postoperative stratified mortality in patients undergoing aortic valve replacement, Reynaud, S., et al., P-33

Aprotinin, Aprotinin is superior to tranexamic acid in paediatric cardiac surgery, Breuer, T., et al., O-34

ARDS, Evaluating the “safe mean airways pressure” in patients with ARDS: an echocardiographic retrospective approach, Di Nardo, M., et al., P-38

Arterial hypertension, The effect of hypertension on neurocognitive function tests in patients who undergo CABG operations done with TIVA, Muslu, S., et al., O-60

Autologous blood transfusion, Assessment of the haematological and haemodynamic influence of colloid and crystalloid solutions in acute normovolaemic haemodilution in coronary artery bypass grafting surgery, Aghadavoudi, O., et al., O-04

Autonomic nervous system, Low arterial pressure after low dose propofol administration: haemodynamic and autonomic nervous system activity, Akselrod, A., et al., P-54

Beta-blocker, Sodium and calcium channel blockade by esmolol, Fallouh, H.B., et al., P-07

Bispectral index

- , Esmolol attenuates BIS response to intubation during fentanyl-midazolam anaesthesia for cardiac surgery, Yazicioglu, H., et al., O-29
- , Central nervous system dysfunction after cardiac surgery: role of bispectral index (BIS), cerebral oxygen saturation and aortic atheromas, Fita, G., et al., P-20
- , Do Bispectral Index values from two sensors placed on each side of the forehead always provide the same value? Case report, Choi, I.C., et al., P-01

Blood cardioplegia, The cardioprotective effects of N-acetylcysteine as an additive to blood cardioplegia during coronary artery bypass grafting, Aghadavoudi, O., et al., O-71

Brain damage, Narcotrend® monitoring is highly specific for detection of brain damage in patients undergoing hypothermic circulatory arrest, Groesdonk, H.V., et al., P-21

Brain natriuretic peptide, Diagnostic value of brain natriuretic peptide in healthy young subjects, Bolliger, D., et al., O-24

Cardiac anaesthesia

- , CHANGE STUDY (Changing intraoperative Anaesthesia strategy based on cerebral oximetry data: effect on incidence of neurological dysfunction in adult cardiac surgery patients), Kobzova, E., et al., P-18
- , Perioperative predictors of postoperative pain at coronary artery bypass grafting, Siciliano, T., et al., O-57
- , Esmolol attenuates BIS response to intubation during fentanyl-midazolam anaesthesia for cardiac surgery, Yazicioglu, H., et al., O-29
- , An investigation of oxidative stress and DNA damage in patients exposed to cardiopulmonary bypass surgery, Mordeniz, C., et al., O-40

- , Comparison of the effects of different fentanyl protocols on postoperative pain control after coronary artery surgery, Tur, H., et al., P-42
- Cardiac output**
- , Comparison of pressure waveform analysis with thermodilution via pulmonary artery catheter (PAC) in valve replacement surgery, Kazanci, D., et al., P-34
- , Haemodynamic monitoring during off-pump CABG: validation of the FloTrac TM-VigileoTM System, Turani, F., et al., O-21
- , Continuous arterial pressure waveform analysis accurately detects cardiac output in cardiac surgery: a prospective comparison with thermodilution, echocardiography and magnetic resonance techniques, Senay, S., et al., O-13
- Cardiac remodelling**, Effects of sildenafil on left ventricular hypertrophy in a porcine model, Dahl Christensen, S., et al., P-48
- Cardioprotection**, Cardioprotective effect of volatile anaesthetics on mitochondrial respiratory function, Sirvinskas, E., et al., O-67
- Cardiopulmonary bypass**
- , Gastrointestinal complications in patients undergoing open heart surgery, Donmez, A., et al., P-43
- , Low haematocrit during cardiopulmonary bypass is not associated with adverse outcome after coronary bypass surgery, Senay, S., et al., O-62
- , Oxygen saturation variability in the frontal lobe cortex during cardiopulmonary bypass, Wolski, P., et al., O-08
- , Cardioprotective effect of volatile anaesthetics on mitochondrial respiratory function, Sirvinskas, E., et al., O-67
- Carotid stenosis**, Management of patients with asymptomatic carotid stenosis undergoing CABG, Dobias, M., et al., O-53
- Case report**
- , The use of real-time TOE in the intra-operative management of renal cell carcinoma, inferior vena cava thrombus and pulmonary metastasis: case report, Mahdy, S., et al., P-52
- , Do Bispectral Index values from two sensors placed on each side of the forehead always provide the same value? Case report, Choi, I.C., et al., P-01
- , Spinal cord stimulator implantation for treatment of angina pectoris gave dysaesthetic pain and burning skin sensations: case reports, Andersen, C., et al., P-59
- , Perioperative management for complex interventions in patients with renal tumours and cavo-atrial thrombosis, Ghenu, O., et al., P-51
- Cavo-atrial thrombosis**, Perioperative management for complex interventions in patients with renal tumours and cavo-atrial thrombosis, Ghenu, O., et al., P-51
- Central venous catheterization**
- , Comparison of the accuracy of central venous catheter placements between the landmark-guided and electrocardiogram-guided techniques, Lee, J.-H., et al., P-04
- , External jugular vein catheterization using 'intra-atrial ECG', Karaaslan, D., et al., P-03
- Central venous pressure**, High preoperative central venous pressure is a risk factor for mortality and adverse neurological outcome after repair of acute type A aortic dissection, Basciani, R.M., et al., O-44
- Cerebral monitoring**, Do Bispectral Index values from two sensors placed on each side of the forehead always provide the same value? Case report, Choi, I.C., et al., P-01
- Cerebral oxygen saturation**
- , CHANGE STUDY (Changing intraoperative Anaesthesia strategy based on cerebral oximetry data: effect on incidence of neurological dysfunction in adult cardiac surgery patients), Kobzova, E., et al., P-18
- , Intraoperative regional cerebral oxygen saturation monitoring in risk patients undergoing cardiac surgery: predictors of desaturation, Lourenço, A.P., et al., O-06
- , Oxygen saturation variability in the frontal lobe cortex during cardiopulmonary bypass, Wolski, P., et al., O-08
- , Central nervous system dysfunction after cardiac surgery: role of bispectral index (BIS), cerebral oxygen saturation and aortic atheromas, Fita, G., et al., P-20
- Cerebral oxymetry**
- , Intraoperative regional cerebral oxygen saturation monitoring in risk patients undergoing cardiac surgery: predictors of desaturation, Lourenço, A.P., et al., O-06
- , Pre-induction value of cerebral oximetry as a predictor of neurological outcome from coronary artery bypass surgery: a pilot study, Argiriadou, H., et al., P-19
- , Cerebral oxygenation monitoring in patients undergoing deep hypothermic circulatory arrest for cardiothoracic surgery, Heringlake, M., et al., O-07
- Cerebral perfusion pressure**, The effect of hypertension on neurocognitive function tests in patients who undergo CABG operations done with TIVA, Muslu, S., et al., O-60
- Cerebral perfusion**, Intraoperative EEG and prognosis in aortic reconstruction surgery when using selective cerebral perfusion, Demir, A., et al., P-22
- Coagulation**
- , Comparison of clinical response to platelet transfusion and bedside impedance aggregometry, Jambor, C., et al., O-03
- , In vitro reversal of warfarin effects with prothrombin complex concentrate vs. fresh frozen plasma, Szlam, F., et al., P-10
- , Anaesthesia and coagulation status in cardiac surgery patients, Levit, A., et al., P-11
- , Kaolin activated and heparinase modified TEG predict post-operative bleeding in neonatal cardiac surgery, Del Sarto, P., et al., P-09
- Colloids**
- , Crystalloid (Ringer's solution) versus colloid 6% HES (130/04) for pump priming in CABG surgery, Sogut, F., et al., O-38
- , Assessment of the haematological and haemodynamic influence of colloid and crystalloid solutions in acute normovolaemic haemodilution in coronary artery bypass grafting surgery, Aghadavoudi, O., et al., O-04
- Combined anaesthesia**, Combined general and regional anaesthesia in pulmonary endarterectomy, Rubes, D., et al., O-45
- Complications**
- , Features of ventilator in surgical treatment of lung echinococcosis patients, Miyerbekov, Y., et al., P-35
- , Prevalence of early postoperative complications and mortality in patients undergoing cardiac surgery, Musiał-Swiętlikiewicz, V., et al., P-25
- , Risk factors for selected early postoperative complications after cardiac surgery, Musiał-Swiętlikiewicz, V., et al., O-65
- , Dexmedetomidine in cardiac surgery patients who failed extubation and in delirium statement, Yapici, N., et al., O-31
- Congenital cardiac surgery**, Effects of low body weight on congenital heart surgery and risk classifications, Akpek, E.A., et al., P-32
- Contrast media**, Angiography timing and contrast media amount: new risk factors for renal dysfunction after cardiac operations, Ranucci, M., et al., O-09
- Coronary artery bypass surgery**
- , Possible technique of myocardial preconditioning by sevoflurane in patients undergoing coronary surgery, Zadorozh, M., et al., O-68
- , A study of cardiac bypass surgery outcome related to patient's age, Massó, E., et al., P-26
- , Perioperative allogeneic blood transfusion is associated with increased thirty day mortality after isolated coronary bypass surgery, Senay, S., et al., O-61
- Coronary artery disease**, Quality of life in coronary artery bypass graft candidates: the SF-36 compared to the WHOQOL-BREF, Najafi, M., et al., O-26

- C-reactive protein**, Preoperative high sensitivity C-reactive protein predicts cardiovascular morbidity after coronary artery bypass grafting, Balciunas, M., et al., P-30
- Cristalloids**, Assessment of the haematological and haemodynamic influence of colloid and crystalloid solutions in acute normovolaemic haemodilution in coronary artery bypass grafting surgery, Aghadavoudi, O., et al., O-04
- Cytokines**
- , Improved analgesic effect in peridural thoracic versus total intravenous anaesthesia during cardiac surgery: role of epicardial fat-secreted cytokines, Cappelletti, M., et al., P-41
 - , Continuous renal replacement therapy decreases pro BNP and IL 6 in cardiac surgery patients with postoperative sepsis C 102 The effects of sevoflurane and desflurane on serum S100 β and NSE levels in cyanotic children undergoing open heart surgery, Turani, F., et al., P-45
- Deep hypothermic cardiac arrest**
- , Combined general and regional anaesthesia in pulmonary endarterectomy, Rubes, D., et al., O-45
 - , The effects of deep hypothermic circulatory arrest on tracheal tube cuff pressure dynamics, Rubes, D., et al., P-08
 - , Cerebral oxygenation monitoring in patients undergoing deep hypothermic circulatory arrest for cardiothoracic surgery, Heringlake, M., et al., O-07
- Depth of anaesthesia**, Assessment of the index of consciousness during cardiac anaesthesia, Galán, J., et al., P-02
- Desflurane**
- , Neutrophil activation during off-pump coronary artery bypass grafting: a comparison between sevoflurane, desflurane and total intravenous anaesthesia, Durak, P., et al., O-41
 - , Comparison of right ventricle haemodynamic parameters between patients anaesthetized with propofol or desflurane for off-pump CABG, Mroziński, P., et al., P-55
- Dexmedetomidine**
- , Dexmedetomidine in cardiac surgery patients who failed extubation and in delirium statement, Yapici, N., et al., O-31
 - , Propofol-dexmedetomidine and propofol-ketamine combinations for anaesthesia in paediatric patients undergoing transcatheter atrial septal closure, Koruk, S., et al., O-30
- Diabetes mellitus**, Predictors of postoperative length of stay of CABG patients: comparison of diabetics and non-diabetics, Najafi, M., et al., O-59
- Diagnostic value**
- , Impact of the observance of the ACC/AHA guidelines for interpretability of ambulatory electrocardiography on the association of silent ischaemia with troponin release, Lurati Buse, G., et al., O-66
 - , Diagnostic value of brain natriuretic peptide in healthy young subjects, Bolliger, D., et al., O-24
- DNA damage**, An investigation of oxidative stress and DNA damage in patients exposed to cardiopulmonary bypass surgery, Mordeniz, C., et al., O-40
- Dopamine**, Effects of low dose dopamine on renal oxygen consumption, blood flow and glomerular filtration rate after cardiac surgery, Redfors, B., et al., O-51
- Doppler ultrasound**, Use of audio-Doppler ultrasonography guidance for paediatric internal jugular vein catheterization: effect on experience of the operator, Akpek, E.A., et al., O-18
- ECG**
- , Comparison of the accuracy of central venous catheter placements between the landmark-guided and electrocardiogram-guided techniques, Lee, J.-H., et al., P-04
 - , External jugular vein catheterization using 'intra-atrial ECG', Karaaslan, D., et al., P-03
- Echinococcosis**, Features of ventilatin in surgical treatment of lung echinococcosis patients, Miyerbekov, Y., et al., P-35
- Echocardiography**
- , The use of real-time TOE in the intra-operative management of renal cell carcinoma, inferior vena cava thrombus and pulmonary metastasis: case report, Mahdy, S., et al., P-52
 - , Evaluating the "safe mean airways pressure" in patients with ARDS: an echocardiographic retrospective approach, Di Nardo, M., et al., P-38
 - , Feasibility of measuring myocardial performance index of the right ventricle in anaesthetized patients, Michaux, I., et al., O-11
 - , Assessment of right ventricular functions and pulmonary haemodynamics during lung resection guided by TOE, Galal, A., et al., O-12
 - , Comparison of dP/dt estimates measured by PiCCO® and TOE during cardiac surgery, Velghe, D., et al., P-14
 - , Anaesthetic management of endovascular repair of type B thoracic aortic dissection, Matute-Jimenez, P., et al., P-06
 - , Transoesophageal echocardiography and tricuspid annular dilation: a prospective comparison with surgical measurement. a preliminary study, Parra, V., et al., P-13
 - , Placing a saline bag during cardiac positioning and stabilization aids transoesophageal echocardiography in off-pump coronary artery bypass surgery, Kim, T.-Y., et al., P-15
 - , Assessment of mitral valve pathologies with intraoperative transoesophageal echocardiography: comparison of the 2D mode with the realtime 3D technique, Koncar-Zeh, J., et al., O-16
 - , Visibility of cardiac valves using real-time 3D-transoesophageal echocardiography (RT3D-TOE) in the OR: a feasibility study, Tschernich, H., et al., O-17
- Echocardiography, speckle tracking**, Myocardial deformation assessed by speckle tracking echocardiography: in vitro investigation of potential limitations and pitfalls, Sivesgaard, K., et al., O-15
- Echocardiography, strain**, Two dimensional strain echocardiography: a feasibility and repeatability study, Krøijer, R., et al., O-14
- EEG**
- , Intraoperative EEG and prognosis in aortic reconstruction surgery when using selective cerebral perfusion, Demir, A., et al., P-22
 - , Neurological monitoring during cardiac surgery: is it sufficient?, Boran, E., et al., O-10
 - , Narcotrend® monitoring is highly specific for detection of brain damage in patients undergoing hypothermic circulatory arrest, Groesdonk, H.V., et al., P-21
- Ejection fraction**, Effects of levosimendan in cardiac surgery, Basaran, B., et al., P-47
- Epidural anaesthesia**
- , Improved analgesic effect in peridural thoracic versus total intravenous anaesthesia during cardiac surgery: role of epicardial fat-secreted cytokines, Cappelletti, M., et al., P-41
 - , Can skin resistance be used to evaluate the sympathetic block of a thoracic epidural blockade?, Andersen, C., et al., P-05
 - , Thoracic epidural anaesthesia impairs the haemodynamic response to acute pulmonary hypertension by depressing right ventricular contractility in pigs, Rex, S., et al., O-47
- Esmolol**
- , Esmolol attenuates BIS response to intubation during fentanyl-midazolam anaesthesia for cardiac surgery, Yazicioglu, H., et al., O-29
 - , Sodium and calcium channel blockade by esmolol, Fallouh, H.B., et al., P-07
- Euroscore**, Health related quality of life following cardiac surgery—correlation with EuroSCORE, Colak, Z., et al., O-25
- Extubation**, Very early extubation in children after cardiac surgery, Karahan, N., et al., P-36
- Factor VIIa**, Thrombelastographic analysis of recombinant factor VIIa in cardiac surgery, Prati, P., et al., O-33

Factor XIII, Predictors of clot firmness after high risk cardiac surgery, Jambor, C., et al., O-01

Feasibility

- , Placing a saline bag during cardiac positioning and stabilization aids transoesophageal echocardiography in off-pump coronary artery bypass surgery, Kim, T.-Y., et al., P-15
- , I-gel supraglottic airway for ventilation during on-pump coronary artery bypass grafting surgery, Samalavicius, R., et al., P-17
- , Two dimensional strain echocardiography: a feasibility and repeatability study, Krøijer, R., et al., O-14

Fentanyl

- , Comparison of the effects of different fentanyl protocols on postoperative pain control after coronary artery surgery, Tur, H., et al., P-42
- , Fentanyl causes vasodilation via K⁺ channels, Na⁺-K⁺-ATPase and nitric oxide: an in vitro study, Sasmaz, S., et al., O-27

Fibrinogen, Predictors of clot firmness after high risk cardiac surgery, Jambor, C., et al., O-01

Fresh frozen plasma, In vitro reversal of warfarin effects with prothrombin complex concentrate vs. fresh frozen plasma, Szlam, F., et al., P-10

Gastrointestinal complications, Gastrointestinal complications in patients undergoing open heart surgery, Donmez, A., et al., P-43

Geriatric surgery

- , Age is not a contraindication for repair of thoracic aortic aneurysm, Silvaj, G., et al., O-43
- , Is heart surgery a possible option for octogenarians?, Corona, A., et al., P-27
- , A study of cardiac bypass surgery outcome related to patient's age, Massó, E., et al., P-26

Haematocrit, Low haematocrit during cardiopulmonary bypass is not associated with adverse outcome after coronary bypass surgery, Senay, S., et al., O-62

Haemodilution, The influence of profound haemodilution on indocyanine green plasma disappearance rate and hepatic function after normothermic cardiopulmonary bypass, von Heymann, C., et al., O-36

Haemodynamics, Low arterial pressure after low dose propofol administration: haemodynamic and autonomic nervous system activity, Akselrod, A., et al., P-54

Haemofiltration, Factors determining long term survival in cardiac surgery patients with acute renal failure treated with haemofiltration, Klajbor, K., et al., O-52

Haemostasis

- , Thrombelastographic analysis of recombinant factor VIIa in cardiac surgery, Prati, P., et al., O-33
- , Determinants of postoperative bleeding after coronary bypass surgery, Senay, S., et al., O-05

Haemotransfusion, Impact of autologous shed mediastinal blood reinfusion on clinical outcome in the early postoperative period after cardiac surgery, Andrejaitiene, J., et al., P-39

Heart failure, Perioperative inflammatory time-course in patients receiving left ventricular assist for end-stage heart failure, Mondino, M., et al., O-58

Heparin, Minimum discard volume to obtain reliable activated clotting time from the heparinized arterial line, Erden, I.A., et al., P-12

Hepatic function, The influence of profound haemodilution on indocyanine green plasma disappearance rate and hepatic function after normothermic cardiopulmonary bypass, von Heymann, C., et al., O-36

HES 130/0.4, Priming of the cardiopulmonary bypass with HES 130/0.4 compared to lactated Ringer's solution reduces volume load but does not change subcutaneous tissue, Schiferer, A., et al., O-37

Holter ECG, Impact of the observance of the ACC/AHA guidelines for interpretability of ambulatory electrocardiography on the association of silent ischaemia with troponin release, Lurati Buse, G., et al., O-66

Hyperfibrinolysis, Anaesthesia and coagulation status in cardiac surgery patients, Levit, A., et al., P-11

Hyperglycaemia

- , Effectiveness of predictive algorithm eMPC2 with variable sampling interval in perioperative glycaemia control, Kopecky, P., et al., O-55
- , Blood glucose control in cardiac surgery patients: a comparative study of different insulin protocols, Blaha, J., et al., O-54

Iloprost, Effects of inhaled iloprost on right ventricular contractility and ventricular interdependence in acute pulmonary hypertension in pigs, Rex, S., et al., O-28

Index of consciousness, Assessment of the index of consciousness during cardiac anaesthesia, Galán, J., et al., P-02

Inflammation

- , Preoperative high sensitivity C-reactive protein predicts cardiovascular morbidity after coronary artery bypass grafting, Balciunas, M., et al., P-30
- , The effects of propofol and sevoflurane on modulation of inflammation and oxidative stress in the kidney following aortic cross clamping, Rodriguez-Lopez, J.M., et al., O-48

Inotropic agents, Effects of levosimendan in cardiac surgery, Basaran, B., et al., P-47

Insulin

- , Effectiveness of predictive algorithm eMPC2 with variable sampling interval in perioperative glycaemia control, Kopecky, P., et al., O-55
- , Blood glucose control in cardiac surgery patients: a comparative study of different insulin protocols, Blaha, J., et al., O-54

Intensive care medicine

- , Effects of low dose dopamine on renal oxygen consumption, blood flow and glomerular filtration rate after cardiac surgery, Redfors, B., et al., O-51
- , Continuous monitoring of ScvO₂ in paediatric patients undergoing cardiac surgery –, the pediasat technology, Ranucci, M., et al., O-19
- , Blood glucose control in cardiac surgery patients: a comparative study of different insulin protocols, Blaha, J., et al., O-54

Intrapleural analgesia, Effects of bilateral intrapleural block as pre-emptive analgesia on haemodynamic changes in coronary artery graft surgery, Mansouri, M., et al., P-57

Intrathoracic chemotherapy, Anaerobic metabolism during thoracic anaesthesia for hyperthermic intrathoracic chemotherapy, Scolletta, S., et al., O-50

Ischaemia-reperfusion injury, Protective effects of N-acetylcysteine infusion on ischaemia– reperfusion injury during CABG surgery, Cinar, D., et al., O-72

Isoflurane

- , Pharmacological preconditioning effect of isoflurane in the model of cardiac ischaemia-reperfusion in guinea pigs, Sahin, N., et al., O-74
- , A randomized comparison of isoflurane versus propofol TIVA in coronary artery bypass surgery: effects on postoperative cardiac troponin I release and clinical outcome, Flier, S., et al., O-69

Ketamine, Propofol-dexmedetomidine and propofol-ketamine combinations for anaesthesia in paediatric patients undergoing transcatheter atrial septal closure, Koruk, S., et al., O-30

Lactate, Tissue perfusion during minimized extracorporeal circulation (MECC), Sercia, G., et al., P-53

Left ventricular assist device, Perioperative inflammatory time-course in patients receiving left ventricular assist for end-stage heart failure, Mondino, M., et al., O-58

Length of stay, Predictors of postoperative length of stay of CABG patients: comparison of diabetics and non-diabetics, Najafi, M., et al., O-59

Levosimendan

- , Effects of levosimendan in cardiac surgery, Basaran, B., et al., P-47
- , The use of levosimendan associated with norepinephrine in a cardiac postoperative care unit, Coruh, T., et al., O-56
- , Levosimendan attenuates ischaemia-reperfusion injury of the heart better than ischaemic preconditioning during moderate hypothermia, Yapici, D., et al., O-70

Lobectomy, Assessment of right ventricular functions and pulmonary haemodynamics during lung resection guided by TOE, Galal, A., et al., O-12

Low body weight, Effects of low body weight on congenital heart surgery and risk classifications, Akpek, E.A., et al., P-32

Low cardiac output, The use of levosimendan associated with norepinephrine in a cardiac postoperative care unit, Coruh, T., et al., O-56

Lung protection, The effects of pulmonary ventilation during cardio-pulmonary bypass on pulmonary function after open heart surgery, Gorgu, A., et al., O-39

Minimized extracorporeal circulation

- , Aortic valve replacement with a mini-extracorporeal circulation circuit, Flo, A., et al., O-35
- , Coronary artery surgery with mini-CPB system vs. off-pump technique: comparison in terms of haemodilution and postoperative outcome. A prospective randomized study, Maestre Hittinger, M.L., et al., P-16
- , Tissue perfusion during minimized extracorporeal circulation (MECC), Sercia, G., et al., P-53

Mitral valve pathology, Assessment of mitral valve pathologies with intraoperative transoesophageal echocardiography: comparison of the 2D mode with the realtime 3D technique, Koncar-Zeh, J., et al., O-16

Monitoring

- , Oxygen saturation variability in the frontal lobe cortex during cardiopulmonary bypass, Wolski, P., et al., O-08
- , Feasibility of measuring myocardial performance index of the right ventricle in anaesthetized patients, Michaux, I., et al., O-11
- , Minimum discard volume to obtain reliable activated clotting time from the heparinized arterial line, Erden, I.A., et al., P-12
- , Myocardial deformation assessed by speckle tracking echocardiography: in vitro investigation of potential limitations and pitfalls, Sivesgaard, K., et al., O-15

Mucociliary transport velocity, Effect of smoking on bronchial mucus transport velocity assessed by the bronchoscope under total intravenous anaesthesia, Kim, Y.H., et al., P-29

Multiorgan Failure Syndrome, Perioperative inflammatory time-course in patients receiving left ventricular assist for end-stage heart failure, Mondino, M., et al., O-58

Myeloperoxidase

- , Neutrophil activation during off-pump coronary artery bypass grafting: a comparison between sevoflurane, desflurane and total intravenous anaesthesia, Durak, P., et al., O-41
- , Neutrophil activation and oxidative stress: a comparative study between off-pump and on-pump coronary artery bypass grafting, Durak, P., et al., O-42

Myocardial contractility, Comparison of dP/dt estimates measured by PiCCO® and TOE during cardiac surgery, Velghe, D., et al., P-14

Myocardial infarction, Postconditioning reduced infarct size in an open-chest porcine acute ischaemia-reperfusion model, Haarup Lie, R., et al., O-73

Myocardial ischaemia-reperfusion

- , Pharmacological preconditioning effect of isoflurane in the model of cardiac ischaemia-reperfusion in guinea pigs, Sahin, N., et al., O-74
- , Postconditioning reduced infarct size in an open-chest porcine acute ischaemia-reperfusion model, Haarup Lie, R., et al., O-73
- , Levosimendan attenuates ischaemia-reperfusion injury of the heart better than ischaemic preconditioning during moderate hypothermia, Yapici, D., et al., O-70

Myocardial protection

- , Possible technique of myocardial preconditioning by sevoflurane in patients undergoing coronary surgery, Zadorozh, M., et al., O-68
- , Protective effects of N-acetylcysteine infusion on ischaemia-reperfusion injury during CABG surgery, Cinar, D., et al., O-72
- , The cardioprotective effects of N-acetylcysteine as an additive to blood cardioplegia during coronary artery bypass grafting, Aghadavoudi, O., et al., O-71
- , Levosimendan attenuates ischaemia-reperfusion injury of the heart better than ischaemic preconditioning during moderate hypothermia, Yapici, D., et al., O-70
- , Effects of sevoflurane and propofol on myocardial recovery in coronary surgery, Sungur Ulke, Z., et al., P-24

N-acetylcysteine

- , Protective effects of N-acetylcysteine infusion on ischaemia-reperfusion injury during CABG surgery, Cinar, D., et al., O-72
- , The cardioprotective effects of N-acetylcysteine as an additive to blood cardioplegia during coronary artery bypass grafting, Aghadavoudi, O., et al., O-71

Natriuretic peptides

- , NT-proBNP as a marker of pulmonary hypertension in cardiac surgical patients with aortic stenosis, Kozlov, I., et al., P-58
- , Relationship between left ventricular wall stress, estimated by a pulse contour system, and pro-B-type natriuretic peptide in cardiac surgery, Scolletta, S., et al., O-22
- , The effects of physical exercise on plasma levels of relaxin, NTproANP, and NTproBNP in patients with ischaemic heart disease, Heringlake, M., et al., O-23

Near-infrared spectroscopy, Intraoperative regional cerebral oxygen saturation monitoring in risk patients undergoing cardiac surgery: predictors of desaturation, Lourenço, A.P., et al., O-06

Neonatal, Kaolin activated and heparinise modified TEGÒ predict post-operative bleeding in neonatal cardiac surgery, Del Sarto, P., et al., P-09

Neurocognitive function, The effect of hypertension on neurocognitive function tests in patients who undergo CABG operations done with TIVA, Muslu, S., et al., O-60

Neurologic complications, Dexmedetomidine in cardiac surgery patients who failed extubation and in delirium statement, Yapici, N., et al., O-31

Neurologic dysfunction

- , CHANGE STUDY (Changing intraoperative Anaesthesia strategy based on cerebral oximetry data: effect on incidence of neurologic dysfunction in adult cardiac surgery patients), Kobzova, E., et al., P-18
- , Central nervous system dysfunction after cardiac surgery: role of bispectral index (BIS), cerebral oxygen saturation and aortic atheromas, Fita, G., et al., P-20
- , High preoperative central venous pressure is a risk factor for mortality and adverse neurological outcome after repair of acute type A aortic dissection, Basciani, R.M., et al., O-44

Neurologic monitoring

- , Neurological monitoring during cardiac surgery: is it sufficient?, Boran, E., et al., O-10

- , Narcotrend® monitoring is highly specific for detection of brain damage in patients undergoing hypothermic circulatory arrest, Groesdonk, H.V., et al., P-21

Neurologic outcome

- , Cerebral oxygenation monitoring in patients undergoing deep hypothermic circulatory arrest for cardiothoracic surgery, Heringlake, M., et al., O-07
- , Pre-induction value of cerebral oximetry as a predictor of neurological outcome from coronary artery bypass surgery: a pilot study, Argiriadou, H., et al., P-19

Norepinephrine, The use of levosimendan associated with norepinephrine in a cardiac postoperative care unit, Coruh, T., et al., O-56

Off-pump coronary bypass surgery

- , Neutrophil activation and oxidative stress: a comparative study between off-pump and on-pump coronary artery bypass grafting, Durak, P., et al., O-42
- , I-gel supraglottic airway for ventilation during on-pump coronary artery bypass grafting surgery, Samalavicius, R., et al., P-17
- , Does off-pump coronary artery bypass have reno-protective effect in patients with abnormal preoperative renal function?, Papagiannopoulou P., et al., P-44
- , Coronary artery surgery with mini-CPB system vs. off-pump technique: comparison in terms of haemodilution and postoperative outcome. A prospective randomized study, Maestre Hittinger, M.L., et al., P-16
- , Comparison of two different total intravenous techniques for off-pump coronary artery bypass graft surgery: a randomized double-blind study, Askin, T., et al., P-56
- , Troponin I is an independent predictor of long term outcome after off-pump coronary surgery, Timmermans, A., et al., O-64
- , Haemodynamic monitoring during off-pump CABG: validation of the FloTrac TM-VigileoTM System, Turani, F., et al., O-21
- , Placing a saline bag during cardiac positioning and stabilization aids transoesophageal echocardiography in off-pump coronary artery bypass surgery, Kim, T.-Y., et al., P-15

One-lung ventilation, Lower tidal volumes do not impair oxygenation during pressure controlled one-lung ventilation, Sungur Ulke, Z., et al., O-49

On-pump coronary bypass surgery, Neutrophil activation and oxidative stress: a comparative study between off-pump and on-pump coronary artery bypass grafting, Durak, P., et al., O-42

Opioids, Comparison of two different total intravenous techniques for off-pump coronary artery bypass graft surgery: a randomized double-blind study, Askin, T., et al., P-56

Outcome

- , Impact of autologous shed mediastinal blood reinfusion on clinical outcome in the early postoperative period after cardiac surgery, Andrejaitiene, J., et al., P-39
- , Prevalence of early postoperative complications and mortality in patients undergoing cardiac surgery, Musiał-Świątkiewicz, V., et al., P-25
- , Risk factors for selected early postoperative complications after cardiac surgery, Musiał-Świątkiewicz, V., et al., O-65
- , Association of postoperative troponin T release and 12-month mortality after adult cardiac surgery, Lurati Buse, G., et al., O-63
- , Coronary artery surgery with mini-CPB system vs. off-pump technique: comparison in terms of haemodilution and postoperative outcome. A prospective randomized study, Maestre Hittinger, M.L., et al., P-16
- , Preoperative high sensitivity C-reactive protein predicts cardiovascular morbidity after coronary artery bypass grafting, Balciunas, M., et al., P-30

–, Troponin I is an independent predictor of long term outcome after off-pump coronary surgery, Timmermans, A., et al., O-64

–, Low haematocrit during cardiopulmonary bypass is not associated with adverse outcome after coronary bypass surgery, Senay, S., et al., O-62

–, Determinants of postoperative bleeding after coronary bypass surgery, Senay, S., et al., O-05

–, Perioperative allogeneic blood transfusion is associated with increased thirty day mortality after isolated coronary bypass surgery, Senay, S., et al., O-61

–, Predictors of postoperative length of stay of CABG patients: comparison of diabetics and non-diabetics, Najafi, M., et al., O-59

–, Preoperative statin treatment is not associated with reduced postoperative stratified mortality in patients undergoing aortic valve replacement, Reynaud, S., et al., P-33

–, Health related quality of life following cardiac surgery – correlation with EuroSCORE, Colak, Z., et al., O-25

Oxidative stress, An investigation of oxidative stress and DNA damage in patients exposed to cardiopulmonary bypass surgery, Mordeniz, C., et al., O-40

Oxygenation, Continuous positive airway pressure at 10 cm H₂O during cardiopulmonary bypass does not improve postoperative oxygenation in paediatric patients, Kim, J.-T., et al., P-37

Paediatric cardiac anaesthesia

- , Continuous positive airway pressure at 10 cm H₂O during cardiopulmonary bypass does not improve postoperative oxygenation in paediatric patients, Kim, J.-T., et al., P-37
- , Propofol-dexmedetomidine and propofol-ketamine combinations for anaesthesia in paediatric patients undergoing transcatheter atrial septal closure, Koruk, S., et al., O-30
- , Use of audio-Doppler ultrasonography guidance for paediatric internal jugular vein catheterization: effect on experience of the operator, Akpek, E.A., et al., O-18
- , Continuous monitoring of ScvO₂ in paediatric patients undergoing cardiac surgery – the pediasat technology, Ranucci, M., et al., O-19
- , Use of sildenafil for treatment of pulmonary hypertension after paediatric cardiac surgery: case report, Altuntas, Y., et al., P-50
- , The effects of sevoflurane and desflurane on brain tissue injury markers in cyanotic children undergoing open heart surgery, Kanbak, M., et al., P-23

Paediatric cardiac surgery

- , Aprotinin is superior to tranexamic acid in paediatric cardiac surgery, Breuer, T., et al., O-34
- , Very early extubation in children after cardiac surgery, Karahan, N., et al., P-36
- , Cardiac troponin I in paediatric cardiac operations with a minor degree of surgical myocardial damage, Iantcheva, I., et al., P-28

Pain

- , Perioperative predictors of postoperative pain at coronary artery bypass grafting, Siciliano, T., et al., O-57
- , Comparison of the effects of different fentanyl protocols on postoperative pain control after coronary artery surgery, Tur, H., et al., P-42
- , Thoracic paravertebral block with bupivacaine 0.25% versus intrathecal morphine for pain relief after thoracotomy, Kocabas, S., et al., O-46
- , Does pregabalin improve the analgesia in cardiac surgery? A preliminary study, de Cosmo, D., et al., P-40

Paravertebral block, Thoracic paravertebral block with bupivacaine 0.25% versus intrathecal morphine for pain relief after thoracotomy, Kocabas, S., et al., O-46

- Pharmacokinetic infusion models**, Comparison of Schnider and Marsh pharmacokinetic models for propofol during anaesthesia induction in elective cardiac surgery, Viterbo, J.F., et al., O-32
- Physical exercise**, The effects of physical exercise on plasma levels of relaxin, NTproANP, and NTproBNP in patients with ischaemic heart disease, Heringlake, M., et al., O-23
- Platelet function monitoring**, In vitro platelet function of platelet concentrates transfused to high risk cardiac surgery patients as determined by bedside impedance aggregometry, Jambor, C., et al., O-02
- Platelet transfusion**
- , In vitro platelet function of platelet concentrates transfused to high risk cardiac surgery patients as determined by bedside impedance aggregometry, Jambor, C., et al., O-02
 - , Comparison of clinical response to platelet transfusion and bedside impedance aggregometry, Jambor, C., et al., O-03
- Point of care aggregometry**
- , In vitro platelet function of platelet concentrates transfused to high risk cardiac surgery patients as determined by bedside impedance aggregometry, Jambor, C., et al., O-02
 - , Comparison of clinical response to platelet transfusion and bedside impedance aggregometry, Jambor, C., et al., O-03
- Postconditioning**, Postconditioning reduced infarct size in an open-chest porcine acute ischaemia-reperfusion model, Haarup Lie, R., et al., O-73
- Preconditioning**, Pharmacological preconditioning effect of isoflurane in the model of cardiac ischaemia-reperfusion in guinea pigs, Sahin, N., et al., O-74
- Preemptive analgesia**, Effects of bilateral intrapleural block as pre-emptive analgesia on haemodynamic changes in coronary artery graft surgery, Mansouri, M., et al., P-57
- Pregabalin**, Does pregabalin improve the analgesia in cardiac surgery? A preliminary study, de Cosmo, D., et al., P-40
- Pressure wave form analysis**, Comparison of pressure waveform analysis with thermodilution via pulmonary artery catheter (PAC) in valve replacement surgery, Kazanci, D., et al., P-34
- Priming of cardiopulmonary bypass**
- , Crystalloid (Ringer's solution) versus colloid 6% HES (130/04) for pump priming in CABG surgery, Sogut, F., et al., O-38
 - , Priming of the cardiopulmonary bypass with HES 130/0.4 compared to lactated Ringer's solution reduces volume load but does not change subcutaneous tissue, Schiferer, A., et al., O-37
- Propofol**
- , The effects of propofol and sevoflurane on modulation of inflammation and oxidative stress in the kidney following aortic cross clamping, Rodriguez-Lopez, J.M., et al., O-48
 - , Comparison of Schnider and Marsh pharmacokinetic models for propofol during anaesthesia induction in elective cardiac surgery, Viterbo, J.F., et al., O-32
 - , Low arterial pressure after low dose propofol administration: haemodynamic and autonomic nervous system activity, Akseleod, A., et al., P-54
 - , Comparison of right ventricle haemodynamic parameters between patients anaesthetized with propofol or desflurane for off-pump CABG, Mroziński, P., et al., P-55
 - , Effects of sevoflurane and propofol on myocardial recovery in coronary surgery, Sungur Ulke, Z., et al., P-24
 - , A randomized comparison of isoflurane versus propofol TIVA in coronary artery bypass surgery: effects on postoperative cardiac troponin I release and clinical outcome, Flier, S., et al., O-69
- Prothrombin complex concentrate**, In vitro reversal of warfarin effects with prothrombin complex concentrate vs. fresh frozen plasma, Szlam, F., et al., P-10
- Pulmonary endarterectomy**, Combined general and regional anaesthesia in pulmonary endarterectomy, Rubes, D., et al., O-45
- Pulmonary function**, The effects of pulmonary ventilation during cardio-pulmonary bypass on pulmonary function after open heart surgery, Gorgu, A., et al., O-39
- Pulmonary hypertension**
- , NT-proBNP as a marker of pulmonary hypertension in cardiac surgical patients with aortic stenosis, Kozlov, I., et al., P-58
 - , Use of sildenafil for treatment of pulmonary hypertension after paediatric cardiac surgery: case report, Altuntas, Y., et al., P-50
 - , Thoracic epidural anaesthesia impairs the haemodynamic response to acute pulmonary hypertension by depressing right ventricular contractility in pigs, Rex, S., et al., O-47
 - , Effects of inhaled iloprost on right ventricular contractility and ventricular interdependence in acute pulmonary hypertension in pigs, Rex, S., et al., O-28
 - , Acute haemodynamic effects of tezosentan in rats with monocrotaline-induced pulmonary hypertension, Lourenço, A.P., et al., P-49
- Pulse contour analysis**
- , Relationship between left ventricular wall stress, estimated by a pulse contour system, and pro-B-type natriuretic peptide in cardiac surgery, Scolletta, S., et al., O-22
 - , Haemodynamic monitoring during off-pump CABG: validation of the FloTrac TM-VigileoTM System, Turani, F., et al., O-21
 - , Continuous arterial pressure waveform analysis accurately detects cardiac output in cardiac surgery: a prospective comparison with thermodilution, echocardiography and magnetic resonance techniques, Senay, S., et al., O-13
 - , Comparison of dP/dt estimates measured by PiCCO® and TOE during cardiac surgery, Velghe, D., et al., P-14
- Quality of life**
- , Quality of life in coronary artery bypass graft candidates: the SF-36 compared to the WHOQOL-BREF, Najafi, M., et al., O-26
 - , Treatment of chronic refractory angina pectoris, Dobias, M., et al., P-46
 - , Health related quality of life following cardiac surgery – correlation with EuroSCORE, Colak, Z., et al., O-25
- Real time 3D echocardiography**
- , Assessment of mitral valve pathologies with intraoperative transoesophageal echocardiography: comparison of the 2D mode with the realtime 3D technique, Koncar-Zeh, J., et al., O-16
 - , Visibility of cardiac valves using real-time 3D-transoesophageal echocardiography (RT3D-TOE) in the OR: a feasibility study, Tschernich, H., et al., O-17
- Renal cell carcinoma**, The use of real-time TOE in the intra-operative management of renal cell carcinoma, inferior vena cava thrombus and pulmonary metastasis: case report, Mahdy, S., et al., P-52
- Renal complications**
- , Preoperative risk factors for the development of acute renal failure in cardiac surgery: a preliminary study, Persi, B., et al., P-31
 - , Angiography timing and contrast media amount: new risk factors for renal dysfunction after cardiac operations, Ranucci, M., et al., O-09
- Renal failure**, Factors determining long term survival in cardiac surgery patients with acute renal failure treated with haemofiltration, Klajbor, K., et al., O-52
- Renal protection**
- , Effects of low dose dopamine on renal oxygen consumption, blood flow and glomerular filtration rate after cardiac surgery, Redfors, B., et al., O-51
 - , Does off-pump coronary artery bypass have reno-protective effect in patients with abnormal preoperative renal function?, Papagiannopoulou P., et al., P-44

Renal replacement therapy

- , Preoperative risk factors for the development of acute renal failure in cardiac surgery: a preliminary study, Persi, B., et al., P-31
- , Continuous renal replacement therapy decreases pro BNP and IL 6 in cardiac surgery patients with postoperative sepsis C 102 The effects of sevoflurane and desflurane on serum S100 β and NSE levels in cyanotic children undergoing open heart surgery, Turani, F., et al., P-45

Respiratory failure, Prevalence of early postoperative complications and mortality in patients undergoing cardiac surgery, Musiał-Świątkiewicz, V., et al., P-25

Right ventricle

- , Assessment of right ventricular functions and pulmonary haemodynamics during lung resection guided by TOE, Galal, A., et al., O-12
- , Thoracic epidural anaesthesia impairs the haemodynamic response to acute pulmonary hypertension by depressing right ventricular contractility in pigs, Rex, S., et al., O-47
- , Effects of inhaled iloprost on right ventricular contractility and ventricular interdependence in acute pulmonary hypertension in pigs, Rex, S., et al., O-28
- , Comparison of right ventricle haemodynamic parameters between patients anaesthetized with propofol or desflurane for off-pump CABG, Mroziński, P., et al., P-55

Ringer's solution

- , Crystalloid (Ringer's solution) versus colloid 6% HES (130/04) for pump priming in CABG surgery, Sogut, F., et al., O-38
- , Priming of the cardiopulmonary bypass with HES 130/0.4 compared to lactated Ringer's solution reduces volume load but does not change subcutaneous tissue, Schiferer, A., et al., O-37

Risk factors

- , Preoperative risk factors for the development of acute renal failure in cardiac surgery: a preliminary study, Persi, B., et al., P-31
- , Risk factors for selected early postoperative complications after cardiac surgery, Musiał-Świątkiewicz, V., et al., O-65
- , Determinants of postoperative bleeding after coronary bypass surgery, Senay, S., et al., O-05

Sepsis, Continuous renal replacement therapy decreases pro BNP and IL 6 in cardiac surgery patients with postoperative sepsis C 102 The effects of sevoflurane and desflurane on serum S100 β and NSE levels in cyanotic children undergoing open heart surgery, Turani, F., et al., P-45

Sevoflurane

- , The effects of propofol and sevoflurane on modulation of inflammation and oxidative stress in the kidney following aortic cross clamping, Rodriguez-Lopez, J.M., et al., O-48
- , Neutrophil activation during off-pump coronary artery bypass grafting: a comparison between sevoflurane, desflurane and total intravenous anaesthesia, Durak, P., et al., O-41
- , Possible technique of myocardial preconditioning by sevoflurane in patients undergoing coronary surgery, Zadorozh, M., et al., O-68
- , Assessment of the index of consciousness during cardiac anaesthesia, Galán, J., et al., P-02
- , Cardioprotective effect of volatile anaesthetics on mitochondrial respiratory function, Sirvinskas, E., et al., O-67
- , Effects of sevoflurane and propofol on myocardial recovery in coronary surgery, Sungur Ulke, Z., et al., P-24

Sildenafil

- , Effects of sildenafil on left ventricular hypertrophy in a porcine model, Dahl Christensen, S., et al., P-48
- , Use of sildenafil for treatment of pulmonary hypertension after paediatric cardiac surgery: case report, Altuntas, Y., et al., P-50

Skin resistance, Can skin resistance be used to evaluate the sympathetic block of a thoracic epidural blockade?, Andersen, C., et al., P-05

Smoking, Effect of smoking on bronchial mucus transport velocity assessed by the bronchoscope under total intravenous anaesthesia, Kim, Y.H., et al., P-29

Spinal cord stimulator, Spinal cord stimulator implantation for treatment of angina pectoris gave dysaesthetic pain and burning skin sensations: case reports, Andersen, C., et al., P-59

Statins, Preoperative statin treatment is not associated with reduced postoperative stratified mortality in patients undergoing aortic valve replacement, Reynaud, S., et al., P-33

Sympathectomy, Treatment of chronic refractory angina pectoris, Dobias, M., et al., P-46

Sympathic block, Can skin resistance be used to evaluate the sympathetic block of a thoracic epidural blockade?, Andersen, C., et al., P-05

Tei index, Feasibility of measuring myocardial performance index of the right ventricle in anaesthetized patients, Michaux, I., et al., O-11

Tezosentan, Acute haemodynamic effects of tezosentan in rats with monocrotaline-induced pulmonary hypertension, Lourenço, A.P., et al., P-49

Thermodilution cardiac output

- , Pulmonary artery catheter cardiac output measurements using cold or tepid injectate during spontaneous or controlled ventilation, Kirkeby-Garstad, I., et al., O-20
- , Comparison of pressure waveform analysis with thermodilution via pulmonary artery catheter (PAC) in valve replacement surgery, Kazanci, D., et al., P-34

Thoracic anaesthesia

- , Anaerobic metabolism during thoracic anaesthesia for hyperthermic intrathoracic chemotherapy, Scolletta, S., et al., O-50
- , Lower tidal volumes do not impair oxygenation during pressure controlled one-lung ventilation, Sungur Ulke, Z., et al., O-49
- , Thoracic paravertebral block with bupivacaine 0.25% versus intrathecal morphine for pain relief after thoracotomy, Kocabas, S., et al., O-46

Thrombelastography

- , Thrombelastographic analysis of recombinant factor VIIa in cardiac surgery, Prati, P., et al., O-33
- , Kaolin activated and heparinise modified TEG \dot{O} predict post-operative bleeding in neonatal cardiac surgery, Del Sarto, P., et al., P-09
- , Predictors of clot firmness after high risk cardiac surgery, Jambor, C., et al., O-01

Tidal volume, Lower tidal volumes do not impair oxygenation during pressure controlled one-lung ventilation, Sungur Ulke, Z., et al., O-49

Tissue injury markers, The effects of sevoflurane and desflurane on brain tissue injury markers in cyanotic children undergoing open heart surgery, Kanbak, M., et al., P-23

Tissue perfusion, Tissue perfusion during minimized extracorporeal circulation (MECC), Sercia, G., et al., P-53

Total intravenous anaesthesia

- , Neutrophil activation during off-pump coronary artery bypass grafting: a comparison between sevoflurane, desflurane and total intravenous anaesthesia, Durak, P., et al., O-41
- , Effect of smoking on bronchial mucus transport velocity assessed by the bronchoscope under total intravenous anaesthesia, Kim, Y.H., et al., P-29
- , Comparison of Schnider and Marsh pharmacokinetic models for propofol during anaesthesia induction in elective cardiac surgery, Viterbo, J.F., et al., O-32

- , Improved analgesic effect in peridural thoracic versus total intravenous anaesthesia during cardiac surgery: role of epicardial fat-secreted cytokines, Cappelletti, M., et al., P-41
- , Comparison of two different total intravenous techniques for off-pump coronary artery bypass graft surgery: a randomized double-blind study, Askin, T., et al., P-56
- , A randomized comparison of isoflurane versus propofol TIVA in coronary artery bypass surgery: effects on postoperative cardiac troponin I release and clinical outcome, Flier, S., et al., O-69

Tracheal cuff pressure, The effects of deep hypothermic circulatory arrest on tracheal tube cuff pressure dynamics, Rubes, D., et al., P-08

Tranexamic acid, Aprotinin is superior to tranexamic acid in paediatric cardiac surgery, Breuer, T., et al., O-34

Transfusion

- , Aortic valve replacement with a mini-extracorporeal circulation circuit, Flo, A., et al., O-35
- , Perioperative allogeneic blood transfusion is associated with increased thirty day mortality after isolated coronary bypass surgery, Senay, S., et al., O-61

Tricuspid annular dilation, Transoesophageal echocardiography and tricuspid annular dilation: a prospective comparison with surgical measurement. A preliminary study, Parra, V., et al., P-13

Troponin I

- , Troponin I is an independent predictor of long term outcome after off-pump coronary surgery, Timmermans, A., et al., O-64
- , Cardiac troponin I in paediatric cardiac operations with a minor degree of surgical myocardial damage, Iantcheva, I., et al., P-28

Troponin T, Association of postoperative troponin T release and 12-month mortality after adult cardiac surgery, Lurati Buse, G., et al., O-63

Valvular pathology, Visibility of cardiac valves using real-time 3D-transoesophageal echocardiography (RT3D-TOE) in the OR: a feasibility study, Tschernich, H., et al., O-17

Venous catheterization, Use of audio-Doppler ultrasonography guidance for paediatric internal jugular vein catheterization: effect on experience of the operator, Akpek, E.A., et al., O-18

Venous oxygen saturation, Continuous monitoring of ScvO₂ in paediatric patients undergoing cardiac surgery – the pediasat technology, Ranucci, M., et al., O-19

Ventilation, airway pressure

- , Features of ventilator in surgical treatment of lung echinococcosis patients, Miyerbekov, Y., et al., P-35
- , The effects of pulmonary ventilation during cardio-pulmonary bypass on pulmonary function after open heart surgery, Gorgu, A., et al., O-39

Volatile anaesthetics, The effects of sevoflurane and desflurane on brain tissue injury markers in cyanotic children undergoing open heart surgery, Kanbak, M., et al., P-23