

- Thangathurai, D., 730, 898
 Theusinger, O.M., 1
 Thompson, A., 568, 629
 Thunberg, C.A., 656
 Tibble, A.D., 883
 Tini, L., 921
 Tisol, W.B., 519
 Tolani, K., 644
 Tonini, S., 867
 Torp, P., 37
 Tousignant, C., 275
 Trace, C.L., 780
 Tranquilli, M., 776
 Tritapepe, L., 210
 Tulman, G., 644
 Turi, S., 219
- Ueno, M., 730
 Umakanthan, R., 568
 Ural, N., 119
 Urzúa, J., 210
- Vahanian, A., 752
 Valfrè, C., 43
 van Barneveld, L.J.M., 422
 van Straten, A.H.M., 413
 van Zundert, A.J., 413
 Varghese, R., 890
- Veerhoek, D., 422
 Verstappen, F., 413
 Vidal-Melo, M.F., 1002
 Virzo, I., 265, 898
 Vitale, D., 210
 Vitzthum, K., 730
 Vonk, A.B.A., 422
 Vuddagiri, V., 314
- Waikar, H.D., 532
 Wallace, J., 921
 Wallace, S., 939
 Wartier, D.C., 560, 964
 Wasnick, J.D., 121, 196
 Wax, D.B., 927
 Weibel, L., 257
 Weiss, A., 988
 Wells, C., 73
 Welsby, I., 633
 Welsby, I.J., 427
 Wexler, S., 210
 Willcox, T., 164
 Williams, G., 471
 Williams, G.D., 834
 Williams, M., 633
 Wong, G.T.C., 790
 Woosenraft, D., 519
 Wyckoff, T., 892
- Xavier, A., 712
 Xia, V.W., 80
- Yamada, T., 306
 Yamazaki, T., 898
 Yanfang, C., 99
 Yang, B.-X., 1022
 Yang, C., 210
 Yared, J.-P., 580
 Yeh, C., 467
 Yoganathan, T., 69
 Young, M., 513
 Yu, C., 913
 Yuan, H., 1022
 Yue, Y., 3
- Zaky, S.S., 447
 Zambon, M., 51
 Zangrillo, A., 51, 219, 265, 586, 898, 931, 969
 Zhang, M., 644
 Zhang, R.-J., 3
 Zhang, W., 99
 Zhao, Y., 814
 Zhou, J., 314
 Zollinger, A., 257

SUBJECT INDEX

- AbioMed AB5000, 660
 AbioMed Biventricular Support 5000, 659–660
 Abscess, aortic, in dental infections, 513–515
 Acdesine, for deep hypothermic circulatory arrest, 648
 Accessory leaflet, in mitral valve, 890–891
 N-Acetylcysteine
 for acute lung injury, 686
 for hemostasis, 13
 Acid-base management, in deep hypothermic circulatory arrest, 650–651
 Acute lung injury, after thoracic surgery, 681–690
 vs. acute respiratory distress syndrome, 681–682
 etiology of, 681–682
 incidence of, 681
 prognosis for, 686–687
 reduction of
 anesthesia for, 682–683
 blood transfusions for, 683
 fluid therapy for, 683
 intraoperative management for, 682–683
 strategies for, 685–686
 ventilation techniques for, 683–685
 risk factors for, 682
 treatment of, 686
 Acute respiratory distress syndrome, vs. acute lung injury, 681–682
 Air embolism
 in left ventricle, after aortic cannulation, 824–827
 after radiofrequency ablation for atrial fibrillation, 716–718
 Albumin, in priming solutions, clot formation and, 422–426
 Alcohol users, CABG surgery in, morbidity and mortality in, 580–585
 Allodynia, after thoracic surgery, analgesia techniques effects on, 608–616
- Alpha-state management, in deep hypothermic circulatory arrest, 651
 Ambler risk score, for preoperative risk assessment, 692
 AMEDEO information resource, 851–852
 American Society of Anesthesiologists, patient information site of, 851
 American Society of Thoracic Surgeons database, for blood product use, 814–816
 Aminocaproic acid, 875–882
 Amiodarone, for ventricular fibrillation prevention, in CABG surgery, 239–243
 Aneurysm, aortic
 ascending, 198–201, 776–779
 descending, 525–527
 repair of
 in coagulopathy, 828–830
 prediction of, 84–90
 transthoracic two-dimensional echocardiography for, 37–42
 wall motion in, 776–779
 Angiosarcoma, cardiac, coronary artery pseudoaneurysm in, 633–635
 Angiotensin-converting enzyme inhibitors, for CABG surgery, 10
 Animal studies
 alveolar damage in mechanical ventilation, for thoracic surgery, 617–623
 desflurane for preconditioning, 91–98
 isoflurane-enhanced St. Thomas solution, 99–103
 ketamine for neuroprotection, 132
 levosimendan pathologic effects, 624–628
 neurocognitive function after cardiopulmonary bypass, 293–299
 noble gases for cardioprotection, 143–163
 Antiaoptotic agents, for kidney failure, 997
 Anticoagulants
 monitoring of, in sternotomy, 469–470
 for noncardiac surgery, 674

- Anticoagulants (*Continued*)
 after VAD implantation, 673
 Antidiuretic hormone. *See* Vasopressin
 Antiphospholipid antibodies, anticoagulant management with, 469–470
 Antiplatelet agents
 for acute coronary syndrome, 12
 for coronary stent thrombosis prevention, 722–727
 Aorta
 abscess of, in dental infections, 513–515
 aneurysm of
 ascending, 198–201, 776–779
 descending, 525–527
 repair of
 in coagulopathy, 828–830
 prediction of, 84–90
 transthoracic two-dimensional echocardiography for, 37–42
 wall motion in, 776–779
 ascending, diseases of, 198–201
 cannulation of, left ventricular air in, 824–827
 dissection of, 198–201
 cardiopulmonary bypass in, 887–889
 descending, 524–525
 in pregnancy, 116–118
 with preservation of aortic arch, 719–721
 malformations of, in heterotaxy, 839
 retained sponge in, 314–315
 surgery on
 deep hypothermic circulatory arrest, blood loss control during with, 73–79
 neuroprotection in, 316–321
 with tranexamic acid vs. aprotinin, 959–963
 Aortic arch
 diseases of, 201–204
 dissection near, 719–721
 reconstruction of, neuroprotection for, 201–203
 Aortic valve
 calcification of, in Gaucher disease, 471–474
 endocarditis of
 native, 516–518
 prosthetic, 513–515
 normal vs. prolapse or mass, 367–368
 regurgitation of
 classification of, 1016–1017
 imaging of, 373
 repair of, 1016–1020
 repair of, advances in, 1016–1020
 replacement of, 364–365
 anticoagulation for, 831–833
 cardiac output monitoring in, 303–305
 in elderly persons, 18–24, 193–195
 fenoldopam mesylate for, kidney flow and, 58–62
 left atrial appendage thrombus in, 977–979
 liver transplantation with, 285–292
 percutaneous method for, 25–29
 with polycythemia vera, transoxygenator pressure gradient in, 104–108
 in rheumatic disease, 191–192
 transcatheter, 8–9, 498–505, 691–699, 752–761
 stenosis of
 CABG surgery with, 712–713
 imaging of, 372–373
 replacement for, 25–29
 in rheumatic disease, 191–192
 Aortic valve (*Continued*)
 statin effects on, 893
 surgery on
 statins in, 921–926
 with venoarterial extracorporeal membrane oxygenation, 946–951
 Apoptosis, levosimendan effects on, 624–628
 Aprotinin
 for aortic surgery, with deep hypothermic circulatory arrest, 73–79
 suspension of license for, surgeons' opinions of, in United Kingdom, 418–421
 for valvular surgery, blood product use with, 959–963
 withdrawal of, anesthesia practice after, 875–882
 Argatroban, for cardiac surgery, 831–833
 Argon, for cardioprotection, 150
 Arrhythmias, cardiac
 after Fontan procedure, 126
 in heterotaxy syndrome, 841
 Arterial pressure, variation of, fluid therapy based on, 487–497
 Arterial pressure waveform analysis
 vs. cardiac index, 257–264
 vs. thermodilution method, 250–256, 767–771
 Aspergillosis, in tricuspid valve obstruction, 506–512
 Aspirin, for coronary stent thrombosis prevention, 722–727
 Asplenia syndrome, 838, 842
 Atrial appendage, left, transesophageal echocardiography for, 475–477
 Atrial arrhythmias, postoperative, 447–450
 Atrial fibrillation
 after cardiac surgery, 952–958
 after Fontan procedure, 126
 radiofrequency ablation of, air embolism after, 716–718
 in tricuspid regurgitation, 1013–1015
 Atrial flutter, after Fontan procedure, 126
 Atrial natriuretic peptide, for kidney protection, 13
 Atrial septal defects, transcatheter closure of, 374
 Atrioesophageal fistula, after radiofrequency ablation for atrial fibrillation, 716–718
 Atrium
 left, dissection of, in mitral valve repair, 817–819
 left appendage of, thrombus of, 977–979
 right
 aspergillosis of, 506–512
 enlargement of, 1013–1015
 thrombus in, in liver transplantation, 109–111
 Awareness, intraoperative, with fast-track anesthesia, 785–789
 Axillary artery, cannulation of, for neuroprotection, in aortic arch reconstruction, 202
 Banding, pulmonary artery, for single ventricle, 122–123
 Barbiturates, for deep hypothermic circulatory arrest, 647
 Beta-blockers, for deep hypothermic circulatory arrest, 648
 Bicuspid valve, aortic, repair of, 1016–1020
 Bioimpedance device, for cardiac output monitoring, 762–766
 Biopsy, lung, with clozapine therapy, hypotension in, vasopressin for, 467–468
 Bispectral index, for brain monitoring, 544–549
 Bivalirudin, for aortic aneurysm repair, 828–830
 Biventricular assist device, for heart failure, in pheochromocytoma, 985–987
 Blalock-Taussig procedure, 121–122
 Bland-White-Garland syndrome, in pregnancy, 112–115
 Bleeding and blood loss. *See also* Transfusion
 in aortic repair, with deep hypothermic circulatory arrest, control of, 73–79
 in ECMO, 170

- Bleeding and blood loss. *See also* Transfusion (*Continued*)
 endothelial leukocyte adhesion molecule polymorphisms in, 427–433
 during or after cardiac surgery, control of, 13
 in VAD implantation, 666
- Blood gas measurement
 in deep hypothermic circulatory arrest, 650–651
 in one-lung ventilation, with epidural anesthesia, 456–462
 in ventilation weaning, 451–455
- Blood transfusion. *See* Transfusion
- Boolean operators, for literature searches, 850
- Brain. *See also subjects starting with* Cerebral
 cooling of, in deep hypothermic circulatory arrest, 651
 dysfunction of, after cardiopulmonary bypass, 293–299
 ischemia of, pathophysiology of, 644–645
 monitoring of, in cardiac surgery, 544–549
 protection of. *See also* Deep hypothermic circulatory arrest
 in aortic arch reconstruction, 201–203
 in aortic surgery, 316–321
 ketamine for, 131–142
 statins for, 893
- Brain natriuretic peptide
 after aortic valve replacement, 25–29
 in cardiac transplantation, 939–945
- Breathing exercises, before thoracic surgery, 686
- Bronchodilators, before thoracic surgery, 685
- Bronchus, shunt impinging on, after tetralogy of Fallot correction, 369–370
- Bundle-branch block, left, resynchronization/defibrillation implantation for, outcomes of, 30–36
- Bupivacaine, with one-lung ventilation, pulmonary shunt fraction and, 456–462
- CABG surgery. *See* Coronary artery bypass graft (CABG) surgery
- Calcification, valvular, in Gaucher disease, 471–474
- Calcium antagonists, for deep hypothermic circulatory arrest, 648
- Cancellation, of surgery, in hypertension, 927–930
- Capnographic cardiogenic oscillations, during cardiopulmonary bypass, 369–370
- Carbon dioxide, measurement of
 in deep hypothermic circulatory arrest, 650–651
 in ventilation weaning, 451–455
- Cardiac arrest
 transesophageal echocardiography for, 184
 vasopressin for, 337–338
- Cardiac arrhythmias
 after Fontan procedure, 126
 in heterotaxy syndrome, 841
- Cardiac index
 validation of, in cardiac surgery, 265–269
 vasopressin effects on, 340
- Cardiac output, fluid therapy based on, 490
- Cardiac output monitoring
 arterial pressure waveform analysis for, 250–256, 257–264, 767–771
 endotracheal vs. thermodilution, 762–766
 FloTrac device for, 706–711, 767–771
 methods for, response time of, 306–308
 pulse contour method for, in aortic valve replacement, 303–305
 thermodilution vs. endotracheal, 762–766
- Cardiac Risk Index, Revised, for mortality prediction, in aortic aneurysm repair, 84–90
- Cardiac surgery. *See also specific procedures (Continued)*
 in alcohol users, morbidity and mortality in, 580–585
- Cardiac surgery. *See also specific procedures (Continued)*
 for angiosarcoma, 633–635
 anticoagulation for, problems with, 831–833
 aortic valve implantation, 752–761
 aprotinin license suspension for, surgeons' opinions of, in United Kingdom, 418–421
 aprotinin withdrawal for, 875–882
 atrial arrhythmias after, 447–450
 atrial fibrillation after, 952–958
 bleeding after, endothelial leukocyte adhesion molecule polymorphisms in, 427–433
 blood product use in, temporal changes in, 814–816
 brain monitoring in, 544–549
 brain natriuretic peptide biomarker in, 939–945
 bypass priming solutions for, 422–426
 cardiac index validation in, 265–269
 cardiac output monitoring in
 arterial pressure waveform analysis for, 257–264, 767–771
 FloTrac device for, 706–711
 methods for, 762–766
 combined procedures for, 356–363
 continuous ultrafiltration during, 63–68
 coronary sinus catheter for, 746–751
 delirium after
 dispositional optimism and, 560–567
 risk factors for, 555–559
 in elderly persons, risks of, 18–24
 endovascular. *See* Endovascular techniques
 epicardial three-dimensional echocardiography during, 43–50
 with epidural analgesia, outcomes of, 586–597
 fast-track anesthesia for, intraoperative awareness in, 785–789
 fenoldopam mesylate for, kidney flow and, 58–62
 in Gaucher disease, 471–474
 heparin in, thrombocytopenia due to, 69–72
 with hereditary spherocytosis, 636–638
 with history of post-traumatic stress disorder, 964–968
 hydroxyethyl starch for
 balanced, 399–407
 plasma-adapted, 394–398
 immediate extubation after, length of stay and, 780–784
 kidney injury in, 990–1001
 left atrial appendage thrombus in, 977–979
 left ventricular rupture and left atrial dissection after, 817–819
 levosimendan for, mortality risk and, 51–57
 in liver disease, continuous ultrafiltration during, 63–68
 liver transplantation with, 285–292
 minimally invasive
 coronary sinus catheter for, 746–751
 pacing pulmonary artery catheter for, 568–573
 overview of 2009, 7–17
 pacing pulmonary artery catheter for, 568–573
 with polycythemia vera, transoxygenator pressure gradient in, 104–108
 postoperative pain management in, 808–813, 867–874
 pulmonary artery hypertension management in, 797–801
 pulmonary embolism in, 309–311
 ramipril discontinuation or continuation with, vasopressin and, 230–238
 remifentanyl for, myocardial damage reduction due to, 790–796
 with renal failure, outcomes of, 574
 resynchronization/defibrillation implantation, outcomes of, 30–36
 for single ventricle, 127–128
 statins in
 for kidney injury reduction, 913–920

- Cardiac surgery. *See also specific procedures (Continued)*
 for mortality reduction, 921–926
 systolic anterior motion tolerance after, 883–884
 transcatheter valve repair, 8–9
 transesophageal echocardiography in, inadequate visualization and reporting of, 280–284
 valvular. *See* Valvular heart disease, surgery for
 with venoarterial extracorporeal membrane oxygenation, 946–951
 ventilation weaning after, 451–455
 ventricular fibrillation in, amiodarone for, 239–243
- Cardiogenic shock
 oxygen saturation measurement in, 434–439
 venoarterial extracorporeal membrane oxygenation and, 946–951
- Cardiomyopathy, Takotsubo, during liver transplantation, 629–632
- Cardioplegia solution, emulsified isoflurane with, for cardioprotection, 99–103
- Cardioprotection. *See* Myocardial protection; Preconditioning
- Cardiopulmonary bypass
 in aortic dissection, 887–889
 brain monitoring in, 544–549
 capnographic cardiogenic oscillations during, 369–370
 delirium after
 dispositional optimism and, 560–567
 risk factors for, 555–559
 diastolic function in, milrinone for, 244–249
 in elderly persons, 18–24
 heparin in, thrombocytopenia due to, 69–72
 hypotension after, vasopressin for, 338–339
 levosimendan for, mortality risk and, 51–57
 in liver disease, continuous ultrafiltration during, 63–68
 neurocognitive function after, 293–299
 with polycythemia vera, transoxygenator pressure gradient in, 104–108
 priming solutions for, fibrin clot formation and, 422–426
 separation from, after VAD implantation, 669–670
 troponin release after, 931–938
- Cardiotomy, hypotension after, vasopressin for, 338–339
- Cardioversion, atrial fibrillation after, 952–958
- Carotid arteries
 iatrogenic injury of, 322–324
 tumor of, removal of, regional anesthesia for, 312–313
- Catheters
 for aortic valve replacement, 8–9
 central venous, 119–120, 434–437
 coronary sinus, for minimally invasive valve surgery, 746–751
 pulmonary artery. *See* Pulmonary artery catheter
- Cavopulmonary connection, in Fontan procedure, 124–127
- Central venous catheter
 insertion of, internal jugular vein evaluation for, 119–120
 for oxygen saturation, circulatory failure, 434–439
- CentriMag device, for right ventricular assist, 673
- Cerebral blood flow, ketamine effects on, 132–138
- Cerebral metabolic rate for oxygen, ketamine effects on, 132–138
- Cerebral perfusion
 antegrade, for aortic arch reconstruction, 202
 in CABG surgery, intra-aortic balloon pump effects on, 300–302
 with deep hypothermic circulatory arrest, 649
 retrograde
 for aortic arch reconstruction, 201–202
 for neuroprotection, 317–320
 selective antegrade, for neuroprotection, 316–320
- Cerebrospinal fluid drainage, for spinal cord protection, in aortic aneurysm repair, 526
- Cesarean section, aortic dissection diagnosed during, 116–118
- Ciliary dysfunction, 837, 839
- Circulatory arrest, for surgery. *See* Deep hypothermic circulatory arrest
- CircuLite Synergy device, 676
- Cirrhosis
 cardiac surgery in, continuous ultrafiltration during, 63–68
 vasopressin for, 335
- Clinical Evidence resource, 848
- Clopidogrel
 for acute coronary syndrome, 12
 for coronary stent thrombosis prevention, 722–727
- Clot formation, in cardiopulmonary bypass, priming solutions effects on, 422–426
- Clozapine, lung biopsy with, hypotension in, vasopressin for, 467–468
- Coagulation, hydroxyethyl starch preparation effects on, 394–398
- Coagulopathy
 in aortic aneurysm repair, 828–830
 in ECMO, 169
 hemostatic components in, 408–412
- Cochrane Database of Systematic Reviews, 847
- Cognitive function
 after cardiac surgery, optimism and, 560–567
 with history of post-traumatic stress disorder, 964–968
- Computed tomography
 for anomalous origin of left coronary artery, 112–115
 for atriopharyngeal fistula, 716–718
 for cardiac aspergillosis, 506–512
 for infective endocarditis, 374–375
 for tracheal laceration, 519–522
- Computerized coronary angiography, for ischemia risk assessment, 348–355
- Conduction abnormalities, in heterotaxy syndrome, 841
- Confusion Assessment Method, for delirium, after cardiac surgery, 555–559
- Congenital heart disease
 accessory mitral valve leaflet, 193–195
 carotid artery surgery in, regional anesthesia for, 312–313
 double-chambered right ventricle, 356–363
 etomidate use in, hemodynamic effects of, 802–807
 in heterotaxy syndrome, 834–844
 imaging of, 372
 left atrial appendage, 475–477
 pulmonary artery hypertension in, inhaled drugs for, 797–801
 surgery for, single-ventricle physiology, 121–130
 tetralogy of Fallot, childhood correction of, 369–370
- Congestive heart failure, vasopressin for, 335
- Conivaptan, 334
 adverse effects of, 340
 for syndrome of inappropriate antidiuretic hormone, 335
- Cooling phase, of deep hypothermic circulatory arrest, 651
- CoreValve revalving system, for transcatheter implantation, 500–502
- Coronary angiography, computerized, for ischemia risk assessment, 348–355
- Coronary artery(ies)
 anomalous origin of, endocarditis with, 516–518
 left, anomalous origin of, in pregnancy, 112–115
 pseudoaneurysm of, in angiosarcoma, 633–635
 stents in, thrombosis prevention in, 722–727
 vasospasm of, during liver transplantation, 629–632
- Coronary artery bypass graft (CABG) surgery
 in alcohol users, morbidity and mortality in, 580–585
 anticoagulation management in, with antiphospholipid antibodies, 469–470
 with aortic stenosis, 712–713
 aortic valve appearance during, 367–368

- Coronary artery bypass graft (CABG) surgery (*Continued*)
 aortic valve replacement with, in elderly persons, 18–24
 atrial fibrillation after, 952–958
 blood product use in, temporal changes in, 814–816
 brain monitoring in, 544–549
 coronary computerized tomography before, 350
 delirium after, risk factors for, 555–559
 with double-chambered right ventricle repair and mitral valve replacement, 356–363
 in elderly persons, 18–24
 with epidural analgesia, outcomes of, 586–597
 fenoldopam mesylate for, kidney flow and, 58–62
 with hereditary spherocytosis, 636–638
 with history of post-traumatic stress disorder, 964–968
 levosimendan for, mortality risk and, 51–57
 liver transplantation with, 285–292
 minimally invasive, ventilation techniques for, 602–607
 off-pump
 arterial pressure waveform analysis in, 767–771
 coronary computerized tomography before, 351–353
 diastolic function in, milrinone for, 244–249
 intra-aortic balloon pump with, 300–302
 vs. on-pump, 11
 ventilation weaning after, 451–455
 outcomes of, 10–11
 with polycythemia vera, 104–108
 pulmonary embolism during or before, 309–311, 639–640
 ramipril discontinuation or continuation with, vasopressin and, 230–238
 remifentanyl for, myocardial damage reduction due to, 790–796
 with renal failure, outcomes of, 574
 statins in, for kidney injury reduction, 913–920
 techniques for, 10–11
 transfusion after, risk factors for, 413–417
 with venoarterial extracorporeal membrane oxygenation, 946–951
 ventricular fibrillation in, amiodarone for, 239–243
- Coronary sinus catheter, for minimally invasive valve surgery, 746–751
- Corticosteroids
 for deep hypothermic circulatory arrest, 647–648
 before thoracic surgery, 685–686
- C-reactive protein, in aortic dissection, 524
- Cryoprecipitate, clot formation and, 408–412
- Cyanosis, after Fontan procedure, 126
- Cystoscopy, with left ventricular assist device, 1002–1009
- Cytokines, in bleeding, 427–433
- DARE (Database of Reviews of Effects), 847–848
- Databases, for anesthesia information, 845–858
- 1-Deamino-8-D-arginine vasopressin. *see* Desmopressin
- Deep hypothermic circulatory arrest, 644–655
 acid-base management during, 650–651
 adjunctive protection in, 648–649
 alternatives to, 644
 for aortic arch reconstruction, 202–203
 benefits of, 644
 cooling in, 651
 disadvantages of, 644
 duration of, 652
 history of, 644
 hypothermia action in, 646–647
 monitoring during, 649–650
 neurologic injury caused by, 652
 pathophysiologic considerations in, 644–645
- Deep hypothermic circulatory arrest, 644–655 (*Continued*)
 pharmacologic protection in, 647–648
 rewarming in, 651–652
 temperature management in, 651–652
- Defibrillators
 for bundle-branch block, 30–36
 lead extraction for, 974–976
- Delirium, after cardiac surgery
 dispositional optimism and, 560–567
 risk factors for, 555–559
- Demeclocycline, for syndrome of inappropriate antidiuretic hormone, 335
- Dental infections, endocarditis due to, 513–515
- Depression, before cardiac surgery, delirium risk and, 560–567
- Desflurane
 for preconditioning, 91–98
 for thoracic surgery, 682–683
- Desmopressin, 333
 for diabetes insipidus, 334
 for perioperative bleeding, 336
 for von Willebrand disease, 335–336
- Developing countries, publication trends in, 969–973
- Dexamethasone, for deep hypothermic circulatory arrest, 647–648
- Dexmedetomidine, for deep hypothermic circulatory arrest, 648
- Diabetes insipidus, desmopressin for, 334
- Dialysis, CABG surgery with, outcomes of, 574
- Diastole, aortic valve appearance during, 367–368
- Diastolic function, in CABG surgery, milrinone for, 244–249
- DICOM (Digital Imaging and Communication in Medicine), 859–866
- Digital storage, of echocardiographic images, 859–866
- Dissection
 aortic, 198–201
 cardiopulmonary bypass in, 887–889
 descending, 524–525
 in pregnancy, 116–118
 with preservation of aortic arch, 719–721
 carotid artery, iatrogenic, 322
- Duraheart LVAS, 676
- Echocardiography
 electronic image storage in, 859–866
 epiaortic, for wall motion detection, in aneurysm, 776–779
 epicardial, for pulmonary embolism, 309–311
 intraoperative, pediatric transthoracic probe for, 43–50
 for kidney flow measurement, fenoldopam mesylate effects on, 58–62
 three-dimensional, 43–50, 371–378
 transesophageal. *See* Transesophageal echocardiography
 transthoracic two-dimensional, for aortic aneurysm repair, 37–42
- ECMO (extracorporeal membrane oxygenation), for severe cardiac and respiratory failure, 164–172
- Edema, pulmonary, after thoracic surgery, 681–690
- Editorials
 Can Meta-analysis of the Current Literature Help Determine if Perioperative β -Blockers Improve Outcome of High-Risk Patients Undergoing Noncardiac Surgery?, 217–218
 The Cardiothoracic and Vascular Anesthesia Session at the 2009 Annual Meeting of the Chinese Society of Anesthesiologists, 3–4
 Combined Valvular Disease: When Echocardiography Provides the Questions and the Answers, 366
 East Joins West, 1–2

Editorials (*Continued*)

- The New-Generation Hydroxyethyl Starch Solutions: The Holy Grail of Fluid Therapy or Just Another Starch?, 389–393
- Ordinary Images--Extraordinary Stories: Echo Challenges and Clinical Decisions, 5–6
- Percutaneous Coronary Sinus Catheterization for Minimally Invasive Cardiac Surgery--More Questions Than Answers?, 743–745
- Statins and Cardiac Surgery, 909–912
- Stuck With a Decision: What Is the "True" Aortic Valve Area--Anatomic, Geometric, or Effective Orifice Area?, 714–715
- Systolic Anterior Motion and Mitral Valve Reserve Function: Which One, Should We Care About?, 885–886
- 2010 Standard of Care for Central Nervous System Monitoring During Cardiac Surgery, 541–543

Education

- on information management, 857
- on transesophageal echocardiography, 183–190
- Edwards valve, transcatheter implantation of, 499–500
- Edwards-Cribier valve, transcatheter implantation of, 501
- Edwards-Sapien valve, transcatheter implantation of, 501

Elderly persons

- aortic dissection in, 201
- aortic valve replacement in, 193–195
- cardiac surgery in
 - delirium risk factors for, 555–559
 - risks of, 18–24
- transesophageal echocardiography for, 184

Electroencephalography

- for brain monitoring, 544–549
- for deep hypothermic circulatory arrest, 650

Elephant trunk approach, to aortic stent placement, 203–204

EMBASE database, 848–849

Embolism. *See also* Pulmonary embolism

- air
 - in left ventricle, after aortic cannulation, 824–827
 - after radiofrequency ablation for atrial fibrillation, 716–718
- gas, in ECMO, 168–169
- intracardiac, in liver transplantation, 109–111
- from left ventricle, 196–197

eMedicine knowledge base, 849

Emphysema, subcutaneous, from tracheal laceration, 519–522

Emulsified isoflurane, with St. Thomas solution, for cardioprotection, 99–103

Endocarditis

- of aortic valve, 516–518
- imaging of, 374–375
- of prosthetic valves, 513–515

Endoscopic techniques

- for aortic valve replacement, 25–29
- saphenous vein harvesting, 10–11
- valvular replacement, 8–9

Endothelial leukocyte adhesion molecule polymorphisms, in bleeding, 427–433

Endotracheal bioimpedance cardiac output monitoring, 762–766

Endotracheal tube

- double-lumen, performance comparison for, 598–601
- removal of
 - epoprostenol for, 988–989
 - immediate, length of stay after, 780–784
- tracheal laceration from, 519–522

Endovascular techniques

Endovascular techniques (*Continued*)

- for aortic aneurysm repair, 84–90, 526–527
- for aortic arch stent placement, 203–204
- for aortic dissection repair, 524–525
- for aortic valve implantation, 752–761
- for valve surgery, coronary sinus catheter for, 746–751
- Epi-aortic echocardiography, for wall motion detection, in aneurysm, 776–779
- Epicardial echocardiography
 - pediatric transthoracic probe for, 43–50
 - for pulmonary embolism, 309–311
- Epidural anesthesia or analgesia
 - for cardiac surgery, outcomes of, 586–597
 - with one-lung ventilation, pulmonary shunt fraction and, 456–462
 - for thoracotomy, allodynia after, 608–616
- Epoprostenol, for extubation, in ventricular assist device implantation, 988–989
- Erythropoietin, for kidney failure, 997
- Esmolol, for noncardiac surgery, 219–229
- Esophagectomy
 - intensive care for, 440–446
 - transesophageal echocardiography after, 700–705
- Esophagus, fistula of, after radiofrequency ablation for atrial fibrillation, 716–718
- Etomidate, hemodynamic response to, in cardiac shunt lesions, 802–807
- EuroSCORE, for preoperative risk assessment, 691–692
- Evidence-based medicine, literature searches for, 845–858
- Excitotoxicity, mechanisms of, 131–132
- Extracorporeal membrane oxygenation
 - for cardiac surgery, 946–951
 - for left ventricular assist, 483–484
 - for severe cardiac and respiratory failure, 164–172
- Extubation
 - epoprostenol for, 988–989
 - immediate, length of stay after, 780–784
- Facial anomalies, in heterotaxy syndrome, 842
- Factor VII, for hemostasis, 13
- Factor VIIa, for aortic surgery, 828–830, 959–963
- Fast-track anesthesia, for cardiac surgery, intraoperative awareness in, 785–789
- Femoral arterial pressure waveform analysis, 257–264
- Fenoldopam, for kidney protection, 13, 58–62
- Fentanyl, postoperative delirium due to, 555–559
- Fetus, heterotaxy syndrome in, 836
- Fibrinogen, deficiency of, hemostatic components effects on, 408–412
- Fibrinolysis, vs. primary percutaneous coronary intervention, 11
- Fibrosis, of intracardiac leads, 974–976
- Fistulas
 - atrioesophageal, after radiofrequency ablation for atrial fibrillation, 716–718
 - carotid artery, 324
 - subclavian, 325–326
 - tracheo-innominate, 325
- FloTrac device, for cardiac output monitoring, 706–711, 767–771, 772–775
- Fluid therapy
 - balanced hydroxyethyl starch for, 399–407
 - goal-directed, based on arterial pressure variation, 487–497
 - plasma-adapted hydroxyethyl starch for, 394–398
 - responsiveness to, stroke volume variation and, 772–775
 - for thoracic surgery, 683
- Fontan procedure, 124–127, 173–182
- Fresh frozen plasma, clot formation and, 408–412

- Gabapentin, for postoperative pain, 808–813
- Gas embolism, in ECMO, 168–169
- Gastrointestinal anomalies, in heterotaxy syndrome, 841
- Gaucher disease, cardiac surgery in, 471–474
- Gelatin, in priming solutions, clot formation and, 422–426
- Genetic factors, antiplatelet drug resistance, 724–725
- Glenn shunt, for single ventricle, 124
- Glycemic control, perioperative, 14
- Glycogen synthase kinase-3 β , in cardioprotection, 151–153
- Google Alerts information resource, 852
- Head cooling, in deep hypothermic circulatory arrest, 651
- Health Information Access to Research Initiative project, 856
- Heart
- embolism in, during liver transplantation, 109–111
 - in heterotaxy syndrome, 834–844
 - surgery on. *See* Cardiac surgery; *specific procedures*
 - transplantation of
 - brain natriuretic peptide biomarker in, 939–945
 - donor heart evaluation for, 351
 - after failed Fontan procedure, 127
- Heart failure
- extracorporeal membrane oxygenation for, 164–172
 - after Fontan procedure, 126
 - in pheochromocytoma, biventricular assist device for, 985–987
 - resynchronization/defibrillation implantation for, outcomes of, 30–36
 - vasopressin for, 335
 - ventricular assist devices for
 - left. *See* Left ventricular assist devices
 - right, 673
- HeartMate I (extended lead vented electric VAD), 658–659
- HeartMate II left VAD, 662–663
- HeartMate III, 676
- HeartWare HVAD, 676
- Helium, for cardioprotection, 149–157
- Hematomas, intramural, in aortic dissection, 200
- Hemodilution, clot formation and, 422–426
- Hemodynamics
- in CABG surgery, with ramipril, 230–238
 - in congenital shunt lesions, etomidate effects on, 802–807
 - instability of, in ECMO, 170
 - renal, in cardiac surgery, fenoldopam mesylate effects on, 58–62
 - in transcatheter valve replacement, 694–695
- Hemorrhagic shock, vasopressin for, 336
- Hemostasis, perioperative, 13
- Heparin
- thrombocytopenia induced by, aortic aneurysm repair in, 828–830
 - for VAD implantation, 666
- Hepatorenal syndrome, tolvaptan for, 226
- Hereditary spherocytosis, cardiac surgery with, 636–638
- Heterotaxy syndrome, 834–844
- High-frequency jet ventilation
- vs. one-lung ventilation, for CABG surgery, 602–607
 - for thoracic surgery, 685
- Histamine levels, heparin effects on, in cardiac surgery, 69–72
- Hybrid procedures
- for aortic aneurysm repair, 526–527
 - for aortic arch repair, 204
 - operating room for, 8–9, 693
 - for single ventricle, 124
- Hydroxyethyl starch
- balanced, 399–407
 - plasma-adapted, 394–398
- Hydroxyethyl starch (*Continued*)
- in priming solutions, clot formation and, 422–426
 - responsiveness to, stroke volume variation and, 772–775
- Hypertension
- pulmonary artery
 - inhaled drugs for, 797–801
 - in ventricular assist device implantation, 988–989
 - systemic, preoperative, 927–930
- Hypoglycemia, in glucose management, 14
- Hypokalemia, control of, in liver transplantation, 80–83
- Hyponatremia, vasopressin for, 334–335
- Hypoperfusion, vasopressin for, 336–339
- Hypoplastic left heart syndrome, 121
- Hypotension
- heparin-induced, in cardiac surgery, 69–72
 - in liver transplantation, methylene blue for, 453–466
 - in lung biopsy, with clozapine therapy, vasopressin for, 467–468
- Hypothermia, for surgery. *See* Deep hypothermic circulatory arrest
- Hypoxemia, in ECMO, 170
- Iatrogenic injury, of supra-aortic arteries, 322–329
- Imaging, 371–378. *See also specific imaging techniques*
- Immune-mediated hypotension, heparin-induced, in cardiac surgery, 69–72
- Impella circulatory support system, 481–483
- Impella Recover LP 2.5, 660–661
- Implanted cardioverter-defibrillator, laser-lead extraction for, 974–976
- Infants, newborn
- heterotaxy syndrome in, 836–837
 - single ventricle in, first-stage repairs for, 121–124
- Infections, in ECMO, 170–171
- Information retrieval and management, 845–858, 859–866
- Innominate artery, iatrogenic injury of, 325–326
- Insulin
- for deep hypothermic circulatory arrest, 648
 - for glycemic control, 14
 - for hypokalemia prevention, in liver transplantation, 80–83
- Intensive care
- for cardiac surgery, immediate extubation effects on, 780–784
 - ECMO use in, 167–168
 - for esophagectomy, 440–446
 - statin use in, 894–896
- Intercellular adhesion molecule, in bleeding, 427–433
- Interleukins, in bleeding, 427–433
- Intra-aortic balloon pump
- for myocardial infarction, 11–12
 - for off-pump CABG, cerebral perfusion benefits in, 300–302
- Ischemia, brain, protection against, ketamine for, 131–142
- Isoflurane, emulsified, with St. Thomas solution, for cardioprotection, 99–103
- Isoproterenol, for right ventricular dysfunction, 672
- Jarvik 2000 Flowmaker, 663–664
- Journal of Cardiothoracic and Vascular Anesthesia*, publication trends in, 969–973
- Jugular veins
- internal, echocardiography probe for, 119–120
 - oxygen saturation of, in deep hypothermic circulatory arrest, 650
- Ketamine, for neuroprotection, 131–142
- Kidney
- dysfunction or failure of
 - CABG surgery with, outcomes of, 574
 - new developments in, 990–1001

Kidney (*Continued*)

- non-volume--responsive, 993
- pathophysiology of, 996–997
- prevention of, 13–14
 - fenoldopam mesylate for, 58–62
 - statins for, 893–894, 913–920
- renal blood flow in, 993
- treatment of, 997–998
- tubular injury in, 995–996
- volume-responsive, 991–993
- hydroxyethyl starch effects on, 399–407
- protection of, in aortic aneurysm repair, 525

Laser-lead extraction, for implanted cardioverter-defibrillator, 974–976

Leads, for defibrillators, extraction of, 974–976

Lee Revised Cardiac Risk Index, for mortality prediction, in aortic aneurysm repair, 84–90

Left ventricular assist devices, 656–680

- anesthetic management for, 664–666
- anticoagulation after, 673
- cardiopulmonary bypass separation after placement, 669–670
- classification of, 656–664
- complications of, 675
- first-generation, 657–660
- new technology for, 675–676
- noncardiac procedures after implantation, 673–675
- outcomes of, 673–675
- percutaneous, 478–486
 - extracorporeal membrane oxygenation, 483–484
 - Impella circulatory support system, 481–483
 - rationale for, 479
 - TandemHeart, 479–481
- potential patients for, 656
- preoperative assessment for, 664–665
- right ventricular dysfunction after, 671–673
- second-generation, 660–664
- selection of, 656, 665–666
- third-generation, 657, 676
- transesophageal echocardiography for, 666–669
- types of, 656

Leg, embolization to, from left ventricle, 196–197

Leipzig Fast-Track Concept, for cardiac surgery, intraoperative awareness in, 785–789

Length of stay, immediate extubation effects on, 780–784

Letters to the editor

- Anesthetic Management of a Patient with Giant Right Atrial Myxoma, 532–533
- Any Rationale for Propofol Use in Cardiac Surgery?, 903
- Argatroban "Reversal" Is Caused by Nonphysiologic Stimulation of Coagulation Not Activated Factor VII, 1026–1027
- Assessing Agreement in Cardiac Output Monitoring Validation Studies, 741–742
- Assessment of Neurocognitive Function and Neuroprotective Strategies in Cardiac Surgery, 536–537
- Bispectral Index Monitor Prevented Intraoperative Awareness During Partial Cardiopulmonary Bypass, 740
- Cardiac Insufficiency: A Critical Analysis of the Current Publication Procedures Under Quantitative and Qualitative Aspects, 731–734
- Cetacaine-Induced Complication During Transesophageal Echocardiography Placement, 736
- Congenital Pleuropericardial Fistula Complicating Awake Cardiac Surgery, 901–903

Letters to the editor (*Continued*)

- Considerations About the Association Between Preoperative Statin Administration and Outcomes in Patients Undergoing CABG, 904–906
- Coopdech Bronchial Blocker Is Useful in Abnormalities of the Tracheobronchial Tree, 735–736
- Coronary Stents and Antiplatelet Therapy: Patients' and Health Professionals' Understanding of the Implications, 383–384
- Decreased Nitric Oxide Products in the Urine of Patients Undergoing Cardiac Surgery, 213–215
- Epidural Anesthesia for Cesarean Section in a Patient With Severe Mitral Stenosis and Pulmonary Hypertension, 1022–1023
- Excessive Use of Phenylephrine by the Perfusionist and Hemodynamic Instability, 899
- Failure of Action of Neuromuscular Blocking Agents in a Patient With Severe Vasoconstriction Caused by Infused Vasopressors Following a Mitral Valve Replacement, 533–534
- Failure of Double-Lumen Tube Cuff Deflation on the Cuff Leak Test, 1025
- Guidewire Entrapment in a Tricuspid Valve Apparatus, 898–899
- Iatrogenic Oropharyngeal Injury and Hemorrhage Requiring Blood Transfusions During Insertion of Transesophageal Echocardiographic Probes in Patients Undergoing Cardiac Surgery, 1023–1024
- Intraoperative Use of Activated Recombinant Factor VII for Refractory Bleeding in Complicated Lung Transplantations, 382–383
- Levosimendan and Cardiac surgery, 210
- Levosimendan and Platelet Function, 532
- Missing Sponge: Clinical Signs, 534
- Monitoring Anticoagulation of Patients With Antiphospholipid Antibodies on Cardiopulmonary Bypass, 734–735
- A Novel Technique for Easy Identification of the Subclavian Vein During Ultrasound-Guided Cannulation, 210–211
- One-Lung Ventilation Can Alter the Severity of Aortic Regurgitation, 736–738
- The Parker Flex-Tip Tube Is Useful in a Bougie-Assisted Endotracheal Tube Exchange After Lung Lavage, 901
- Postoperative Pain After Cardiac Surgery, 1025–1026
- The Predictability of Blood Loss with Thromboelastometry, 381–382
- Pulmonary Artery Catheter Use with Left-Sided Active Endocarditis: Deal or No Deal?, 903–904
- Remifentanyl for Awake Thoracoscopic Bullectomy, 386–387
- Removal of Pulmonary Artery Catheter Knotted During Placement by Using Transesophageal Echocardiography, 1027–1028
- In Response to "A Modified 'Over-the-Needle' Technique for Arterial Cannulation," 384
- A Rounded Image Inside the Left Ventricle: The Mechanism of the Artifact Formation, 384–385
- A Simple Method for Obtaining a "Ramped" Laryngoscopy Position After Anesthetic Induction, 537–539
- Surgery for Pseudoaneurysm of the Ascending Aorta: Role of Intraoperative 2-Dimensional and Real-time 3-Dimensional Transesophageal Echocardiography, 906–908
- Surgical Management of the Calcified Patent Ductus Arteriosus Under Hypotensive Anesthesia, 739–740
- Temperature Management During Cardiopulmonary Bypass, 385–386

- Letters to the editor (*Continued*)
- Transesophageal Echocardiography in Detecting Tricuspid Valve Pathology in an Intensive Care Unit Patient, 211–213
 - Transesophageal Echocardiography in Identifying Fungal Vegetations in the Right Cardiac Chambers in an Intensive Care Unit Patient, 535–536
 - Transesophageal Echocardiography in the Identification of Subtle Aortic Dissection in a Trauma Patient, 900–901
 - Use of the Proseal Laryngeal Mask Airway and Arndt Bronchial Blocker for Lung Separation in a Patient With a Tracheal Mass and Aspiration Risk, 215–216
 - Utility of Intraoperative Three-Dimensional Transesophageal Echocardiography in a Patient With Embolized Atrial Septal Defect Closure Device for Emergency Cardiac Surgery, 730–731
 - Which Protein C After Cardiac Surgery?, 738–739
- Levacor LVAS, 676
- Levitronix CentriMag blood pump, 661–662
- Levosimendan
- for apoptosis attenuation, 624–628
 - for cardiac surgery, mortality statistics with, 51–57
 - for preconditioning, 9–10
 - for right ventricular dysfunction, 673
- Lidocaine, for deep hypothermic circulatory arrest, 648
- LIFEBRIDGE system, for circulatory support, 484
- Literature
- publication of, in *Journal of Cardiothoracic and Vascular Anesthesia*, 969–973
 - searches of, 845–858
- Liver
- disorders of
 - cardiac surgery in, continuous ultrafiltration during, 63–68
 - in heterotaxy syndrome, 841–842
 - transplantation of
 - cardiac surgery with, 285–292
 - hyperkalemia prevention during, 80–83
 - intracardiac embolism during, 109–111
 - pulmonary embolism during, 641–643
 - Takotsubo cardiomyopathy and coronary vasospasm during, 629–632
 - vasoplegia syndrome in, methylene blue for, 453–466
- Lixivaptan, 334
- Lobectomy
- fluid responsiveness measurement in, 772–775
 - postoperative pain management in, 867–874
- Loeys-Dietz syndrome, aortic disorders in, 200
- Losartan, for aortic disease, in Marfan syndrome, 199–200
- Lung
- acute injury of, after thoracic surgery, 681–690
 - alveolar damage in, in thoracic surgery, with ventilation, 617–623
 - anomalies of, in heterotaxy syndrome, 842
 - biopsy of, hypotension in, vasopressin for, 467–468
 - embolism in. *See* Pulmonary embolism
 - resection of
 - acute lung injury after, 681–690
 - fluid responsiveness measurement in, 772–775
 - postoperative pain management in, 867–874
- Lysosomal storage disease, cardiac surgery in, 471–474
- Magnesium, for deep hypothermic circulatory arrest, 648
- Magnetic resonance imaging
- for anomalous origin of left coronary artery, 112–115
 - cardiac, 371–378
- Malperfusion, in aortic dissection, 200–201
- Mannitol, for deep hypothermic circulatory arrest, 648
- Marfan syndrome, aortic disease in, 199–201
- MD Consult resource, 849–850
- Medical Subject Headings, 850
- Medline, 848
- MedlinePlus, 851
- Megakaryocytes, stimulation of, 13
- MESH (Medical Subject Headings), 850
- Meta-analysis
- of epidural analgesia for cardiac surgery, 586–597
 - of esmolol for ischemia prevention, 219–229
- N-Methyl-D-aspartate receptor antagonists, neurotoxic effects of, 133, 136
- Methylprednisolone
- for deep hypothermic circulatory arrest, 647–648
 - before thoracic surgery, 685–686
- MicroMed DeBakey axial flow pump, 664
- Midline defects, in heterotaxy syndrome, 842
- Milrinone
- for diastolic function, in CABG surgery, 244–249
 - inhaled, for pulmonary artery hypertension, 797–801
 - for right ventricular dysfunction, 672
- Minimally invasive cardiac surgery, valvular
- coronary sinus catheter for, 746–751
 - pacing pulmonary artery catheter for, 568–573
- Mitogen-activated protein kinase, in cardioprotection, 146–148
- Mitral valve
- accessory leaflet in, 890–891
 - calcification of, in Gaucher disease, 471–474
 - endocarditis of, in dental infections, 513–515
 - leaflet abnormalities in, 1010–1012
 - regurgitation of
 - due to accessory valve leaflet, 193–195
 - imaging of, 374
 - repair of
 - coronary sinus catheter for, 746–751
 - fenoldopam mesylate for, kidney flow and, 58–62
 - left ventricular rupture and left atrial dissection after, 817–819
 - systolic anterior motion tolerance after, 883–884
 - transcatheter, 9
 - with tricuspid valve repair, 375
 - replacement of, 364–365
 - anticoagulation for, 831–833
 - with CABG surgery and double-chambered right ventricle repair, 356–363
 - coronary sinus catheter for, 746–751
 - left atrial appendage thrombus in, 977–979
 - in rheumatic disease, 191–192
 - stenosis of, in rheumatic disease, 191–192
 - surgery on
 - statins in, 921–926
 - troponin release after, 931–938
 - transcatheter repair for, 374
- Mixed venous oxygen saturation, measurement of, in circulatory failure, 434–439
- Morphine, postoperative, gabapentin influence on, 808–813
- Mortality
- in aortic dissection, in elderly persons, 201
 - aprotinin use and, 875–882
 - after CABG surgery, blood transfusion and, 413–417
 - in CABG surgery
 - in alcohol users, 580–585
 - with kidney failure, 574
 - in cardiac surgery

- Mortality (*Continued*)
 in elderly persons, 18–24
 levosimendan effects on, 51–57
 resynchronization/defibrillation implantation, 30–36
 statins and, 892–896
 with venoarterial extracorporeal membrane oxygenation, 946–951
 in deep hypothermic circulatory arrest, 652
 in endovascular aortic aneurysm repair, prediction of, 84–90
 in esophagectomy, 440–446
 in noncardiac surgery, esmolol effects on, 219–229
 preoperative statin therapy and, 921–926
 in pulmonary embolism, during liver transplantation, 641–643
 reduction of, transesophageal echocardiography for, 184–185
- Motor-evoked potentials, for spinal cord protection, in aortic aneurysm repair, 525–526
- Mozavaptan, 334
- Multicenter Evaluation of Tolvaptan Effect on Remodeling trial, 335
- Myocardial infarction, primary percutaneous coronary interventions for, 11–12
- Myocardial ischemia
 in cardiac transplantation, brain natriuretic peptide biomarker for, 939–945
 levosimendan attenuation of, 624–628
 in noncardiac surgery, esmolol effects on, 219–229
 for preconditioning, 10
 risk assessment for, coronary computerized angiography for, 348–355
 statins effects on, 892–896
- Myocardial performance index, of right ventricle, 270–274
- Myocardial protection. *See also* Preconditioning
 noble gases for, 143–163
 remifentanyl for, 790–796
 St. Thomas cardioplegia solution with isoflurane for, 99–103
 statins for, 893
- Myocardial reperfusion
 brain natriuretic peptide in, 939–945
 salvage kinase pathway in, 147–149, 151–152
 ventricular fibrillation in, 239–243
- National Guidelines Clearinghouse, 848
- Near-infrared spectroscopy
 for cerebral perfusion, in CABG surgery, 300–302
 for deep hypothermic circulatory arrest, 650
- Neon, for cardioprotection, 149–150
- Nesiritide, for right ventricular dysfunction, 672–673
- Neurocognitive function, after cardiopulmonary bypass, 293–299
- Neurologic disorders, in ECMO, 171
- Neuroprotection. *See also* Deep hypothermic circulatory arrest
 in aortic aneurysm repair, 828–830
 in aortic arch reconstruction, 201–203
 in aortic surgery, 316–321
 ketamine for, 131–142
 statins for, 893
- Newborns
 heterotaxy syndrome in, 836–837
 single ventricle in, first-stage repairs for, 121–124
- Nitric oxide
 for acute lung injury, 686
 for right ventricular dysfunction, 672
- Nitroglycerin, inhaled, for pulmonary artery hypertension, 797–801
- Noble gases, for cardioprotection, 143–163
- Noncardiogenic pulmonary edema, after thoracic surgery, 681–690
- Norwood procedure, 123–124
- Novacor II LAD, 676
- Off-pump CABG. *See* Coronary artery bypass graft (CABG) surgery, off-pump
- One-lung ventilation
 epidural anesthesia with, pulmonary shunt fraction and, 456–462
 vs. high-frequency jet ventilation, for CABG surgery, 602–607
 performance comparison for, 598–601
 stroke volume variation vs. fluid responsiveness in, 772–775
 for thoracic surgery, alveolar damage in, 617–623
- Optimism, before cardiac surgery, delirium risk and, 560–567
- Outcomes. *See also* Mortality
 of aortic valve repair, 1016–1020
 of aprotinin withdrawal, 875–882
 of CABG surgery, 10–11
 in alcohol users, 580–585
 with amiodarone, 239–243
 with history of post-traumatic stress disorder, 964–968
 with kidney failure, 574
 with milrinone, 244–249
 with ramipril, 230–238
 of cardiac surgery
 with epidural analgesia, 586–597
 with hydroxyethyl starch, 399–407
 with liver transplantation, 285–292
 resynchronization/defibrillation implantation, 30–36
 statins and, 892–896
 with venoarterial extracorporeal membrane oxygenation, 946–951
 of coronary stent thrombosis, 722–727
 of deep hypothermic circulatory arrest, 652
 of endovascular aortic aneurysm repair, prediction of, 84–90
 of esophagectomy, 440–446
 of fast-track anesthesia, 785–789
 of Fontan procedure, 125
 of gabapentin for postoperative pain, 808–813
 of immediate extubation, 780–784
 of left ventricular assist device implantation, 675
 of mitral valve surgery, troponin release and, 931–938
 of noncardiac surgery, esmolol effects on, 219–229
 preanesthesia hypertension effects on, 927–930
 of remifentanyl myocardial protection, 790–796
 of statin therapy
 postoperative, 913–920
 preoperative, 921–926
 of transcatheter aortic valve implantation, 752–761
- L-2-Oxothiazolidine-4-carboxylate, for acute lung injury, 686
- Oxygen, measurement of
 in deep hypothermic circulatory arrest, 650–651
 for ventilation weaning, 451–455
- Oxygen saturation of jugular vein, in deep hypothermic circulatory arrest, 650
- Oxygen therapy, for acute lung injury, 686
- Oxygenation, in one-lung ventilation, with epidural anesthesia, 456–462
- Pacing pulmonary artery catheter, for minimally invasive valve surgery, 568–573
- Paravertebral blockade, for thoracotomy, 867–874
- Partial carbon dioxide rebreathing, for cardiac output monitoring, during resuscitation, 306–308
- Pediatric patients. *See also* Congenital heart disease
 cardiac surgery in
 with Gaucher disease, 471–474
 pulmonary artery hypertension management in, 797–801
 heterotaxy syndrome in, 834–844
 single ventricle in, first-stage repairs for, 121–124

- PediVAS, 676
- Pentastarch, for kidney protection, 14
- Percutaneous techniques. *See also* Transcatheter procedures
 aortic valve replacement, cardiac output monitoring in, 303–305
 coronary interventions, for myocardial infarction, 11–12
 for left ventricular assist devices, 478–486
 valve surgery, aortic valve replacement, 25–29
- Pericardial tamponade, in lead extraction, 974–976
- Pheochromocytoma, heart failure due to, biventricular assist device for, 985–987
- pH-stat management, in deep hypothermic circulatory arrest, 651
- PIER (Physician Information Education Resource), 848
- Plasma-adapted hydroxyethyl starch, 394–398
- Platelet(s)
 clot formation and, 408–412
 inhibitors of
 for acute coronary syndrome, 12
 for coronary stent thrombosis prevention, 722–727
- Pneumomediastinum, from tracheal laceration, 519–522
- Pneumectomy, acute lung injury after, 681–690
- Polycythemia vera, cardiac surgery with, transoxygenator pressure gradient in, 104–108
- Polysplenia syndrome, 838, 842
- Positioning, for acute lung injury, 686
- Positive end-expiratory pressure
 for Fontan procedure, 125
 for thoracic surgery, 684
- Postoperative pain
 in cardiac surgery, gabapentin for, 808–813
 in thoracotomy, 867–874
- Post-traumatic stress disorder, coronary artery surgery in, 964–968
- Potassium channels, in cardioprotection, 147, 151
- Prasugrel, for acute coronary syndrome, 12
- Preconditioning
 desflurane for, 91–98
 ischemic, 9–10
 levosimendan for, 51–57
 noble gases in, 143–163
 pharmacologic, 9–10
 St. Thomas cardioplegia solution with isoflurane for, 99–103
- Preeclampsia, anomalous origin of left coronary artery with, 112–115
- Pregnancy
 anomalous origin of left coronary artery in, 112–115
 aortic dissection in, 116–118
- Preload, fluid therapy based on, 487–489
- Pressure recording analytic method, 257–264
- Primary ciliary dyskinesia disorders, 837, 839
- Progenitor cells, for kidney failure, 997–998
- Prone positioning, for acute lung injury, 686
- Propofol, for deep hypothermic circulatory arrest, 647
- Prostacyclin, for right ventricular dysfunction, 672
- Prosthetic valves, endocarditis of, 513–515
- Protection
 brain. *See* Brain, protection of
 kidney, statins for, 913–920
 myocardial. *See also* Preconditioning
 remifentanyl for, 790–796
 statins for, 893
 spinal cord, in aortic aneurysm repair, 828–830
- Protein losing enteropathy, after Fontan procedure, 125–127
- P-selectin, in bleeding, 427–433
- Pseudoaneurysms
 coronary artery, in angiosarcoma, 633–635
 supra-aortic arteries, 322–329
- Publication trends, in *Journal of Cardiothoracic and Vascular Anesthesia*, 969–973
- PubMed Clinical Queries, 848, 854–855
- Pulmonary artery
 diseases of, after Fontan procedure, 126–127
 left coronary artery origin in, in pregnancy, 112–115
 in single ventricle physiology, 123–125
- Pulmonary artery catheter
 for cardiac output monitoring
 during resuscitation, 306–308
 vs. thermodilution method, 250–256
 uncalibrated, 257–264
 pacing, for minimally invasive valve surgery, 568–573
- Pulmonary artery hypertension
 inhaled drugs for, 797–801
 in ventricular assist device implantation, 988–989
- Pulmonary edema, after thoracic surgery, 681–690
- Pulmonary embolism
 in cardiac surgery, 309–311
 after Fontan procedure, 127, 173–182
 intraoperative diagnosis of, 639–640
 during liver transplantation, 641–643
 thrombolysis for, 980–984
- Pulmonary shunt fraction, in one-lung ventilation, with epidural anesthesia, 456–462
- Pulse contour method, for cardiac output monitoring, in aortic valve replacement, 303–305
- Pulse-pressure variations, fluid therapy based on, 492–493
- Pumps, for ECMO, 165–166
- Radial arterial pressure waveform analysis, 257–264
- Radiofrequency ablation, for atrial fibrillation, air embolism after, 716–718
- Radiography, for tracheal laceration, 519–522
- Ramipril, discontinuation or continuation of, with cardiac surgery, vasopressin and, 230–238
- Rapid ventricular pacing, for valve replacement, 695
- Red blood cell(s), augmentation of, for CABG surgery, 10
- Relcovaptan, 339
- Remacemide, for deep hypothermic circulatory arrest, 648
- REMATCH study, of left ventricular assist device implantation, 675
- Remifentanyl
 for CABG surgery, myocardial damage reduction due to, 790–796
 with one-lung ventilation, pulmonary shunt fraction and, 456–462
 for thoracotomy, allodynia after, 608–616
- Reperfusion, myocardial
 brain natriuretic peptide in, 939–945
 salvage kinase pathway in, 147–149, 151–152
 ventricular fibrillation in, 239–243
- Reperfusion injury salvage kinase pathway, in cardioprotection, 147–149, 151–152
- Resistance, to antiplatelet drugs, for coronary stents, 722–727
- Respiratory support, extracorporeal membrane oxygenation for, 164–172
- Respiratory tract, anomalies of, in heterotaxy syndrome, 842
- Respiratory variations, in arterial pulse pressure, fluid therapy based on, 490–491
- Resuscitation, cardiac output monitoring in, methods for, response time of, 306–308
- Resynchronization/defibrillation implantation, for heart failure, outcomes of, 30–36
- Retrograde transcatheter procedures, for aortic valve implantation, 503
- Revascularization. *See also specific techniques*
 techniques for, 12–13

- Revised Cardiac Risk Index, for mortality prediction, in aortic aneurysm repair, 84–90
- Rewarming, in deep hypothermic circulatory arrest, 651–652
- Rheumatic heart disease, valve replacement in, 191–192, 364–365
- Risk stratification, hypertension in, 927–930
- Sano operation, for single ventricle, 123–124
- Saphenous vein, endoscopic harvesting of, 10–11
- Satavapan, 334
- Scopus database, 849
- Searches, of anesthesia literature, 845–858
- Seizures, from air embolism, after radiofrequency ablation, 716–718
- Sepsis, in ECMO, 170–171
- Septic shock, vasopressin for, 336–337
- Sevoflurane, for thoracic surgery, 682–683
- Shock
 - from air embolism, after radiofrequency ablation, 716–718
 - hemorrhagic, vasopressin for, 336
 - septic
 - oxygen saturation measurement in, 434–439
 - vasopressin for, 336–337
 - transesophageal echocardiography for, 184
- Shunts, for single ventricle
 - Glenn procedure, 124
 - pulmonary artery, 121–124
- Sildenafil, for pulmonary embolism, 980–984
- Single ventricle, 121–130
 - general surgery with, 127–128
 - surgery for
 - failure of, 125–127
 - first-stage repairs, 121–124
 - outcome of, 125
 - physiology after, 125
 - second-stage palliation, 124
 - third-stage palliation, 124–125
 - variants of, 121
- Situs abnormalities (heterotaxy), 834–844
- Smoking, cessation of, before thoracic surgery, 685
- Social support, before cardiac surgery, delirium risk and, 560–567
- Society of Thoracic Surgeons (STS) Predicted Risk of Mortality (Prom) score, for preoperative risk assessment, 691–692
- Sodium bicarbonate, for kidney protection, 13–14
- Speckle tracking, for right ventricle function assessment, 275–279
- Spectral entropy method, for brain monitoring, 544–549
- Spherocytosis, hereditary, cardiac surgery with, 636–638
- Spinal cord, protection of, in aortic aneurysm repair, 525–526, 828–830
- Splanchnic hypoperfusion, vasopressin-induced, 339
- Spleen, anomalies of, 838
- Sponge, retained, in aorta, 314–315
- SSR-149415, for anxiety, 339
- St. Thomas cardioplegia solution, emulsified isoflurane with, for cardioprotection, 99–103
- Statins
 - postoperative, for kidney injury reduction, 913–920
 - preoperative, for mortality reduction, 921–926
 - surgical outcomes and, 892–896
- Stem cells, for kidney failure, 997–998
- Stents
 - aortic arch, 203–204
 - coronary, thrombosis of, 722–727
- Sternotomy
 - anticoagulation management in, with antiphospholipid antibodies, 469–470
- Sternotomy (*Continued*)
 - for VAD implantation, 666
- Stress, before cardiac surgery, delirium risk and, 560–567
- Stroke volume, vs. fluid responsiveness, in one-lung ventilation, 772–775
- Subclavian artery
 - coverage of, in aortic aneurysm repair, 527
 - iatrogenic injury of, 324–325
- Sufentanil, with one-lung ventilation, pulmonary shunt fraction and, 456–462
- SumSearch resource, 850
- Syndrome of inappropriate antidiuretic hormone, vasopressin for, 334–335
- Systolic anterior motion tolerance test, 883–884
- Takotsubo cardiomyopathy, during liver transplantation, 629–632
- TandemHeart percutaneous VAD, 662
- TandemHeart transeptal VAD, 479–481
- Temperature, monitoring of, in deep hypothermic circulatory arrest, 649
- Tetracyclines, for kidney failure, 997
- Tetralogy of Fallot
 - childhood correction of, bronchus impingement after, 369–370
 - repair of, imaging after, 372
- Thermodilution
 - vs. arterial pressure waveform analysis, 250–256, 767–771
 - vs. endotracheal bioimpedance device, 762–766
- Thiopental, for deep hypothermic circulatory arrest, 647
- Thoracic paravertebral block, for postoperative pain management, 867–874
- Thoracic surgery
 - acute lung injury after, 681–690
 - allodynia after, analgesia techniques effects on, 608–616
 - atrial arrhythmias after, 447–450
 - double-lumen tubes for, performance comparison for, 598–601
 - left atrial appendage discovery before, 475–477
 - overview of 2009, 7–17
 - postoperative pain management in, 867–874
 - ventilation for, alveolar damage in, 617–623
- Thoracoscopy, with left ventricular assist device, 1002–1009
- Thoracotomy
 - allodynia after, analgesia techniques effects on, 608–616
 - postoperative pain management in, 867–874
- Thoratec Dual Drive Console, 658
- Thoratec implantable VAD, 657–658
- Thoratec paracorporeal VAD, 657
- Three-dimensional transesophageal echocardiography, for bilobar left atrial appendage, 475–477
- Thrombocytopenia, heparin-induced
 - aortic aneurysm repair in, 828–830
 - in cardiac surgery, 69–72
- Thromboelastometry
 - for coagulopathy, 408–412
 - hydroxyethyl starch and, 394–398
 - for liver transplantation, 641–643
 - for priming solution studies, 422–426
- Thrombolysis, for pulmonary embolism, 980–984
- Thrombosis and thrombi. *See also* Pulmonary embolism
 - coronary stent, 722–727
 - after Fontan procedure, 127, 173–182
 - intracardiac, in liver transplantation, 109–111
 - left atrial appendage, 977–979
- Ticagrelor, for acute coronary syndrome, 12
- Tidal volumes, for thoracic surgery, 683–684
- Tolvaptan, 334
 - for heart failure, 335
 - for hepatorenal syndrome, 226
 - for variceal bleeding, 336

- Trachea, laceration of, from endotracheal tube, 519–522
- Tranexamic acid, 875–882
for aortic surgery, with deep hypothermic circulatory arrest, 73–79
for valvular surgery, blood product use with, 959–963
- Transapical procedure, for aortic valve implantation, 502–503
- Transcatheter procedures
aortic valve implantation, 752–761
coronary sinus catheter for, 746–751
pulmonary embolism thrombolysis, 980–984
for valvular heart disease, 8–9, 373–374, 498–505
anesthesia management in, 691–699
CoreValve for, 500–502
Edwards device for, 499–500
Edwards-Cribier valve for, 501
Edwards-Sapien valve for, 501
hemodynamic monitoring in, 694–695
intraoperative considerations in, 693–697
percutaneous transvenous, 498–499
postoperative considerations in, 697–698
preoperative considerations in, 691–693
retrograde, 503
transapical, 502–503
transarterial, 498–499
transesophageal echocardiography for, 695–697
transfemoral, 501–502
- Transesophageal echocardiography
for accessory mitral valve leaflet, 193–195
for aortic dissection, 116–118, 887–889
for aortic regurgitation, 1016–1020
for aortic stenosis, 712–713
of aortic valve, 367–368
for cardiac aspergillosis, 506–512
for cardiac output monitoring, during resuscitation, 306–308
for deep hypothermic circulatory arrest, 649
for emboli from left ventricle, 196–197
for endocarditis, 516–518
after esophagectomy, 700–705
inadequate visualization and reporting of results for, 280–284
for mitral valve leaflet abnormalities, 1010–1012
for myocardial performance index, 270–274
probe from, for internal jugular vein surface evaluation, 119–120
for pulmonary embolism, 639–640
for retained sponge in aorta, 314–315
for rheumatic valve disease, 191–192
for right ventricle assessment, 671
vs. speckle tracking, for right ventricular function assessment, 275–279
three-dimensional, 371–378
for bilobar left atrial appendage, 475–477
for left atrial appendage thrombus, 977–979
training for, 183–190
for tricuspid regurgitation, 1013–1015
for VAD implantation, 666–669
for valve replacement, 695–697
for valvular endocarditis, 513–515
- Transfemoral procedure, for aortic valve implantation, 501–502
- Transfusion
aprotinin use and, 875–882
after CABG surgery, risk factors for, 413–417
for CABG surgery, temporal changes in, 814–816
for hypokalemia prevention, in liver transplantation, 80–83
risks of, 13
for thoracic surgery, 683
with tranexamic acid vs. aprotinin, 959–963
- Transplantation
Transplantation (*Continued*)
heart
brain natriuretic peptide biomarker in, 939–945
donor heart evaluation for, 351
after failed Fontan procedure, 127
liver
cardiac surgery with, 285–292
hyperkalemia prevention during, 80–83
intracardiac embolism during, 109–111
pulmonary embolism during, 641–643
Takotsubo cardiomyopathy and coronary vasospasm during, 629–632
vasoplegia syndrome in, methylene blue for, 453–466
- Tricuspid valve
obstruction of, in aspergillosis, 506–512
regurgitation of, 375, 1013–1015
repair of
anticoagulation for, 831–833
coronary sinus catheter for, 746–751
replacement of, coronary sinus catheter for, 746–751
stenosis of, imaging of, 373
- Triglycyl lysine vasopressin, 333
- TRIP search system, 850
- Troponin, release of, after mitral valve surgery, 931–938
- Tumor(s), carotid artery, removal of, regional anesthesia for, 312–313
- Tumor necrosis factor, in bleeding, 427–433
- Ultrafiltration, during cardiac valve surgery, in liver disease, 63–68
- Ultrasonography, speckle tracking method for, for right ventricle function assessment, 275–279
- UpToDate expert opinion resource, 849
- VADs. *See* Ventricular assist devices
- Valvular heart disease. *See also specific valves*
leaflet abnormalities in, 1010–1012
rheumatic, valve replacement in, 191–192
surgery for
anticoagulation for, problems with, 831–833
atrial fibrillation after, 952–958
continuous ultrafiltration during, 63–68
coronary computerized tomography before, 350
in elderly persons, 18–24, 193–195
with Gaucher disease, 471–474
left atrial appendage thrombus in, 977–979
left ventricular rupture and left atrial dissection after, 817–819
levosimendan for, mortality risk and, 51–57
liver transplantation with, 285–292
minimally invasive
coronary sinus catheter for, 746–751
pacing pulmonary artery catheter for, 568–573
outcomes of, 1016–1020
in rheumatic disease, 191–192, 364–365
statins in, 893, 913–920, 921–926
systolic anterior motion tolerance after, 883–884
transcatheter, 8–9, 373–374, 498–505, 752–761
troponin release after, 931–938
- Vasoplegia syndrome, in liver transplantation, methylene blue for, 453–466
- Vasopressin, 330–347
abnormal levels of, osmolality disorders in, 334–335
adverse effects of, 339–340
analogs of, 333
antagonists of, 333–334, 340
for cardiac arrest, 337–338

- Vasopressin, 330–347 (*Continued*)
for coagulation disorders, 335–336
for hemorrhagic shock, 336
for hypoperfusion, 336–339
normal levels of, 330–331
physiology of, 330–334
ramipril discontinuation or continuation with, in CABG surgery, 230–238
receptors for, 331–332
for septic shock, 336–337
synthesis of, 330–331
for variceal bleeding, 336
for von Willebrand disease, 335–336
- Vasospasm, coronary, during liver transplantation, 629–632
- Venoarterial extracorporeal membrane oxygenation, 164–170, 946–951
- Venous desaturation, in ECMO, 170
- Venovenous extracorporeal membrane oxygenation, 164–170
- Ventilation
for Fontan procedure, 125
high-frequency jet
vs. one-lung ventilation, for CABG surgery, 602–607
for thoracic surgery, 685
one-lung. *See* One-lung ventilation
postoperative delirium due to, 555–559
for thoracic surgery, 617–623, 683–685
weaning from, after off-pump CABG surgery, 451–455
- Ventricle(s)
dysfunction of, after Fontan procedure, 127
left
air in, after aortic cannulation, 824–827
diastolic function of, after aortic valve replacement, 25–29
dysfunction of, imaging of, 373
embolization of, 196–197
- Ventricle(s) (*Continued*)
hypertrophy of, imaging of, 373
rupture of, in mitral valve repair, 817–819
transthoracic two-dimensional echocardiography of, in aortic aneurysm repair, 37–42
right
double-chambered, repair of, with CABG surgery and mitral valve replacement, 356–363
dysfunction of, after left VAD implantation, 671–673
function of, speckle tracking assessment of, 275–279
myocardial performance index of, 270–274
single, 121–130, 839
- Ventricular assist devices. *See also* Left ventricular assist devices
for heart failure, in pheochromocytoma, 985–987
implantation of, extubation in, 988–989
left. *See* Left ventricular assist devices
noncardiac surgery with, 1002–1009
perioperative management with, 820–823
right, 673
- Ventricular fibrillation
arrest of, during minimally invasive valve surgery, 568–573
in CABG surgery, amiodarone for, 239–243
- Ventricular pacing, rapid, for valve replacement, 695
- Volatile anesthetics, for preconditioning, 9, 91–98
- von Willebrand disease, desmopressin for, 335–336
- Wall motion, aortic, in aneurysm, 776–779
- Warfarin, plasma containing, hemostatic components effects on, 408–412
- World Health Organization, Health Information Access to Research Initiative project, 856
- Xenon, for cardioprotection, 143–163