

- Xia, V.W., e56, 1092
 Xia, Z., 444
 Xiong, L., 908
 Xu, J., 592
 Xu, W., 610
- Yamak, B., 637
 Yamazaki, T., 578, 575
 Yang, L., 908
- Yang, X.X., 536
 Yao, L., 444
 Yate, P., 581
 Yazigi, A., 386
 Yedlin, A., 402
 Yildiz, O., 856
 Yilmaz, A., 250
 Yilmaz, A.T., 824
 Young, G.B., 299
 Yue, Y., 288
- Zalaquett, R., 615
 Zanatta, P., 1076
 Zangrillo, A., e19, e54, 120, 389, 437, 476, 579 804, 902
 Zhang, J., 592
 Zhang, L., 625
 Zhang, X., 288
 Zhou, J., 590
 Zhou, W., 592
 Zhou, X., 625

SUBJECT INDEX

- Abiomed AbiocoR IHR, 1182–1184
 Abscess, aortic root, 192–195
 Academic performance, 761–765
 Accreditation, 761–765
 N-Acetylcysteine, for inflammation control, in cardiac surgery, 268–275
 Acromegaly, cardiovascular manifestations of, 707–708
 Activated coagulation time
 with bivalirudin, 415–418
 measurement of, in pediatric patients, 395–401
 ACTR II device, for activated coagulation time, 395–401
 Acute respiratory distress syndrome
 cardiac output measurement in, 481–485
 in tetralogy of Fallot repair, 867–873
 Adenosine receptors, in preconditioning, 444–448
 Adenosine-monophosphate--activated protein kinase, activation of, in preconditioning, 66–71
 Adrenal gland, dysfunction of, cardiovascular manifestations of, 709–710
 Aggregometry, in CABG surgery, 42–47
 Air embolism, after mitral valve replacement, 1219–1220
 Albumin
 for cardiac surgery, 847–855
 deficiency of, after off-pump CABG surgery, 462–468
 Alpha-adrenergic receptors, dexmedetomidine interactions with, 856–862
 American Society of Anesthesiologists, echocardiographic education program of, 212–215
 ϵ -Aminocaproic acid
 clinical value of, 16–19
 kidney injury and, 402–406
 side effects of, 20–25
 transfusion requirements with, 36–41
 5-Aminoimidazole-4-carboxamide-1-riboside, desflurane interactions with, 66–71
 Amiodarone, for atrial fibrillation prevention, 90–94
 Analgesia
 after cardiac surgery, 110–119
 for electrophysiology laboratory, 647–659
 Androgenic anabolic steroids, cardiovascular manifestations of, 712
 Aneurysm, aortic
 quadricuspid aortic valve with, 196–198
 repair of, 8
 endovascular, 166–176
 in Marfan syndrome, 625–631
 spinal cord injury in, 120–126
 spinal cord perfusion monitoring in, 316–319
 Angina, in anomalous origin of left coronary artery, 501–504
 Angiotensin blockers, for heart failure, 9
 Angiotensin-converting enzyme inhibitors, for hemodynamic control, 517
 Animal studies
 desflurane preconditioning, 66–71, 276–281
 myocardial protection with remifentanyl, 926–930
 preconditioning, 444–448
 reperfusion hemodynamics, 780–783
 vascular effects of JM-1232(-), 72–77
 Anticoagulants
 in cardiac surgery, 340–341
 dabigatran, 1208–1212
 Antidiuretic hormone (vasopressin), deficiency of, in liver disease, 665–670
 Antifibrinolytics
 clinical value of, 16–19
 kidney injury and, 402–406
 side effects of, 20–25
 Antiplatelet agents
 in cardiac surgery, 340–341
 monitoring of, 42–47
 Aorta
 aneurysm of. *See* Aneurysm, aortic
 ascending, diseases of, 8
 calcification of, aortic stenosis with, 368–369
 coarctation of, repair of, Shone's anomaly with, 838–840
 descending, diseases of, 8–9
 dissection of, in lung transplantation, 1121–1124
 pseudoaneurysm of, 1098–1100
 repair of
 coma after, 961–967
 delirium in, 968–974
 in Marfan syndrome, 625–631
 transfusion requirements for, 36–41
 retained broken wire in, 894–896
 rupture of, endovascular repair of, 169
 Aortic arch, diseases of, 8
 Aortic root, abscess of, 192–195
 Aortic valve
 bicuspid, 891–893
 bypass procedure for, after previous cardiac surgery, 134–139
 central calcification of, 1223–1225
 endocarditis of, fistula formation in, 751–752
 quadricuspid
 short-axis view of, 199–201
 without stenosis, 196–198
 regurgitation of
 in bicuspid valve, 891–893
 in quadricuspid aortic valve, 196–198
 repair of, 7–8
 replacement of

- Aortic valve (*Continued*)
- CABG surgery with, 368–369
 - in complicated case, 1193–1199
 - EuroSCOREs in, 1071–1075
 - left atrial dissection and hematoma after, 309–310
 - with mediastinal radiation therapy pathology, 1200–1203
 - mitral regurgitation after, 885–888
 - paravalvular leaks after, 610–614
 - percutaneous, 1109–1112
 - transcatheter, 437–443
 - transfemoral, 1036–1043
 - stenosis of
 - coronary artery disease with, 368–369
 - double orifice appearance of, 1223–1226
 - repair of, mitral regurgitation after, 885–888
 - von Willebrand disease and, 615–618
 - transcatheter implantation of, 7–8
 - warfarin ridge in, 380–381
- Aortopulmonary collateral vessels, 569–571
- Aprotinin, renal injury from, 402–406
- Argatroban, 371–375
- Arndt endobronchial blocker, for thoracic surgery, in pediatric patients, 683–686
- Arterial pressure waveform, in pulsus alternans, 567–568
- Arterial pulse power method, for passive leg raising response assessment, 48–52
- Arterial switch repair, levosimendan in, 419–424
- Arterial waveform analysis, in abnormal left ventricular function, 53–58
- Artificial heart
 - anesthetic management for, 1184–1188
 - devices for, 1180–1184
 - history of, 1179–1180
 - indications for, 1184
 - outcomes of, 1188–1189
- Aspirin, monitoring of, 42–47
- Atrial fibrillation
 - cardiac index in, 476–480
 - in cardiac surgery
 - corticosteroids for, 156–165
 - cryoablation for, 1030–1035
 - dabigatran for, 1209–1210
 - prevention of, after cardiac surgery, 90–94
 - in thoracic surgery, 817–823
- Atrial septal defect
 - mitral valve deformity with, 1106–1108
 - patch closure of, 378–379
 - surgery for, eustachian valve interfering with, 680–682
- Atrial septostomy, for pulmonary hypertension, 695
- Atrioventricular block, in off-pump CABG, 311–312
- Atrium
 - left
 - coronary sinus hematoma in, 376–377
 - dissection of, after valve replacement, 309–310, 498–500
 - fistula of, abscess at, 192–195
 - hematoma encircling, 1221–1222
 - mass in, during pneumonectomy, 833–835
 - postoperative air in, 1219–1220
 - right, pericardial patch in, 378–379
- Autonomic dysregulation, in sepsis, 531
- Bariatric surgery, cardiac performance during, 943–949
- Barrett's esophagus, dissection of, in transesophageal echocardiography, 150–152
- BART (Blood Conservation Using Antifibrinolytics in an Randomized Trial), 16–19
- Benzodiazepine derivatives, vascular effects of, 72–77
- Beta blockers
 - for atrial fibrillation prevention, 90–94
 - for endovascular aortic aneurysm repair, 170
 - for hemodynamic control, 515–518
- Bibliometrics, 761–765
- Bicuspid valve, aortic, 891–893
- Bioimpedance cardiography, for cardiac output monitoring, 770–775
- Bivalrudin, 371–375, 415–418
- BiVent tube, case studies of, 505–508
- Biventricular pacing, hemodynamics of, 238–242
- Bland-White-Garland syndrome, 501–504
- Bleeding and blood loss. *See also* Transfusions
 - antifibrinolytic therapy for, 20–25
 - conservation methods for, 16–19
 - gastrointestinal
 - after CABG surgery, 637–641
 - in transesophageal echocardiography, 678–679
 - in pediatric cardiac surgery, 987–994
 - platelet monitoring for, 42–47
 - prevention of, in cardiac surgery, 340–341
 - pulmonary, in tetralogy of Fallot repair, 313–315
 - recombinant activated factor VII for, 804–810
 - with ventricular assist devices, 10
 - in von Willebrand disease, 131–133
- Blood Conservation Using Antifibrinolytics in an Randomized Trial (BART), 16–19
- Blood gas measurement, postoperative, ear sensor for, 243–249
- Blood pressure, control of, clinical outcomes of, 509–525
- Blood transfusions. *See* Transfusions
- Bosentan, for pulmonary hypertension, 694
- Brain. *See also subjects starting with* Cerebral
 - monitoring of, in cardiac surgery, 1076–1085
 - protection of, hyperbaric oxygen for, 908–916
- Brain natriuretic peptide
 - in heart failure, 9
 - in outcome prediction, for cardiac surgery, 469–475
- Bronchoscopy, for foreign body aspiration, 1005–1008
- B-type natriuretic peptide, in outcome prediction, for cardiac surgery, 256–262, 469–475
- Bupivacaine, for epidural anesthesia, for thoracic surgery, 449–454
- CABG surgery. *See* Coronary artery bypass graft (CABG) surgery
- Calcification
 - aortic, aortic stenosis with, 368–369
 - aortic valve, 1223–1225
- Calcium channel blockers, for hemodynamic control, 517–519
- Calcium flux, in sepsis, 531–532
- Carbon dioxide, measurement of
 - postoperative, ear sensor for, 243–249
 - in robotic cardiac surgery, 85–89
- Carcinoid tumors, cardiovascular manifestations of, 708–709
- Cardiac arrest, therapeutic hyperthermia for, 362–367
- Cardiac Calendar--2011 to 2012, e3, e13, e33, e49
- Cardiac catheterization
 - anesthesia services for, 553–558
 - remifentanyl for, in pediatric patients, 660–664
- Cardiac index, in atrial fibrillation, 476–480
- Cardiac output
 - after biventricular pacing, 238–242
 - control of, clinical outcomes of, 509–525
 - increasing, in cardiac surgery, 1154–1162

- Cardiac output (*Continued*)
 low, after tetralogy of Fallot correction, 569–571
- Cardiac output monitoring
 in abnormal left ventricular function, 53–58
 in acute respiratory distress syndrome, 481–485
 endotracheal, 770–775
 hemodynamic variables in, 1044–1050
 passive leg raising and, 48–52
 plethysmography for, 1044–1050
 trending ability of, 536–546
 in tricuspid valve regurgitation, 830–832
- Cardiac surgery. *See also specific procedures*
 N-acetylcysteine antiinflammatory action in, 268–275
 acquired von Willebrand disease management in, 131–133
 activated coagulation time measurement in, 395–401
 acute respiratory distress syndrome in, 867–873
 alternative central venous catheterization route for, 1018–1023
 anticoagulants for, 415–418
 aortic valve implantation, 437–443
 arterial pulse waveform analysis in, 53–58
 atrial fibrillation in
 cryoablation for, 1030–1035
 prevention after, 90–94
 for atrial septal defect, 378–379
 biventricular pacing in, hemodynamics of, 238–242
 blood gas measurement after, 243–249
 brain monitoring in, 1076–1085
 carbon dioxide tension in, 85–89
 cardiac output monitoring in, 770–775, 776–779
 methods for, 1044–1050
 with tricuspid regurgitation, 830–832
 cefuroxime pharmacokinetics in, 425–430
 cerebral oxygen saturation in, cognitive dysfunction and, 95–104
 Clostridium difficile infections after, 263–267
 coma after, prediction of duration, 961–967
 combined CABG and valve, delirium in, 811–816
 corticosteroid protection in, 156–165
 delirium in, 811–816, 968–974
 dexamethasone for, 950–960
 early extubation in, 431–436
 fluid therapy for, 847–855, 1141–1153
 glycemic variability in, 799–803
 heparin-induced thrombocytopenia in, 371–375
 high-risk, intra-aortic balloon pump for, 632–636
 hydroxyethyl starch for, 407–414
 hyperalgesia reduction in, 917–925
 hyperbaric oxygen preconditioning for, 908–916
 hypoalbuminemia after, 462–468
 insulin during, 1086–1091
 intraoperative electrocardiographic changes in, 791–798
 kidney failure after, 619–624
 left stellate ganglion block for, 78–84
 levosimendan in, 419–424, 1154–1162
 lidocaine pharmacokinetics in, 931–936
 malignant hyperthermia, 731–735
 metabolic syndrome and, 282–287
 microcirculation observation during, 250–255
 mitral valve area assessment during, 221–228
 myocardial velocity measurements on, 216–220
 Norwood procedure, 11
 outcomes of
 benefits in, 880–80f
 lymphocyte count and, 975–980
 prediction of, B-type natriuretic peptide for, 256–262, 469–475
- Cardiac surgery. *See also specific procedures (Continued)*
 paravalvular leaks after, 610–614
 in pediatric patients. *See* Pediatric patients, cardiac surgery in
 perioperative care in, evidence-based medicine for, 335–346
 persistent eustachian valve pitfalls in, 680–682
 platelet function monitoring in, 42–47
 pleural effusion volume measurement in, 228–232
 postoperative pain in, gabapentin for, 824–829
 pulmonary hemorrhage control in, 313–315
 for pulmonary hypertension, 695
 pulse contour analysis in, 48–52
 pulse-pressure variation in, 1056–1062
 renal injury in, in pediatric patients, 402–406
 repeat
 left thoracotomy approach to, 134–139
 without systemic heparinization, 347–352
 research highlights from 2010, 6–15
 sedation and analgesia regimens for, 110–119
 seizures after, 20–25, 299–305
 splanchnic perfusion in, 642–646
 strain measurement in, 216–220
 stroke in, factor VII and, 804–810
 sublingual microcirculation in, 784–790
 superior vena cava obstruction after, 1101–1103
 transfusion requirements for, prospective observational study of,
 36–41
 vasoplegia after, methylene blue for, 736–743
 von Willebrand disease and, 615–618
- Cardiomyocyte depressing factor, in sepsis, 530–531
- Cardiomyopathy
 dilated, aortic valve replacement with, 1193–1199
 hypertrophic
 left ventricular outflow tract obstruction in, 673–677
 septal myectomy for, 306–308
 treatment of, 10
 reverse Takotsubo, 1116–1118
- Cardioprotection. *See* Myocardial protection; Preconditioning
- Cardiopulmonary bypass
 for aortic valve replacement, 1193–1199
 atrial fibrillation in, cryoablation for, 1030–1035
 biventricular pacing after, hemodynamics of, 238–242
 brain monitoring in, 1076–1085
 carbon dioxide tension in, 85–89
 cefuroxime pharmacokinetics in, 425–430
 Clostridium difficile infections after, 263–267
 corticosteroid protection in, 156–165
 delirium in, 811–816
 dexamethasone for, 950–960
 fluid therapy for, 847–855
 gastrointestinal complications after, 637–641
 with heparin-bonded circuits, for reoperations, 347–352
 intra-aortic balloon pump for, 632–636
 levosimendan for, 632–636
 lidocaine pharmacokinetics in, 931–936
 malignant hyperthermia in, 731–735
 metabolic syndrome and, 282–287
 microcirculation observation during, 250–255
 outcome of, lymphocyte count and, 975–980
 paravalvular leaks after, 610–614
 for pulmonary thrombarterectomy, 183–191
 retrograde autologous priming of, 995–999
 for robotic surgery, 723–726
 sedation and analgesia regimens for, 110–119
 splanchnic perfusion in, 642–646

- Cardiopulmonary bypass (*Continued*)
 sublingual microcirculation in, 784–790
- Carotid arteries, endarterectomy of, method comparison for, 1024–1029
- Catheter(s), pulmonary artery, for cardiac output monitoring, 770–775
- Catheterization, cardiac
 anesthesia services for, 553–558
 remifentanyl for, in pediatric patients, 660–664
- Cefuroxime pharmacokinetics, in cardiopulmonary bypass, 425–430
- Celiac artery
 blood flow through, fenoldopam effects on, 642–646
 normal appearance of, 565–566
- Central venous catheter, for cardiac surgery, 1018–1023
- Central venous pressure, in right ventricular insufficiency, 1051–1055
- Cerebral oximetry, for superior vena cava obstruction, 1101–1103
- Cerebral oxygen saturation
 in CABG surgery, cognitive dysfunction and, 95–104
 in congenital heart disease surgery, 11
 in single-lung ventilation, for lobectomy, 127–130
- Cerebral oxygenation, in cardiopulmonary bypass, retrograde autologous priming effects on, 995–999
- Cerebral protection, hyperbaric oxygen for, 908–916
- Chest pain, in anomalous origin of left coronary artery, 501–504
- Children. *See* Pediatric patients
- Cholecystitis, after CABG surgery, 637–641
- Chronotropic agents, for hemodynamic control, 519–520
- Circulatory arrest, deep hypothermic, for thromboembolic pulmonary hypertension, 746
- Cirrhosis, vasopressin deficiency in, 665–670
- Clopidogrel, monitoring of, 42–47
- Clostridium difficile infections, in intensive care, 263–267
- Coagulation factors, in cardiac surgery, stroke and, 804–810
- Coanda effect, 891–893
- Coarctation, aortic, repair of, Shone's anomaly with, 838–840
- Cognitive dysfunction
 after CABG surgery, cerebral oxygen saturation and, 95–104
 in combined CABG and valve surgery, 811–816
 metabolic syndrome and, 282–287
 prevention of, in cardiac surgery, 342–343
- Colloids, for cardiac surgery, 847–855, 1141–1153
- Coma, after cardiac surgery, prediction of duration, 961–967
- Compartment syndrome, in extracorporeal membrane oxygenation, 836–837
- Congenital heart disease
 cardiac output monitoring in, 776–779
 eustachian valve in, 680–682
 Fontan physiology, noncardiac surgery with, 329–334
 quadricuspid aortic valve, 196–198, 199–201
 surgery for
 acute respiratory distress syndrome in, 867–873
 extubation timing after, 874–879
 impaired lung function in, 1000–1004
 levosimendan in, 419–424
 mechanical valvular dysfunction in, 671–672
 pulmonary hemorrhage control in, 313–315
 treatment options for, 10–11
- Continuity equation, for mitral valve area assessment, 221–228
- Coronary artery(ies)
 dissection of, in pregnancy, 353–361
 laceration of, after thoracic spine surgery, 1095–1097
 spasm of, vasodilator-resistant, 1104–1105
- Coronary artery active perfusion system, with off-pump CABG, 311–312
- Coronary artery angiography, retained broken wire after, 894–896
- Coronary artery bypass graft (CABG) surgery
 alternative central venous catheterization route for, 1018–1023
 aortic valve replacement with, 368–369
 carbon dioxide tension in, 85–89
 cardiac output monitoring in, 770–775, 1044–1050
 cerebral oxygen saturation in, cognitive dysfunction and, 95–104
 Clostridium difficile infections after, 263–267
 coma after, 961–967
 dexamethasone for, 950–960
 in diabetes mellitus, prognosis for, 288–298
 epidural anesthesia for, internal thoracic artery endothelium effects of, 1063–1070
 glycemic variability in, 799–803
 high-risk, intra-aortic balloon pump for, 632–636
 hyperbaric oxygen preconditioning for, 908–916
 insulin during, 1086–1091
 kidney failure after, 619–624
 left stellate ganglion block for, 78–84
 lidocaine pharmacokinetics in, 931–936
 metabolic syndrome and, 282–287
 off-pump
 for anomalous origin of left coronary artery, 501–504
 anticoagulants for, 415–418
 arterial pulse waveform analysis in, 53–58
 coronary artery active perfusion system with, 311–312
 coronary artery spasm after, 1104–1105
 early extubation in, 431–436
 glycemic control in, 937–942
 glycemic variability in, 799–803
 hyperbaric oxygen preconditioning for, 908–916
 hypoalbuminemia after, 462–468
 nitroglycerin hypotension in, 140–141
 in pregnancy, 353–361
 pulse-pressure variation in, 1056–1062
 sublingual microcirculation in, 784–790
 tranexamic acid for, 26–35
 outcome of, lymphocyte count and, 975–980
 platelet function monitoring in, 42–47
 pleural effusion volume measurement in, 228–232
 postoperative pain in, gabapentin for, 824–829
 repeat, left thoracotomy approach to, 134–139
 seizures after, 288–298
 splanchnic perfusion in, 642–646
 sublingual microcirculation in, 784–790
 transfusion requirements for, 36–41
 with valve surgery, delirium in, 811–816
- Coronary artery disease
 aortic stenosis in, 368–369
 intraoperative complications with, 791–798
 ventricular septal defect development in, 1213–1216
- Coronary ostium, anomalous displacement of, 196–198
- Coronary sinus
 hematoma of, in mitral valve regurgitation, 376–377
 pericardial patch covering, 378–379
- Corticosteroids
 for cardiac surgery, outcome benefits of, 880–884
 esophageal weakness due to, 150–152
 protective effects of, in cardiac surgery, 156–165
- Cortisol, abnormal levels of, cardiovascular manifestations of, 709–710
- Cryoablation, for atrial fibrillation, for in cardiac surgery, 1030–1035
- Crystalloids, for cardiac surgery, 847–855, 1141–1153
- Cushing syndrome, cardiovascular manifestations of, 709
- Cytokines, *N*-acetylcysteine effects on, in cardiac surgery, 268–275

- Dabigatran, review of, 1208–1212
- Deep hypothermic circulatory arrest, for thromboembolic pulmonary hypertension, 746
- Defibrillation, for robotic cardiac surgery, 728
- Delirium
after cardiac surgery, 968–974
in combined CABG and valve surgery, 811–816
- Desflurane, for preconditioning, 59–65, 276–281
- Dexamethasone
for cardiac surgery, outcome benefits of, 880–884
for postoperative pain, 950–960
protective effects of, in cardiac surgery, 156–165
- Dexmedetomidine
alpha-adrenergic receptor interactions with, 856–862
for transesophageal echocardiography, 233–237
- Dextran, for cardiac surgery, 847–855
- Diabetes mellitus
CABG in, prognosis for, 288–298
glucose variability in, in CABG surgery, 799–803
glycemic control in, during CABG surgery, 937–942
- Dieulafoy lesion, bleeding from, in transesophageal echocardiography, 678–679
- Dilated cardiomyopathy, aortic valve replacement with, 1193–1199
- Dissection
aortic, in lung transplantation, 1121–1124
coronary artery, in pregnancy, 353–361
esophageal, in transesophageal echocardiography, 150–152
left atrial
after aortic valve replacement, 309–310
after mitral valve replacement, 498–500
- Diuresis, for heart failure, 9
- Doppler studies
for cardiac surgery, for brain monitoring, 1076–1085
tissue
for bariatric surgery, 943–949
for myocardial velocity measurement, 216–220
- Double-lumen tube, kinking of, 1119–1120
- Double-outlet right ventricle repair, levosimendan in, 419–424
- Ductus arteriosus, repair of
impaired lung function in, 1000–1004
levosimendan in, 419–424
- Dutch Echocardiographic Cardiac Risk Evaluation Applying Stress Echocardiography (DECREASE) trials, 170
- Ear sensor, for postoperative blood gas measurement, 243–249
- Echocardiography
for mitral valve area assessment, 486–490
transesophageal. *See* Transesophageal echocardiography
- EchoCom transesophageal echocardiography simulator, 178
- Economic issues, anesthesia preparation time, 766–769
- Editorials
Aortic Stenosis and Mitral Regurgitation: Not as Simple as It Looks, 887–888
Cardioprotection by Volatile Anesthetics: What About Noncardiac Surgery?, 899–901
The Education of Anesthesia Residents and Cardiothoracic Fellows: Can They Work Together?, 1207
The Enemy of Good. . ., 607–609
It Is a Tough "ACT" to Follow, 393–394
The Post-BART Anti-Fibrinolytic Dilemma?, 3–5
Postinfarction Ventricular Septal Defects: Surgical or Percutaneous Closure--Between a Rock and a Hard Place, 1217–1218
Scholarly Productivity, Academic Success, and Beyond, 759–760
Simulation: The Importance of "Hands-On" Learning, 209–211
- Editorials (*Continued*)
Stars of the Future, 182
Thinking Beyond the Aortic Valve: Implications of a Diseased Aorta, 370
Twenty-five Years Old and Only Getting Better, 1–2
- Education
anesthesia preparation time in, 766–769
cardiothoracic anesthesiology fellowships, 1204–1206
transesophageal echocardiography, 177–178, 212–215, 559–564
- Elderly persons, CABG surgery in, cognitive dysfunction after, 95–104
- Electroacupuncture, for poststernotomy pain, 1172
- Electrocardiography, intraoperative changes in, 791–798
- Electroencephalography, in cardiac surgery, 1076–1085
- Electrophysiology laboratory
anesthesia services for, 553–558
sedation and analgesia for, 647–659
- Embolism. *See also* Pulmonary embolism
air, after mitral valve replacement, 1219–1220
tumor, in lung, 145–147
- Enalaprilat, for hemodynamic control, 519
- Enderterectomy, carotid, method comparison for, 1024–1029
- Endobronchial blocker, for thoracic surgery, in pediatric patients, 683–686
- Endocarditis
aortic root abscess in, 192–195
aortic valve, fistula formation in, 751–752
internal jugular vein valve and, 749–750
- Endocrinopathies, cardiovascular manifestations of, 705–720
- Endoleaks, in endovascular aortic aneurysm repair, 173
- Endothelin antagonists, for pulmonary hypertension, 694
- Endotracheal cardiac output monitoring, 770–775, 830–832
- Endovascular techniques
for aortic aneurysm repair, 166–176, 316–319
pulmonary thrombendarterectomy, 183–191
- Epidural anesthesia or analgesia
for CABG surgery, internal thoracic artery endothelium effects of, 1063–1070
for poststernotomy pain, 1168–1169
for sympathectomy, for polymorphic ventricular tachycardia, 844–846
for thoracic surgery, 449–454, 817–823
for video-assisted thoracoscopic surgery, 1009–1013
- Epinephrine, for congenital heart disease surgery, 419–424
- Epoprostenol, for pulmonary hypertension, 693
- Esophagus, Barrett's, dissection of, in transesophageal echocardiography, 150–152
- EuroSCORE (European System for Cardiac Risk Evaluation)
in aortic valve replacement, 1071–1075
vs. B-type natriuretic peptide, 256–262
- EUROSTAR collaboration, for endovascular aortic aneurysm repair, 174
- Eustachian valve, persistent, 680–682
- Evidence-based medicine, for perioperative care, in cardiac surgery, 335–346
- Extracorporeal membrane oxygenation
compartment syndrome in, 836–837
for lung transplantation, 1121–1124
for pulmonary hypertension, 696
for tetralogy of Fallot repair, 313–315
- Extrapleural block, for video-assisted thoracoscopic surgery, 1009–1013
- Extravascular lung water, in acute respiratory distress syndrome, 481–485
- Extubation

Extubation (*Continued*)

- after congenital heart disease surgery, timing of, 874–879
- early
 - after cardiac surgery, sedation regimens for, 110–119
 - in off-pump CABG surgery, 431–436
- electrocardiographic changes during, 791–798

Factor VII, activated

- for cardiac surgery, stroke and, 804–810
- for pediatric cardiac surgery, 987–994

Fast-track anesthesia, for CABG surgery, immediate extubation for, 431–436

Fellowships

- accreditation for, 761–765
- cardiothoracic anesthesiology, 1204–1206

Fenoldopam mesylate, for splanchnic perfusion, in cardiac surgery, 642–646

Fentanyl, for postoperative pain, in cardiac surgery, 110–119

Fick technique, for cardiac output monitoring, 776–779

Fistulas, left atrial, in aortic root abscess, 192–195

Fluid responsiveness, after reperfusion, 780–783

Fluid therapy

- for cardiac surgery, 847–855, 1056–1062, 1141–1153
- for heart failure, 9
- hemodynamic response assessment for, 48–52
- for right ventricular insufficiency, 1051–1055

Flumazenil, vascular effects of, 72–77

FMS 2000 Rapid Infuser, malfunction of, 1091–1094

Fontan physiology, palliation of, noncardiac surgery with, 329–334

Foramen ovale, patent

- eustachian valve interfering with, 680–682
- pulmonary embolism with, 841–843

Foreign body aspiration, bronchoscopy for, 1005–1008

Gabapentin

- for postoperative pain, 824–829
- for poststernotomy pain, 1172

Gastroepiploic artery, smooth muscle of, JM-1232(-) effects on, 72–77

Gastrointestinal complications

- after CABG surgery, 637–641
- in transesophageal echocardiography, 678–679

Gastrointestinal tract, decompression of, in echocardiography, 559–561

Gelatin, for cardiac surgery, 847–855

General anesthesia, reverse Takotsubo cardiomyopathy in, 1116–1118

Glucose control

- in CABG surgery, 799–803, 937–942, 1086–1091
- in cardiac surgery, 335, 337

Growth hormone, excess of, cardiovascular manifestations of, 707–708

Heart, artificial, 1179–1192

Heart failure

- acute right ventricular, 1051–1055
- management of, 9–12
- pleural effusion in, volume of, 228–232
- in pulmonary embolism, 183–191
- right, acute respiratory distress syndrome in, 867–873

Heart rate, control of, clinical outcomes of, 509–525

Heart transplantation, pleural effusion volume measurement in, 228–232

Heartworks transesophageal echocardiography simulator, 177–178

Hematoma

- around left atrium, 1221–1222

Hematoma (*Continued*)

- coronary sinus, in mitral valve regurgitation, 376–377
- intramural, after aortic valve replacement, 309–310

Hemodynamics

- in bariatric surgery, 943–949
- in biventricular pacing, 238–242
- in CABG surgery, 1056–1062
- in cardiac output monitoring, 1044–1050
- control of, clinical outcomes and, 509–525
- in dexmedetomidine sedation, for transesophageal echocardiography, 233–237
- in passive leg raising, pulse contour analysis of, 48–52
- after reperfusion, 780–783

Hemofiltration

- aortic valve replacement in, 1193–1199
- for kidney failure, after cardiac surgery, 619–624

Hemoglobin, oxygenation of, in CABG surgery, 784–790

Hemoglobin A1c, in CABG surgery, 799–803

Heparin

- measurement of, 981–986
- in reoperations, 347–352
- thrombocytopenia induced by, 371–375

Hepatic artery, blood flow through, fenoldopam effects on, 642–646

h-index, for performance evaluation, 761–765

Hurwitz-Roberts quadricuspid aortic valve, 196–198

Hydralazine, for hemodynamic control, 519

Hydrocortisone, for cardiac surgery, outcome benefits of, 880–884

Hydroxyethyl starch, 407–414, 847–855

Hyperalgesia, after cardiac surgery, remifentanyl for, 917–925

Hyperbaric oxygen, for preconditioning, 908–916

Hyperglycemia, in CABG surgery, 799–803

Hypertension, pulmonary. *See* Pulmonary hypertension

Hyperthermia

- malignant, in cardiac surgery, 731–735
- therapeutic, for cardiac arrest, 362–367

Hyperthyroidism, cardiovascular manifestations of, 705–706

Hypertrophic cardiomyopathy

- septal myectomy for, 306–308
- treatment of, 10

Hypoalbuminemia, after off-pump CABG surgery, 462–468

Hypotension

- nitroglycerin-induced, microcirculatory response to, 140–141
- in off-pump CABG, 311–312

Hypothyroidism, cardiovascular manifestations of, 706–707

Hypoxemia, in one-lung ventilation, 1113–1115

Iloprost, for pulmonary hypertension, 694

Impedance cardiography, for cardiac output monitoring, 770–775

Infections, in cardiac surgery

- corticosteroid effects on, 156–165
- prevention of, 337–339

Inferior vena cava, renal cell carcinoma extension into, resection of, 153–155

Inflammation

- N-acetylcysteine effects on, in cardiac surgery, 268–275
- in cardiac surgery, corticosteroids for, 156–165

Inotropic agents, for hemodynamic control, 519–520

Inovasodilators, for hemodynamic control, 520

Insulin

- for CABG surgery, 1086–1091
- for glycemic control, in CABG surgery, 937–942

Intensive care, *Clostridium difficile* infections in, 263–267

Intensive Care Delirium Screening Checklist, after cardiac surgery, 968–974

- Internal jugular vein valve, 749–750
- Interventional cardiology, anesthesia services for, 553–558
- Intestinal ischemia, after CABG surgery, 637–641
- Intra-aortic balloon pump, for cardiac surgery, 632–636, 1154–1162
- Intrathecal analgesia, 1169
- Intravenous anesthesia, for noncardiac surgery, 902–907
- Intubation
 - for ductus arteriosus repair, complications of, 1000–1004
 - electrocardiographic changes during, 791–798
 - tube kinking in, 1119–1120
- Isoflurane
 - for myocardial protection, 59–65
 - for preconditioning, 1125–1140
- i-STAT device, for activated coagulation time, 395–401
- Jarvik Heart, 1180
- JM-1232(-) benzodiazepine derivative, vascular effects of, 72–77
- Journal of Cardiothoracic and Vascular Anesthesia, twenty-five years of, 1–2
- Kaolin i-STAT device, for activated coagulation time, 395–401
- Kaolin-activated coagulation time, in CABG surgery, 425–428
- Ketamine, after thoracotomy, 455–461
- Kidney
 - dysfunction or failure of
 - antifibrinolytics in, 402–406
 - aortic valve replacement in, 1193–1199
 - after cardiac surgery, 619–624
 - protection of, in endovascular aortic aneurysm repair, 172–173
- Lactate, intrathecal, in spinal cord injury, in aortic repair, 120–126
- Leakage, paravalvular, after valve replacement, 610–614
- Left ventricular assist devices
 - for heart failure, 9–10
 - insertion of, pleural effusion volume measurement in, 228–232
- Leg raising maneuver, hemodynamics of, 48–52
- Lepirudin, 371–375
- Letters to the editor
 - Acute Hemodynamic Instability in an Infant After Pulsatile Bidirectional Cavopulmonary (Glenn Shunt) Anastomosis: Mechanisms and Resolutions, 573–574
 - Amiodarone and Reperfusion Ventricular Fibrillation, 207
 - Aortic Valve Stenosis and von Willebrand Disease: A Clinical Condition Without Full Explanation, 595
 - Asleep-Awake-Asleep Technique During Carotid Endarterectomy, 580–581
 - Avulsion of a Bronchial Blocker Cuff in the Trachea When Using a Parker Flex-Tip Endotracheal Tube, 391–392
 - A Category 1 Indication for Transesophageal Echocardiography, 206–207
 - Central Venous Catheter Insertion in Patients with Transvenous Pacemaker or Defibrillator Leads, e25
 - Cerebral Oximetry-Guided Antegrade Cerebral Perfusion in Aortic Arch Surgery, 591–592
 - Clinically Relevant Outcomes and Long-term Follow-up in Cardiac Anesthesia, e19
 - CNS Monitoring: The Current Weak State of the Evidence, e15
 - Confirmation of Guidewire Placement by Ultrasound Before Tissue Dilatation Prevented Damage to the Internal Jugular Vein, 584–585
 - Decrease in Bispectral Index in a Patient Undergoing Coronary Artery Bypass Graft Surgery, e39
 - Letters to the editor (*Continued*)
 - Diagnosis of Hemodynamic Instability and Hypoxia in a Cardiac Surgery Patient: The Role of Transthoracic Echocardiography, 382–383
 - Do Volatile Anesthetics Reduce Mortality in Coronary Artery Bypass Graft Surgery?, 593–594
 - Double-Lumen Endobronchial Tube in a 6-Year-Old Child: Using Preoperative Imaging to Guide Airway Management, 602
 - The Effects of Pulsatile Cardiopulmonary Bypass on Microcirculatory Perfusion: Perspective From a Null-Result Study, e24
 - Endoscopically Assisted Transesophageal Echocardiographic Probe Insertion, e26
 - Epidurals and Outcome After Cardiac Surgery: An Example of Wishful Thinking?, e17
 - Evaluation of a Previously Undescribed Transesophageal Echocardiographic View of the Main Pulmonary Artery, e29
 - Evaluation of the Internal Jugular Vein With Transesophageal Echocardiography as a Surface Probe: A Real Alternative to Current Practice?, 574–575
 - Extracorporeal Membrane Oxygenation: A New Way to Study Toxic Drugs in Chronic Diseases, e54
 - Factors Affecting Neurocognitive Outcome After Coronary Artery Bypass Graft Surgery, e52
 - Failure of Double-Lumen Tube Positioning in a Patient With Lingual Tonsil Hypertrophy, e20
 - Hemodynamics with Postoperative Pacing, 396–397
 - Iatrogenic Aortic Dissection, e23
 - Iatrogenic Carotid Artery Pseudoaneurysm Recognized by Ultrasound, 395–396
 - Iatrogenic Mitral Valve Injury During Aortic Valve Replacement: Importance of Intraoperative Transesophageal Echocardiography, 586
 - Incomplete Left Atrial Appendage Ligation Diagnosed Intraoperatively Using Transesophageal Echocardiography Following Mitral Valve Repair, 291–394
 - An Indwelling Nasogastric Tube Interferes With Intubation Assisted by the Pentax Airway Scope, 575–576
 - Initial Anesthetic Experience During Combined Epicardial Treatment of Atrial Fibrillation, e6
 - Intraoperative Detection of a Stuck Biprosthetic Mitral Valve Leaflet Causing Severe Mitral Regurgitation, e44
 - Intraoperative Diagnosis of Acute Pulmonary Embolus by Transesophageal Echocardiogram, 603
 - Intraoperative Evaluation of Visceral Arteries by Transesophageal Echocardiography in Stanford B Thoracic Aortic Aneurysm, 589–590
 - Intraoperative Interaction Between Angiotensin-Converting Enzyme Inhibitors and Nitroglycerin in Major Surgical Cases, 605
 - Levosimendan Does Not Reduce Mortality in Cardiac Surgery and Response, 579–580
 - Lung Ultrasound and Transthoracic Echocardiography in the Differentiation Between Pericardial and Left Pleural Effusion in a Cardiac Surgery Patient, 603–605
 - Methylene Blue Diffusion After Multilevel Thoracic Paravertebral Blocks, e5
 - A Multimodal Approach to Address the Inflammation of Cardiopulmonary Bypass?, e35
 - A New Technique of Peripheral Venous Access Under Surgical Drapes in Thoracic Anesthesia, 578
 - A Novel Method of Treating Hypoxemia During One-Lung Ventilation for Thoracic Surgery, 582–583

(Continued)

- One-Lung Anesthesia Alters the Magnitude of Aortic Regurgitation Because of a Reduction in Intrathoracic Pressure, e5
- One-Ventilation for Radiofrequency Ablation of Pulmonary Lesions Out of the Surgical Area: A Secure Option, 577–578
- Overheating and Rupture of FMS 2000 Rapid Infuser, e55
- The Pharmacokinetics of Ropivacaine in Elderly Patient Undergoing Nerve Stimulator-Guided Thoracic Paravertebral Block, 592–593
- Pheochromocytoma Detected During Anesthesia Induction, e43
- Post-traumatic Stress Disorder and Neuropsychologic Impairment Among Cardiac Surgery Patients, e8
- Pro and Con Ultrasound: Are We Missing the Larger Picture?, 203
- Purpura in a Patient Receiving Vancomycin: A Leukoclastic Vasculitis?, 390–391
- Reducing Post-Cardiopulmonary Bypass Delirium: More Ketamine or Less Etomidate?, 581–582
- Reply to Dr Herzlinger, e56
- Reply to Dr Hessel, e16
- In reply to Dr Koning et al, e24
- Reply to Dr Muralidhar, e26
- Reply to Drs Brezinski et al, 585
- Reply to Drs Courtois and Rubulotta, 583
- Reply to Drs Giroti and Kiran, e53
- Reply to "Post-traumatic Stress Disorder and Neuropsychologic Impairment Among Cardiac Surgery Patients," e9
- Response: Is It Reasonable To Discourage Propofol Use in Cardiac Surgery Patients Until Strong Evidence is Provided?, 202
- Response: Spinal Analgesia with Opioids Has No Clinically Relevant Impact in Cardiac Surgery, 389–390
- Response to Howard et al, e52
- Retractions Lead to Revision of Review Article "Contemporary Fluid Management in Cardiac Anesthesia, e55
- The Role of Continuous Thoracic Paravertebral Block for Fast-Track Anesthesia After Cardiac surgery via Thoracotomy, 205–206
- Seizures After Adult Cardiac Surgery, e25
- Severe Pulmonary Artery Hypertension Caused by Hypoglycemia After Cardiac Surgery, 203–204
- Should an Anesthesiologist Be Present on Site During Cardiologic Interventions?, e51
- Simultaneous Delivery of Inhaled Prostacyclin and Milrinone Through a Double Nebulizer System, 590–591
- Spinal Analgesia in Cardiac Surgery, 388–389
- Surgical Adhesive Incise Drapes and Defibrillation During Cardiac Surgery, 387–388
- Thoracic Epidural Anesthesia and Coronary Artery Bypass Surgery: What Clinicians Have Learned from Clinical Trials, 388
- The Thoracicus Longus and Thoracodorsalis Nerve Block: A Supplemental Block for Thoracic Anesthesia Analgesia, e40
- Transcatheter Aortic Valve Implantation: Is General Anesthesia Superior to Conscious Sedation?, 576–577
- Transcatheter Aortic Valve Implantation for High-Risk Patients Does Not Reduce Mortality Compared with Predicted, 595–596
- Transesophageal Echocardiographic Short-Axis Views of the Left Atrial Appendage, e45
- Transesophageal Echocardiography in the Diagnosis of Hypoxia in the Intensive Care Unit, 583–584
- Transesophageal Echocardiography in the Diagnosis of "Spongy" Myocardium and Bicuspid Aortic Valve in an Intensive Care Unit Patient With Tachyarrhythmia, e21

(Continued)

- Transthoracic Echocardiography: Training Options for Practicing Physicians, e42
- Transthoracic Echocardiography for the Identification of Acute Aortic Regurgitation in the Intensive Care Unit, 204–205
- Transthoracic Echocardiography in the Intensive Care Unit for the Diagnosis of Right-Ventricle Endocarditis, 586–587
- Transvenous Pacemaker Lead Vegetation, 600–602
- Two-Dimensional Echocardiography and M-Mode Doppler of the Interatrial Septum for Assessment of Left Ventricular Diastolic Function, 588–589
- Ultrasound Examination for the Diagnosis of Pneumonia Associated With *Acinetobacter baumannii* in a Cardiac Surgery Patient, e38
- Unexpected Detection of Internal Jugular Vein Thrombus During Ultrasound-Guided Central Venous Cannulation, e36, e37
- Unusual Variance at the Main Carina During Bronchoscopy, 576
- Use of an Inferior Epigastric Arterial Catheter in a Patient With Difficult Vascular Access, e38
- Use of Rapid Ventricular Pacing for Facilitating Left Ventricular Assist Device Implantation, 598–600
- The Use of Transesophageal Echocardiography for Diagnosing a Malfunctioning Newly Implanted Prosthetic Mitral Valve, e57
- Levobupivacaine, for epidural anesthesia, for thoracic surgery, 449–454, 817–823
- Levosimendan
for cardiac surgery, 1154–1162
for congenital heart disease surgery, 419–424
for high-risk cardiac surgery, 632–636
- Lidocaine, pharmacokinetics of, in cardiac surgery, 931–936
- Liver
end-stage disease of, vasopressin deficiency in, 665–670
transplantation of, transfusion device malfunction during, 1092–1094
- Lobectomy
nitroglycerin hypotension in, 140–141
single-lung ventilation for, cerebral desaturation in, 127–130
- Lung
aortopulmonary collateral vessels supplying, 569–571
resection of, left atrial mass assessment during, 833–835
thromboembolic pulmonary hypertension effects on, 746–747
transplantation of
iatrogenic aortic dissection in, 1121–1124
for pulmonary hypertension, 695–696
- Lymphocyte count, preoperative, cardiac surgery outcome and, 975–980
- Malignant hyperthermia, in cardiac surgery, 731–735
- Marfan syndrome, aortic surgery in, prognosis for, 625–631
- MAX-ACT assay, in CABG surgery, 425–428
- Maze procedures, for atrial fibrillation, 1030–1035
- Medtronic ACTR II device, for activated coagulation time, 395–401
- Mesenteric artery, blood flow through, fenoldopam effects on, 642–646
- Meta-analysis, of *N*-acetylcysteine antiinflammatory action, 268–275
- Metabolic modulation therapy, for hypertrophic cardiomyopathy, 10
- Metabolic syndrome, in postoperative cognitive dysfunction, 282–287
- Methylene blue, for vasoplegia, after cardiac surgery, 736–743
- Methylprednisone, protective effects of, in cardiac surgery, 156–165
- Microcirculation, in cardiopulmonary bypass, 250–255
- Midazolam
for postoperative pain, 110–119

- Midazolam (*Continued*)
vascular effects of, 72–77
- Milrinone
for congenital heart disease surgery, 419–424
for pulmonary hypertension, 694
- Minimally invasive cardiac surgery. *See also* Endovascular techniques
anticoagulants for, 415–418
for mitral valve repair, 721–730
for mitral valve replacement, 376–377
- Mitral valve
assessment of, 863–866
cleft of, 1106–1108
deformity of, 1106–1108
endocarditis of, aortic root abscess in, 192–195
myxoma of, 889–890
parachute, 838–840
regurgitation of
after aortic valve replacement, 885–888
coronary sinus hematoma with, 376–377
repair of, 486–490
repair of
intraoperative assessment of, 221–228, 486–490
after previous CABG surgery, 134–139
robot-assisted, 721–730
replacement of
atrial intramural wall dissection after, 498–500
chordal entrapment of, 495–497
coronary sinus hematoma with, 376–377
left atrial hematoma after, 1221–1222
with mediastinal radiation therapy pathology, 1200–1203
paravalvular leaks after, 610–614
prosthesis evaluation in, 491–494
residual air in, 1219–1220
- Modelflow device, for passive leg raising response assessment, 48–52
- Morphine
for postoperative pain, in cardiac surgery, 110–119
for poststernotomy pain, 1170–1171
for preconditioning, 444–448
- Mortality
in aortic valve replacement, 1071–1075
in CABG surgery
with diabetes mellitus, 288–298
hypoalbuminemia and, 462–468
in cardiac surgery
B-type natriuretic peptide measurement and, 469–475
corticosteroid effects on, 156–165
due to kidney failure, 619–624
total lymphocyte count and, 975–980
after carotid endarterectomy, 1024–1029
in endovascular aortic aneurysm repair, 166–176
intraoperative, electrocardiography in, 791–798
- Most-Care system, for cardiac index measurement, 476–480
- Multimodal brain monitoring, in cardiac surgery, 1076–1085
- MULTIPLATE test, for heparin measurement, 981–986
- Multiple-electrode aggregometry, in CABG surgery, 42–47
- Myectomy, septal, transesophageal echocardiography for, 306–308
- Myocardial dysfunction, in sepsis and septic shock, 525–536
- Myocardial infarction
after carotid endarterectomy, 1024–1029
intraoperative, electrocardiographic detection of, 791–798
in pregnancy, CABG surgery for, 353–361
ventricular septal defect in, 1213–1216
- Myocardial ischemia
intraoperative, electrocardiographic detection of, 791–798
- Myocardial ischemia (*Continued*)
prevention of. *See* Myocardial protection; Preconditioning
volume responsiveness after, 780–783
- Myocardial protection. *See also* Preconditioning
in coronary artery bypass graft surgery, 908–916
delayed, 1125–1140
in noncardiac surgery, 902–907
in off-pump CABG surgery, isoflurane for, 53–58
- Myocardial reperfusion
after cardiac arrest, 362–367
volume responsiveness in, 780–783
- Myocardial velocity measurements, 216–220
- Myxoma, mitral valve, 889–890
- Naloxone, vascular effects of, 72–77
- Nasogastric tube, for nausea and vomiting prevention, in cardiac surgery, 105–109
- Natriuretic peptides, in outcome prediction, for cardiac surgery, 256–262, 469–475
- Nausea and vomiting, after cardiac surgery, nasogastric tube effects on, 105–109
- Near-infrared spectroscopy
for compartment syndrome assessment, 836–837
for spinal cord perfusion monitoring, in aortic aneurysm repair, 316–319
- Nellcor Durasensor, for postoperative blood gas measurement, 243–249
- Neuroendocrine tumors, cardiovascular manifestations of, 708–709
- Neurologic disorders
monitoring for, 1076–1085
prevention of, 342–343
- Nitric acid and nitric oxide synthase, in sepsis, 531
- Nitroglycerin, microcirculatory response to, 140–141
- Noncardiac surgery
cardiac protection in, 902–907
in congenital heart disease, 329–334
coronary artery laceration in, 1095–1097
- Nonsteroidal antiinflammatory drugs, for poststernotomy pain, 1172
- Norwood procedure, shunts for, 11
- Obesity, bariatric surgery for, cardiac performance during, 943–949
- Off-pump CABG. *See* Coronary artery bypass graft (CABG) surgery, off-pump
- One-lung ventilation. *See* Ventilation, one-lung
- Open Versus Endovascular Repair trial, 167
- Opioids for preconditioning, 444–448
- Outcomes. *See also* Mortality
of aortic aneurysm repair, in Marfan syndrome, 625–631
of aortic valve replacement, 1071–1075
of atrial fibrillation cryoablation, 1030–1035
of bronchoscopy, with foreign body aspiration, 1005–1008
of CABG surgery
with diabetes mellitus, 288–298
hypoalbuminemia and, 462–468
off-pump, 462–468
of cardiac surgery
with *N*-acetylcysteine, 268–275
B-type natriuretic peptide measurement and, 256–262, 469–475
corticosteroid effects on, 156–165
with metabolic syndrome, 282–287
statins effects on, 880–884
steroid effects on, 880–884
total lymphocyte count and, 975–980
of carotid endarterectomy, 1024–1029

- Outcomes. *See also* Mortality (*Continued*)
of early extubation, off-pump CABG surgery in, 431–436
of fibrinolytic therapy, 20–25
of hemodynamic control, 509–525
of kidney failure, 619–624
of robotic mitral valve repair, 728
of thromboembolic pulmonary hypertension, 744–748
of valve replacement, 610–614
- Oximetry
in CABG surgery, cognitive dysfunction and, 95–104
cerebral, for superior vena cava obstruction, 1101–1103
postoperative, ear sensor for, 243–249
- Oxygen, hyperbaric, for preconditioning, 908–916
- Oxygenation
cerebral, in cardiopulmonary bypass, 995–999
extracorporeal membrane. *See* Extracorporeal membrane oxygenation
sublingual, in CABG surgery, 784–790
- Pacing
biventricular, hemodynamics of, 238–242
redundant right ventricular lead in, 753–754
- Pancreatitis, after CABG surgery, 637–641
- Papworth BiVent tube, case studies of, 505–508
- Parachute mitral valve, deformity of, in adult, 1106–1108
- Paragangliomas, cardiovascular manifestations of, 710–712
- Parecoxib, after thoracotomy, 455–461
- Patent ductus arteriosus, repair of, impaired lung function in, 1000–1004
- Patent foramen ovale
intraoperative detection of, 145–147
pulmonary embolism with, 841–843
- Pediatric patients. *See also* Congenital heart disease
activated coagulation time measurement in, 395–401
anomalous origin of left coronary artery in, 501–504
cardiac catheterization in, remifentanyl for, 660–664
cardiac surgery in
acute respiratory distress syndrome in, 867–873
cardiac output monitoring in, 776–779
extubation timing after, 874–879
impaired lung function in, 1000–1004
levosimendan in, 419–424
mechanical valvular dysfunction in, 671–672
outcome benefits in, 880–884
postoperative bleeding control in, 987–994
renal injury in, 402–406
cefuroxime pharmacokinetics in, 425–430
foreign body aspiration in, bronchoscopy for, 1005–1008
polymorphic ventricular tachycardia in, 844–846
thoracic surgery for, endobronchial blocker for, 683–686
ventricular assist devices for, 11
- Peptic ulcer perforation, after CABG surgery, 637–641
- Percutaneous techniques, aortic valve implantation, 7–8, 437–443, 1036–1043, 1109–1112
- Performance evaluation, 761–765
- Perhexiline, for hypertrophic cardiomyopathy, 10
- Pericardial patch, covering coronary sinus, 378–379
- Pericardial tamponade, after thoracic spine surgery, 1095–1097
- Pericarditis, aortic pseudoaneurysm in, 1098–1100
- Pheochromocytomas, cardiovascular manifestations of, 710–712
- Planimetry, for mitral valve area assessment, 221–228, 486–490
- Platelet(s), functional monitoring of, in CABG surgery, 42–47
- Plethysmography, in cardiac output monitoring, 1044–1050
- Pleural effusion, volume of, transesophageal echocardiography for, 228–232
- Pneumonectomy, left atrial mass assessment in, 833–835
- Pneumoperitoneum, in bariatric surgery, cardiac performance during, 943–949
- Polar plot methodology, for cardiac output monitoring, 536–546
- Polymorphic ventricular tachycardia, in pediatric patients, 844–846
- Portal vein, blood flow through, fenoldopam effects on, 642–646
- Postoperative monitoring
of coma, 961–967
ear sensor for, 243–249
- Postoperative pain
in cardiac surgery
gabapentin for, 824–829
remifentanyl for, 917–925
sedation and analgesia regimens for, 110–119
dexamethasone for, 950–960
in sternotomy, 1163–1178
in thoracic surgery, levobupivacaine for, 817–823
in thoracotomy, 455–461
- Post-traumatic stress disorder, reverse Takotsubo cardiomyopathy with, 1116–1118
- Preconditioning
adenosine receptors in, 444–448
for coronary artery bypass graft surgery, 908–916
desflurane for, 66–71, 276–281
for endovascular aortic aneurysm repair, 171
hyperbaric oxygen for, 908–916
remifentanyl for, 926–930
- Prednisolone, protective effects of, in cardiac surgery, 156–165
- Pregnancy
in Fontan palliation, anesthesia for, 330
myocardial infarction in, CABG surgery for, 353–361
- Preload, monitoring of, in right ventricular insufficiency, 1051–1055
- Pressure half-time method, for mitral valve area assessment, 221–228, 486–490
- Pressure-recording method, for cardiac index, 476–480
- Preterm infants, ductus arteriosus repair in, impaired lung function in, 1000–1004
- Propofol
as desflurane inhibitor, 276–281
for noncardiac surgery, 902–907
for postoperative pain, in cardiac surgery, 110–119
vascular effects of, 72–77
- Prostanoids, for pulmonary hypertension, 693–694
- Protamine, for heparin reversal, 981–986
- Protection
brain, hyperbaric oxygen for, 908–916
myocardial. *See* Myocardial protection; Preconditioning
- Pseudoaneurysm, aortic, 1098–1100
- Publications, academic, 761–765
- Pulmonary artery
endarterectomy of, for pulmonary hypertension, 695
thromboendarterectomy of, 183–191
- Pulmonary artery catheter, for cardiac output monitoring, 770–775
- Pulmonary embolism
intraoperative, 1113–1115
patent foramen ovale with, 841–843
thromboendarterectomy for, 183–191
- Pulmonary hemorrhage, in tetralogy of Fallot repair, 313–315
- Pulmonary hypertension, 687–704
classification of, 688
decompensation in, 696–700
definition of, 688

- Pulmonary hypertension, 687–704 (*Continued*)
 demographics of, 688–689
 diagnosis of, 691–692
 historical view of, 687–688
 incidence of, 688–689
 pathology and pathogenesis of, 689–691
 prognosis for, 696
 thromboembolic, 744–748
 treatment of, 692–696
- Pulsatile perfusion, in cardiopulmonary bypass, 250–255
- Pulse contour analysis, for passive leg raising response assessment, 48–52
- Pulse oximetry, postoperative, ear sensor for, 243–249
- Pulse-pressure
 in CABG surgery, 1056–1062
 after reperfusion, 780–783
- Pulsus alternans, 567–568
- Quadracuspid aortic valve
 short-axis view of, 199–201
 without stenosis, 196–198
- Radiation therapy, pathology due to, valve replacement with, 1200–1203
- Recovery, quality of, dexamethasone effects on, 950–960
- Reflectance spectrophotometry, for oxygenation measurement, in CABG surgery, 784–790
- Remifentanyl
 for cardiac surgery, 917–925
 for preconditioning, 926–930
- Renal cell carcinoma
 pulmonary tumor embolism due to, 145–147
 surgery for, transesophageal echocardiography for, 153–155
- Reoperation
 left thoracotomy approach to, 134–139
 without systemic heparinization, 347–352
- Reperfusion, myocardial
 after cardiac arrest, 362–367
 volume responsiveness in, 780–783
- Research highlights from 2010, 6–15
- Resident education
 anesthesia preparation time in, 766–769
 transesophageal echocardiographic simulator for, 177–178, 212–215
- Resynchronization therapy, hemodynamics of, 238–242
- Retractions
 of De Hert, et al, paper published in *JCVA*, Volume 22: 699-705 (2008), 897
 list of, 755–757
- Retrograde autologous priming, in cardiopulmonary bypass, 995–999
- Revascularization. *See specific techniques*
- Reverse Takotsubo cardiomyopathy, 1116–1118
- Richmond Agitation-Sedation Index, after cardiac surgery, 968–974
- Right ventricular assist devices, for pulmonary hypertension, 696
- Rigid bronchoscopy, for foreign body aspiration, 1005–1008
- Robotic cardiac surgery
 carbon dioxide tension in, 85–89
 for mitral valve repair, 721–730
- Ropivacaine, after thoracotomy, 455–461
- ROTEM test, for heparin measurement, 981–986
- Safety, of transesophageal echocardiography, 559–564
- Scholarly productivity, 761–765
- Sedation
 for cardiac catheterization, 660–664
- Sedation (*Continued*)
 after cardiac surgery, 110–119
 for electrophysiology laboratory, 647–659
 for transesophageal echocardiography, 233–237
- Seizures
 in antifibrinolytic therapy, 20–25
 after cardiac surgery, 299–305
- SenTec V-Sign sensor, for postoperative blood gas measurement, 243–249
- Sepsis and septic shock
 myocardial dysfunction in, 525–536
 paravalvular leaks and, 610–614
- Septal myectomy, transesophageal echocardiography for, 306–308
- Sevoflurane
 cardioprotective effects of, in noncardiac surgery, 902–907
 for preconditioning, 1125–1140
- Shock
 in end-stage liver disease, 665–670
 septic, myocardial dysfunction in, 525–536
- Shone's anomaly, transesophageal echocardiography for, 838–840
- Sildenafil, for pulmonary hypertension, 694
- Simulation, transesophageal echocardiographic, 212–215
- Single ventricle, Norwood procedure for, 11
- Single-lung ventilation, for lobectomy, cerebral desaturation in, 127–130
- Sitaxsentan, for pulmonary hypertension, 694
- Society of Cardiovascular Anesthesiologists, echocardiographic education program of, 212–215
- Software, for mitral valve assessment, 863–866
- Somatosensory evoked potentials, in cardiac surgery, 1076–1085
- Spasm, coronary artery, vasodilator-resistant, 1104–1105
- Speckle tracking, for myocardial velocity measurement, 216–220
- Spinal cord injury
 in aortic aneurysm repair, intrathecal lactate levels in, 120–126
 monitoring for, in aortic aneurysm repair, 316–319
- Spinal surgery, coronary artery laceration after, 1095–1097
- Splanchnic perfusion, in cardiac surgery, fenoldopam mesylate for, 642–646
- Splenic rupture, after CABG surgery, 637–641
- Staphylococcus aureus infections, aortic pseudoaneurysm in, 1098–1100
- Statins
 for endovascular aortic aneurysm repair, 170–171
 outcome benefits of, 880–884
- Stellate ganglion block, for cardiac surgery, 78–84
- Stents
 for aortic aneurysm, 166–176
 for carotid endarterectomy, 1024–1029
- Sternotomy, postoperative pain in, 1163–1178
- Strain and strain rate measurement, transesophageal echocardiography for, 216–220
- Stress, reverse Takotsubo cardiomyopathy in, 1116–1118
- Stroke
 in cardiac surgery, factor VII and, 804–810
 after carotid endarterectomy, 1024–1029
- Stroke volume
 in right ventricular insufficiency, 1051–1055
 variation of, after reperfusion, 780–783
- Subclavian artery, right aberrant, occlusion of, from transesophageal echocardiography probe, 148–149
- Sublingual microcirculation, in CABG surgery, 784–790
- Superior vena cava obstruction, 1101–1103
- Supraclavicular subclavian vein catheterization, for cardiac surgery, 1018–1023

- Surfactant, for tetralogy of Fallot repair, 313–315
- Surveys, of thoracic surgery anesthesia, 1014–1017
- Sympathectomy, for polymorphic ventricular tachycardia, 844–846
- Sympathetic activation, after thoracic surgery, 817–823
- SynCardia Temporary TAH-t device, 1180–1182
- Takotsubo cardiomyopathy, reverse, 1116–1118
- Teaching. *See* Education
- Testosterone, endogenous, cardiovascular manifestations of, 712
- Tetralogy of Fallot, repair of
 - acute respiratory distress syndrome in, 867–873
 - low cardiac output after, 569–571
 - pulmonary hemorrhage control in, 313–315
- Thermodilution
 - for cardiac index, 476–480
 - for cardiac output monitoring, 770–775
 - for passive leg raising response assessment, 48–52
- Thoracic artery, internal, epidural anesthesia effects on, 1063–1070
- Thoracic impedance, for cardiac output monitoring, 776–779
- Thoracic surgery
 - anesthesia for, United Kingdom survey of, 1014–1017
 - cardiac protection in, 902–907
 - coronary artery laceration after, 1095–1097
 - endobronchial blocker for, in pediatric patients, 683–686
 - epidural anesthesia for, 449–454
 - sympathetic activation after, 817–823
- Thoracotomy
 - left, for repeat cardiac surgery, 134–139
 - postoperative pain management for, 455–461
- Thorascopic surgery, video-assisted, analgesic blocks for, 1009–1013
- Thrombocytopenia, heparin-induced, 371–375
- Thromboembolic pulmonary hypertension, 744–748
- Thromboendarterectomy, pulmonary, 183–191
- Thrombosis and thrombi. *See* Pulmonary embolism
- Thyroid dysfunction, cardiovascular manifestations of, 705–707
- Tilting-disc prosthesis, for mitral valve, 491–494
- Tissue Doppler studies, for myocardial velocity measurement, 216–220
- Tongue, microcirculation of, in CABG surgery, 784–790
- Trachea, tube kinking in, 1119–1120
- Training. *See* Education
- Tranexamic acid
 - clinical value of, 16–19
 - for off-pump CABG surgery, 26–35
 - side effects of, 20–25
 - transfusion requirements with, 36–41
- Transaxillary approach, to transcatheter aortic valve implantation, 437–443
- Transcatheter procedures, aortic valve implantation, 7–8, 437–443, 1036–1043
- Transcranial Doppler studies, in cardiac surgery, for brain monitoring, 1076–1085
- Transcutaneous electrical nerve stimulation, for poststernotomy pain, 1172
- Transesophageal echocardiography
 - aberrant right subclavian artery occlusion during, 148–149
 - advances in, 559–564
 - for aortic stenosis with coronary artery disease, 368–369
 - for bariatric surgery, 943–949
 - for bicuspid aortic valve, 891–893
 - for cardiac output monitoring, 770–775
 - for coronary sinus hematoma, 376–377
 - credentials for, 761–765
 - Transesophageal echocardiography (*Continued*)
 - dexmedetomidine sedation for, 233–237
 - dissection due to, in Barrett's esophagus, 150–152
 - gastrointestinal bleeding in, 678–679
 - for left atrial air, 1219–1220
 - for left atrial dissection, 309–310
 - for mitral prosthesis evaluation, 491–494
 - for mitral valve assessment, 221–228, 863–866
 - for mitral valve myxoma, 889–890
 - for myocardial velocity measurement, 216–220
 - of normal celiac artery, 565–566
 - for paravalvular leaks, 610–614
 - for pericardial patch covering coronary sinus, 378–379
 - for pleural effusion volume, 228–232
 - for pneumonectomy, 833–835
 - for prosthetic valve malfunction, 485–497
 - for pulmonary tumor embolism, 145–147
 - for quadricuspid aortic valve, 199–201
 - for renal cell carcinoma surgery, 153–155
 - for retained broken wire, 894–896
 - for septal myectomy, 306–308
 - for Shone's anomaly, 838–840
 - simulator for, 177–181
 - for strain and strain rate measurement, 216–220
 - three-dimensional, 561–562
 - for traumatic ventricular septal defect, 547–552
 - for valve replacement, 671–672
 - for warfarin ridge, 380–381
 - Transfemoral approach, to aortic valve replacement, 1036–1043
- Transfusions
 - for cardiac surgery, prospective observational study of, 36–41
 - devices for, malfunction of, 1091–1094
 - for off-pump CABG surgery, tranexamic acid effects on, 26–35
 - for ventricular assist devices, 10
- Transplantation
 - liver, transfusion device malfunction during, 1092–1094
 - lung
 - iatrogenic aortic dissection in, 1121–1124
 - for pulmonary hypertension, 695–696
- Transthoracic echocardiography, for mitral valve area assessment, 221–228
- Trauma, ventricular septal defect in, 547–552
- Trending, of cardiac output monitors, 536–546
- Treprostinil, for pulmonary hypertension, 693–694
- Tricuspid valve, regurgitation of
 - vs. aortic valve endocarditis, 751–752
 - cardiac output monitoring in, 830–832
 - variable severity of, 753–754
- Troponin, measurement of, in noncardiac surgery, 902–907
- Truncus arteriosus repair, levosimendan in, 419–424
- Ultrafiltration, for heart failure, 9
- Ultrasound, for splanchnic perfusion, in cardiac surgery, 642–646
- Valvular heart disease. *See also specific valves*
 - internal jugular vein valve, 749–750
 - surgery for. *See also specific valves*
 - alternative central venous catheterization route for, 1018–1023
 - atrial fibrillation in, cryoablation for, 1030–1035
 - CABG surgery with, 368–369
 - delirium in, 811–816
 - cardiac output monitoring in, 830–832
 - Clostridium difficile infections after, 263–267

Valvular heart disease. *See also specific valves (Continued)*

- coma after, 961–967
 - in complicated case, 1193–1199
 - coronary sinus hematoma with, 376–377
 - dexamethasone for, 950–960
 - intraoperative mitral valve assessment during, 221–228
 - kidney failure after, 619–624
 - left stellate ganglion block for, 78–84
 - levosimendan in, 419–424
 - lidocaine pharmacokinetics in, 931–936
 - mechanical valvular dysfunction in, 671–672
 - with mediastinal radiation therapy pathology, 1200–1203
 - mitral regurgitation after, 885–888
 - paravalvular leaks after, 610–614
 - pleural effusion volume measurement in, 228–232
 - robot-assisted, 721–730
 - transcatheter implantation, 7–8, 437–443
 - transfusion requirements for, 36–41
 - von Willebrand disease in, 131–133
- Vascular surgery, cardiac protection in, 902–907
- Vasodilation, in end-stage liver disease, 665–670
- Vasodilators
- for coronary artery spasm, 1104–1105
 - for hemodynamic control, 517–519
- Vasoplegia, after cardiac surgery, methylene blue for, 736–743
- Vasopressin, deficiency of, in liver disease, 665–670
- Vasospasm
- anesthetic agents causing, 72–77
 - coronary artery, vasodilator-resistant, 1104–1105
 - left stellate ganglion block for, 78–84
- Velocity, myocardial, measurement of, 216–220
- Vena cava, superior, obstruction of, 1101–1103
- Venous thromboembolism, prevention of, dabigatran for, 1209–1210
- Ventilation
- for aortic valve implantation, 1109–1112
 - cardiac output monitoring in, 1044–1050
 - for cardiac surgery, corticosteroid effects on, 156–165

Ventilation (*Continued*)

- for ductus arteriosus repair, complications of, 1000–1004
 - one-lung
 - case studies of, 505–508
 - in pediatric patients, 683–686
 - pulmonary embolism during, 1113–1115
 - for robotic cardiac surgery, 727
 - survey on, 1014–1017
 - tube kinking in, 1119–1120
 - for tetralogy of Fallot repair, 313–315
- Ventricle(s)
- left, outflow tract obstruction of, 673–677
 - right
 - dysfunction of, pulmonary hypertension with, 687–704
 - failure of, 1051–1055
- Ventricular assist devices
- bleeding with, 10
 - for coronary artery disease, 1213–1216
 - for heart failure, 9–10
 - insertion of, pleural effusion volume measurement in, 228–232
 - for pediatric patients, 11
- Ventricular septal defect
- closure of, levosimendan in, 419–424
 - traumatic, 547–552
- Ventricular tachycardia, polymorphic, in pediatric patients, 844–846
- Video-assisted thoracoscopic, analgesic blocks for, 1009–1013
- Vimedix transesophageal echocardiography simulator, 178–180
- Volulyte solution, 407–414
- Volume responsiveness, after reperfusion, 780–783
- Volume therapy. *See* Fluid therapy
- von Willebrand disease, acquired
- aortic stenosis and, 615–618
 - cardiac surgery with, 131–133
- Warfarin ridge artifact, 380–381
- Wire, retained, in ascending aorta, 894–896